



U.S. DEPARTMENT OF  
**ENERGY**

PNNL-20084

Prepared for the U.S. Department of Energy  
Under Contract DE-AC05-76RL01830

# Annual Hanford Seismic Report for Fiscal Year 2010

Pacific Northwest National Laboratory  
Hanford Seismic Assessment Team  
AC Rohay                      RE Clayton  
MD Sweeney                JL Devary  
DC Hartshorn

December 2010



**Pacific Northwest**  
NATIONAL LABORATORY

*Proudly Operated by **Battelle** Since 1965*

## DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor Battelle Memorial Institute, nor any of their employees, makes **any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.** Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PACIFIC NORTHWEST NATIONAL LABORATORY  
*operated by*  
BATTELLE  
*for the*  
UNITED STATES DEPARTMENT OF ENERGY  
*under Contract DE-AC05-76RL01830*

**Printed in the United States of America**

**Available to DOE and DOE contractors from the  
Office of Scientific and Technical Information,  
P.O. Box 62, Oak Ridge, TN 37831-0062;  
ph: (865) 576-8401  
fax: (865) 576-5728  
email: [reports@adonis.osti.gov](mailto:reports@adonis.osti.gov)**

**Available to the public from the National Technical Information Service,  
U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161  
ph: (800) 553-6847  
fax: (703) 605-6900  
email: [orders@ntis.fedworld.gov](mailto:orders@ntis.fedworld.gov)  
online ordering: <http://www.ntis.gov/ordering.htm>**



This document was printed on recycled paper.

(9/2003)

# **Annual Hanford Seismic Report for Fiscal Year 2010**

AC Rohay  
MD Sweeney  
DC Hartshorn

RE Clayton  
JL Devary

December 2010

Prepared for  
the U.S. Department of Energy  
under Contract DE-AC05-76RL01830

Pacific Northwest National Laboratory  
Richland, Washington 99352



## Summary

The Hanford Seismic Assessment Program (HSAP) provides an uninterrupted collection of high-quality raw and processed seismic data from the Hanford Seismic Network for the U.S. Department of Energy and its contractors. The HSAP is responsible for locating and identifying sources of seismic activity and monitoring changes in the historical pattern of seismic activity at the Hanford Site. The data are compiled, archived, and published for use by the Hanford Site for waste management, natural phenomena hazards assessments, and engineering design and construction. In addition, the HSAP works with the Hanford Site Emergency Services Organization to provide assistance in the event of a significant earthquake on the Hanford Site. The Hanford Seismic Network and the Eastern Washington Regional Network consist of 44 individual sensor sites and 15 radio relay sites maintained by the Hanford Seismic Assessment Team.

During FY 2010, the Hanford Seismic Network recorded 873 triggers on the seismometer system, which included 259 seismic events in the southeast Washington area and an additional 324 regional and teleseismic events. There were 210 events determined to be local earthquakes relevant to the Hanford Site. One hundred and fifty-five earthquakes were detected in the vicinity of Wooded Island, located about eight miles north of Richland just west of the Columbia River. With regard to the depth distribution, 173 earthquakes were located at shallow depths (less than 4 km, most likely in the Columbia River basalts), 18 earthquakes were located at intermediate depths (between 4 and 9 km, most likely in the pre-basalt sediments), and 19 earthquakes were located at depths greater than 9 km, within the crystalline basement. Geographically, 178 earthquakes were located in known swarm areas, 4 earthquakes occurred on or near a geologic structure (Saddle Mountain anticline), and 28 earthquakes were classified as random events.

The highest-magnitude events ( $3.0M_c$ ) were recorded on February 4, 2010 within the Wooded Island swarm (depth 2.4 km) and May 8, 2010 on or near the Saddle Mountain anticline (depth 3.0 km). This latter event is not considered unusual in that earthquakes have been previously recorded at this location, for example, in October 2006 (Rohay et al. 2007).

The Wooded Island events recorded this fiscal year were a continuation of the swarm events observed during fiscal year 2009 and reported in previous quarterly and annual reports (Rohay et al. 2009a, 2009b, 2009c, 2010a, 2010b, and 2010c). Most events were considered minor (coda-length magnitude [ $M_c$ ] less than 1.0) with the largest event recorded on February 4, 2010 ( $3.0M_c$ ). The estimated depths of the Wooded Island events are shallow (averaging approximately 1.5 km deep) placing the swarm within the Columbia River Basalt Group. Based upon the last two quarters (Q3 and Q4) data, activity at the Wooded Island area swarm has largely subsided. Pacific Northwest National Laboratory will continue to monitor for activity at this location.

As reported in the first quarter FY 2010 monitoring report (Rohay, et al. 2010a), strong motion accelerometer units temporarily installed directly above the Wooded Island swarm at ground surface measured peak ground accelerations approaching 15% g during FY 2009, the largest values recorded at Hanford. This corresponds to strong shaking of the ground, consistent with what people nearest the swarm's epicenters had reported. However, the swarm events duration, magnitude and distance from facilities resulted in no structural damage to facilities.

Also reported in the first quarter FY 2010 monitoring report, the USGS performed a geophysical survey (Wicks et al, 2009) using satellite interferometry that detected approximately 1 inch uplift of surface deformation within the Wooded Island swarm. Similar earthquake swarms have been recorded near this same location in 1970, 1975 and 1988 but not with SMA readings or satellite imagery. Prior to the 1970s, earthquake swarms may have occurred at this location or elsewhere in the Columbia Basin, but a dense seismic network was not in place to record those events.

The Hanford Strong Motion Accelerometer (SMA) network was triggered several times by the Wooded Island swarm events and the events located on or near the Saddle Mountain anticline. The maximum acceleration value recorded by the SMA network during fiscal year 2010 occurred February 4, 2010 (Wooded Island swarm event), approximately 2 times lower than the reportable action level for Hanford facilities (2% g) with no action required.

## Abbreviations and Acronyms

BWIP	Basalt Waste Isolation Project
CRBG	Columbia River Basalt Group
DOE	U.S. Department of Energy
ETNA	strong motion accelerometer manufactured by Kinometrics
EWRN	Eastern Washington Regional Network
FY	fiscal year
GPRS	General Packet Radio Service
GPS	Global Positioning System
HSAP	Hanford Seismic Assessment Program
HSN	Hanford Seismic Network
Mc	coda-length magnitude
ML	local magnitude
PNNL	Pacific Northwest National Laboratory
SMA	strong motion accelerometer
USGS	U.S. Geological Survey
UTC	Coordinated Universal Time
UW	University of Washington
WHC	Westinghouse Hanford Company





# Contents

Summary .....	iii
Abbreviations and Acronyms .....	v
1.0 Introduction .....	1.1
1.1 Mission.....	1.1
1.2 History of Monitoring Seismic Activity at Hanford .....	1.1
1.3 Documentation and Reports .....	1.2
2.0 Network Operations.....	2.1
2.1 Seismometer Stations .....	2.1
2.1.1 Station Maintenance .....	2.1
2.1.2 Data Acquisition.....	2.7
2.2 Strong Motion Accelerometer Stations .....	2.7
2.2.1 Location.....	2.7
2.2.2 Station Design .....	2.8
2.2.3 Strong Motion Accelerometer Operations Center.....	2.10
2.2.4 Strong Motion Operational Characteristics .....	2.10
3.0 Geology and Tectonic Analysis.....	3.1
3.1 Earthquake Stratigraphy.....	3.1
3.2 Geologic Structure Beneath the Monitored Area .....	3.1
3.3 Tectonic Pattern.....	3.4
4.0 Earthquake Catalog Description.....	4.1
4.1 Coda-Length Magnitude .....	4.1
4.2 Velocity Model.....	4.1
4.3 Quality Factors .....	4.11
5.0 Seismic Activity – FY 2010 .....	5.1
5.1 Summary .....	5.1
5.2 First Quarter FY 2010 Earthquakes.....	5.2
5.2.1 Location and Depth of Earthquakes .....	5.2
5.2.2 Major Anticlinal Ridges .....	5.3
5.2.3 Earthquake Swarm Areas .....	5.4
5.2.4 Random or Floating Events .....	5.4
5.3 Second Quarter FY 2010 Earthquakes .....	5.4
5.3.1 Location and Depth of Earthquakes .....	5.4
5.3.2 Major Anticlinal Ridges .....	5.4
5.3.3 Earthquake Swarm Areas .....	5.4
5.3.4 Random or Floating Events.....	5.5

5.4.....	Third Quarter FY 2010 Earthquakes	5.5
5.4.1	Location and Depth of Earthquakes .....	5.5
5.4.2	Major Anticlinal Ridges .....	5.5
5.4.3	Earthquake Swarm Areas .....	5.5
5.4.4	Random or Floating Events .....	5.6
5.5	Fourth Quarter FY 2010 Earthquakes .....	5.6
5.5.1	Location and Depth of Earthquakes .....	5.6
5.5.2	Major Anticlinal Ridges .....	5.6
5.5.3	Earthquake Swarm Areas .....	5.6
5.5.4	Random or Floating Events .....	5.7
6.0	Strong Motion Accelerometer Operations.....	6.1
6.1	Fourth Quarter FY 2010 Triggers of the Hanford SMA Network .....	6.1
6.2	FY 2010 Triggers of the Hanford SMA Network .....	6.1
7.0	Capabilities in the Event of a Significant Earthquake .....	7.1
8.0	References .....	8.1

## Figures

2.1	Seismometer and Strong Motion Accelerometer Stations in the Hanford Seismic Network .....	2.3
2.2	Seismometer Stations in the Eastern Washington Regional Network .....	2.5
2.3	Eltopia Station with Silage Handling and Irrigation Equipment Seen in the Distance.....	2.6
2.4	Bickleton Station with Wind Turbine Equipment Seen in the Background .....	2.6
2.5	Schematic Diagram of a Strong Motion Accelerometer Installation .....	2.9
3.1	Physical and Structural Geology of the Hanford Site, Washington.....	3.2
3.2	Geologic Cross Sections through the Columbia Basin .....	3.3
5.1	FY 2010 Earthquakes Recorded Within the Hanford Monitoring Area .....	5.3
5.2	Earthquakes Occurring in the Hanford Monitoring Area Between July 1, 2010 and September 30, 2010 .....	5.8
5.3	Three Dimensional Depiction of Earthquakes Occurring in the Hanford Monitoring Area between July 1, 2010 and September 30, 2010.....	5.9
6.1	Acceleration Records from the February 4, 2010 Wooded Island Swarm Event at the 300 Area SMA .....	6.1
6.2	Acceleration Records from the 200 East SMA May 8, 2010 Event Located On or Near the Saddle Mountain Anticline .....	6.2

## Tables

2.1	Seismometer Stations in the Hanford Seismic Network.....	2.2
2.2	Seismometer Stations in the Eastern Washington Regional Network .....	2.3
2.3	Free Field Strong Motion Accelerometer Sites.....	2.8
2.4	Instrument Parameters for the Kinemetrics ETNA System in the Hanford SMA Network .....	2.9
3.1	Thicknesses of Stratigraphic Units in the Monitoring Area .....	3.4
4.1	Local Seismic Data, October 1, 2009 – September 30, 2010 .....	4.2
4.2	Crustal Velocity Model for Eastern Washington.....	4.12
5.1	Depth Distribution of Earthquakes for FY 2010.....	5.1
5.2	Earthquake Locations for FY 2010.....	5.2



# **1.0 Introduction**

This annual report documents the locations, magnitudes, and seismic interpretations of earthquakes recorded for the Hanford monitoring area of south-central Washington during fiscal year (FY) 2010 (October 2009 through September 2010). The report provides summaries of seismic events recorded during the first three quarters of FY 2010 and a more comprehensive discussion of seismic events for the fourth quarter of FY 2010. Comprehensive discussions of seismic events recorded during the first three quarters of FY 2010 may be found in previously published quarterly reports (Rohay et al. 2010a, 2010b, 2010c).

## **1.1 Mission**

The principal mission of the Hanford Seismic Assessment Program (HSAP) is to maintain the seismometer and strong motion accelerometer (SMA) sites, report data from measured events, and provide assistance in the event of an earthquake. This mission supports the U.S. Department of Energy (DOE) and the other Hanford Site contractors in their compliance with DOE Order 420.1B, Chapter IV, Section 3.d “Seismic Detection” and DOE Order G 420.1-1, Section 4.7, “Emergency Preparedness and Emergency Communications.” DOE Order 420.1B requires facilities or sites with hazardous materials to maintain instrumentation or other means to detect and record the occurrence and severity of the seismic event. The HSAP maintains the seismic network located on and around the Hanford Site. The data collected from the seismic network can be used to support facility or site operations to protect the public, workers, and the environment from the impact of seismic events.

In addition, the HSAP provides an uninterrupted collection of high-quality raw seismic data from the Hanford Seismic Network (HSN) and the Eastern Washington Regional Network (EWRN) and provides interpretations of seismic events from the Hanford Site and the vicinity. The program locates and identifies sources of seismic activity, monitors changes in the historical pattern of seismic activity, and builds a “local” earthquake database (processed data) that is permanently archived. The focus of this report is the precise location of earthquakes proximal to or on the Hanford Site, specifically, between 46-47° north latitude and between 119-120° west longitude. Data from the EWRN and other seismic networks in the Northwest provide the HSAP with necessary regional input for the seismic hazards analysis at the Hanford Site. These seismic data are used to support Hanford Site contractors for waste management activities, natural phenomena hazards assessments, and engineering design and construction.

## **1.2 History of Monitoring Seismic Activity at Hanford**

Assessing seismic activity at the Hanford Site was initiated in 1969 by the U.S. Geological Survey (USGS) under a contract with the U.S. Atomic Energy Commission. In 1975, the University of Washington (UW) assumed responsibility for the network and subsequently expanded it. In 1979, the Basalt Waste Isolation Project (BWIP) became responsible for collecting seismic data for the Hanford Site as part of site characterization activities. Rockwell Hanford Operations, followed by Westinghouse Hanford Company (WHC), operated the local network and were the contract technical advisors for the EWRN operated and maintained by UW. Funding ended for BWIP in December 1988; the seismic program (including the UW contract) was transferred to the WHC Environmental Division. Maintenance

responsibilities for the EWRN also were assigned to WHC, who made major upgrades to EWRN sites. Effective October 1, 1996, all seismic assessment activities were transferred to the Pacific Northwest National Laboratory (PNNL).<sup>1</sup>

The Hanford SMA network was constructed during 1997, becoming operational in May 1997. It was shut down in FY 1998 due to lack of funding but became operational again in FY 1999 and has operated continuously since that time.

### **1.3 Documentation and Reports**

The HSAP issues quarterly reports of local activity, an annual catalog of earthquake activity in southeastern Washington, and special-interest bulletins on local seismic events. This includes information and special reports as requested by DOE and Hanford Site contractors. Earthquake information provided in these reports is subject to revision as new information becomes available. An archive of all seismic data from the HSAP is maintained by PNNL on computer servers.

---

<sup>1</sup> Pacific Northwest National Laboratory is operated by Battelle for the U.S. Department of Energy under Contract DE-AC05-76RL01830.

## **2.0 Network Operations**

### **2.1 Seismometer Stations**

The seismic network consists of two types of earthquake sensors—seismometers and strong motion accelerometers (SMAs). Seismometers are designed primarily to detect microearthquakes near Hanford recording seismograms that are used to determine the magnitudes and locations of seismic events. SMA stations are designed to measure ground motion and are discussed in Section 2.2.

The HSN and the EWRN consist of 44 seismometer stations. Most stations reside in remote locations and require solar panels and batteries for power. The HSN includes 26 stations (Table 2.1 and Figure 2.1), and the EWRN consists of 39 stations (Table 2.2 and Figure 2.2). Twenty-one stations are shared by both networks. Note that the HSN Bickelton (BLT) and Phinney Hill (PHIN) stations are shown on Figure 2.2.

The EWRN is used by the HSAP for two major reasons. A large earthquake located in the Pacific Northwest outside of Hanford could produce significant ground motion and damage at the Hanford Site. For example, the magnitude 7.0 event that occurred in 1872 near Chelan/Entiat or other events located in the region (e.g., eastern Cascade mountain range) could have such an effect. The EWRN would provide valuable information to help determine the impacts of such an event. Additionally, the characterization of seismicity throughout the surrounding areas, as required for the Probabilistic Seismic Hazard Analysis, supports facility safety assessments at the Hanford Site. Both the HSN and the EWRN are fully integrated within the Pacific Northwest Seismic Network managed by the University of Washington.

The HSN and EWRN networks have 54 combined data channels because the 5 three-component sites (Gable Butte, Frenchman Hills East, Cold Creek, Didier Farms, and Phinney Hill) require two additional data channels per station. The tri-axial stations record motion in the vertical, north-south horizontal, and east-west horizontal directions. The Cold Creek, Didier Farms, and Phinney Hill stations are Transportable Arrays (TA) that are broad-band seismometers with digital telemetry via cellular telephone, while the Gable Butte and Frenchman Hills East tri-axial sites have 1-Hz seismometers and analog radio telemetry. The other 39 stations are single vertical component seismometers. Fifteen radio telemetry relay sites are used by both networks to continuously transmit seismogram data to the Seismic Assessment Laboratory in the Sigma V building, Richland, Washington, for processing and archiving.

#### **2.1.1 Station Maintenance**

Site maintenance activities during the fourth quarter of FY 2010 were minimal due to the warm, dry weather typical of Central Washington summers. An item of concern was the continued industrial development near previously quiet and remote seismic locations of the Eastern Washington Regional Network (EWRN). For example, a large dairy purchased land adjacent to the Eltopia (ET3) site near Mesa, Washington (Figure 2.3). Previously used for low tillage wheat production, the land is now dedicated to intensive corn silage production to support the dairy operation.

The Bickleton (BLT) site was also impacted by the development of wind farms on land formerly dedicated to dry land wheat farming and hunting reserves. Road improvements for all-weather access for construction and maintenance of the wind towers have dramatically increased background noise levels of the seismic data for this site (see Figure 2.4).


**Table 2.1.** Seismometer Stations in the Hanford Seismic Network

Station <sup>(a)</sup>	Latitude Deg. Min. N.	Longitude Deg. Min. W.	Elevation (m)	Station Name
CCRK <sup>(b)</sup>	46°55.85	119°85.49	560	Cold Creek
DDRF <sup>(b)</sup>	46°49.12	119°05.96	270	Didier Farms
FHE <sup>(b)</sup>	46°57.11	119°29.82	455	Frenchman Hills East
PHIN <sup>(b)</sup>	45°89.52	119°92.78	270	Phinney Hill
GBB <sup>(b)</sup>	46°36.49	119°37.62	177	Gable Butte
BEN	46°31.13	119°43.02	340	Benson Ranch
BLT	45°54.91	120°10.55	659	Bickleton
BRV	46°29.12	119°59.47	920	Black Rock Valley
BVW	46°48.66	119°52.99	670	Beverly
CRF	46°49.50	119°23.22	189	Corfu
ET3	46°34.64	118°56.25	286	Eltopia Three
GBL	46°35.92	119°27.58	330	Gable Mountain
H2O	46°23.75	119°25.38	158	Water
LOC	46°43.02	119°25.85	210	Locke Island
MDW	46°36.79	119°45.66	330	Midway
MJ2	46°33.45	119°21.54	146	May Junction Two
OT3	46°40.14	119°13.98	322	Othello Three
PRO	46°12.73	119°41.15	550	Prosser
RED	46°17.92	119°26.30	366	Red Mountain
RSW	46°23.67	119°35.48	1,045	Rattlesnake Mountain
SNI	46°27.85	119°39.60	312	Snively Ranch
VT2	46°58.04	119°58.95	387	Vantage Two
WA2	46°45.32	119°33.94	244	Wahluke Slope
WIW	46°25.76	119°17.26	128	Wooded Island
WRD	46°58.20	119°08.69	375	Warden
YPT	46°02.93	118°57.73	325	Yellepit

(a) The first column is the alphanumeric seismic station designator. The latitude and longitude, elevation above sea level in meters, and the full station name follow this. The locations of the stations all are in Washington; locations were derived from the Global Positioning System (GPS).

(b) Three-component station.





**Figure 2.1.** Seismometer and Strong Motion Accelerometer Stations in the Hanford Seismic Network

**Table 2.2.** Seismometer Stations in the Eastern Washington Regional Network


Station <sup>(a)</sup>	Latitude Deg. Min. N.	Longitude Deg. Min. W.	Elevation (m)	Station Name
CCRK <sup>(b)</sup>	46°55.85	119°85.49	560	Cold Creek
DDRF <sup>(b)</sup>	46°49.12	119°05.96	270	Didier Farms
FHE <sup>(b)</sup>	46°57.11	119°29.82	455	Frenchman Hills East
PHIN <sup>(b)</sup>	45°89.52	119°92.78	270	Phinney Hill
BLT	45°54.91	120°10.55	659	Bickleton
BRV	46°29.12	119°59.47	920	Black Rock Valley

**Table 2.2.** (contd)

Station <sup>(a)</sup>	Latitude Deg. Min. N.	Longitude Deg. Min. W.	Elevation (m)	Station Name
BVW	46°48.66	119°52.99	670	Beverly
CBS	47°48.26	120°02.50	1,067	Chelan Butte South
CRF	46°49.50	119°23.22	189	Corfu
DPW	47°52.25	118°12.17	892	Davenport
DY2	47°59.11	119°46.28	890	Dyer Hill Two
ELL	46°54.58	120°33.98	789	Ellensburg
EPH	47°21.38	119°35.76	661	Ephrata
ET3	46°34.64	118°56.25	286	Eltopia Three
ETW	47°36.26	120°19.94	1,477	Entiat
GBL	46°35.92	119°27.58	330	Gable Mountain
LNO	45°52.31	118°17.11	771	Linton Mountain Oregon
LOC	46°43.02	119°25.85	210	Locke Island
MDW	46°36.79	119°45.66	330	Midway
MJ2	46°33.45	119°21.54	146	May Junction Two
MOX	46°34.64	120°17.89	501	Moxee City
NAC	46°43.99	120°49.42	728	Naches
NEL	48°04.21	120°20.41	1,500	Nelson Butte
OD2	47°23.26	118°42.58	553	Odessa Two
OT3	46°40.14	119°13.98	322	Othello Three
PAT2	45°53.03	119°45.40	259	Paterson Two
PRO	46°12.73	119°41.15	550	Prosser
RSW	46°23.67	119°35.48	1,045	Rattlesnake Mountain
SAW	47°42.10	119°24.03	701	St. Andrews
TBM	47°10.20	120°35.88	1,006	Table Mountain
TRW	46°17.32	120°32.31	723	Toppenish Ridge
TWW	47°08.29	120°52.10	1,027	Teanaway
VT2	46°58.04	119°58.95	387	Vantage Two
WA2	46°45.32	119°33.94	244	Wahluke Slope Two
WAT	47°41.92	119°57.24	821	Waterville
WIW	46°25.76	119°17.26	128	Wooded Island
WRD	46°58.20	119°08.69	375	Warden
YA2	46°31.60	120°31.80	652	Yakima Two
YPT	46°02.93	118°57.73	325	Yellepit

(a) The first column is the alphanumeric seismic station designator. The latitude and longitude, elevation above sea level in meters, and the full station name follow this. The locations of the stations all are in Washington unless otherwise indicated; locations were determined from the Global Positioning System (GPS).

(b) Three-component station.



**Figure 2.2.** Seismometer Stations in the Eastern Washington Regional Network



**Figure 2.3.** Eltopia (ET3) Station with Silage Handling and Irrigation Equipment Seen in the Distance

Hanford cleanup activities continued in FY 2010 (heavy equipment movement, vehicular traffic, well drilling, demolition and construction) and are expected to impact seismic monitoring performance for at least another 6 months.



**Figure 2.4.** Bickleton (BLT) Station with Wind Turbine Equipment Seen in the Background

### **2.1.2 Data Acquisition**

The signals from the seismometer stations are monitored for changes in signal amplitude that are expected from earthquakes. The seismic network is subdivided into spatial groupings of stations that are monitored for nearly simultaneous amplitude changes, resulting in triggering a permanent recording of the events. The groupings and associated weighting schemes are designed to allow very small seismic events to be recorded and to minimize false triggers. Events are classified as local (south-central Washington near the Hanford Site), regional (western United States and Canada), and teleseisms (from farther distances around the world). Local and regional events are usually earthquakes, but quarry and mining explosions also are recorded. Quarry and mining explosions usually can be identified from wave characteristics and the time of occurrence and may be confirmed with local government agencies and industries. Frequently, military exercises at the U.S. Army Yakima Training Center produce a series of acoustic shocks that trigger the recording system. Sonic booms and thunder also produce acoustic signals that may trigger the recording system.

The HSAP uses Earthworm, a PC-based system developed by the USGS and used by the Pacific Northwest Seismic Network at the UW, to record triggered events. One Earthworm system has been in continuous operation since January 6, 1999. A second system was installed in mid-March 1999. Both systems have been running in parallel since that time, with periodic hardware and software upgrades performed. Seismogram data from triggered events are collected on a SUN workstation (Sun Microsystems, Santa Clara, California) for assessment by HSAP staff. This information is evaluated to determine if the event is “false” (for example, due to a sonic boom) or is an earthquake or ground-surface or underground blast. Earthquake events are evaluated to determine epicenter locations, focal depths, and magnitudes (Section 4).

Although the two Earthworm systems are practically identical, slight differences in the trigger algorithms, combined with the granularity of the signal-measurement time windows, sometimes result in triggered events from one Earthworm system but not the other. These different or exclusive events are generally “false” triggers resulting from acoustical sources and not earthquakes or quarry blasts. Sometimes these exclusive events correspond to barely detectable, distant regional or teleseismic earthquakes.

## **2.2 Strong Motion Accelerometer Stations**

### **2.2.1 Location**

The Hanford SMA network consists of five free-field SMA stations (see Figure 2.1; Table 2.3) located in the 200 East, 200 West, 100 K, 400, and the 300 Areas.

The locations of SMA stations were chosen based on two criteria: 1) density of workers and 2) sites of hazardous facilities (Moore and Reidel 1996). The 200 East and 200 West Areas contain single-shell and double-shell tanks in which high-level radioactive wastes from past processing of fuel rods are stored. In addition, the Canister Storage Facility (holding encapsulated spent fuel rods) and the new Waste Treatment and Immobilization Plant being constructed are both located in the 200 East Area. The 100 K Area includes the K-West and K-East basins that are being remediated. The now inactive Fast Flux Test Facility is located in the 400 Area.

**Table 2.3.** Free Field Strong Motion Accelerometer Sites

Site	Site ID	Location	Latitude Longitude Elevation
100 K Area	H1K	South of the 100 Area fence lines	46° 38.51' 119° 35.53' 152 m
200 East Area	H2E	East of B Plant; northwest of Waste Treatment and Immobilization Plant; north of 7th Street and east of Baltimore Avenue	46° 33.58' 119° 32.00' 210 m
200 West Area	H2W	West of Plutonium Finishing Plant (PFP) and 200 West Area tree barrier	46° 33.11' 119° 38.64' 201 m
300 Area	H3A	South end of 300 Area inside fence lines (NE 1/4, SW 1/4, Sec. 11, T10N, R28E)	46° 21.83' 119° 16.55' 119 m
400 Area	H4A	500 ft from fence line on east side of facility and north of parking area)	46° 26.13' 119° 21.30' 171 m

### 2.2.2 Station Design

All free-field SMA stations consist of a four-panel solar array and two 30-gal galvanized drums that contain equipment. Each panel has a maximum 42-W output. The two 30-gal drums are set in the ground such that the base of each drum is about 1 m below the ground surface. One drum houses only the SMA; the other drum, which is connected via a sealed conduit to the SMA drum, contains the batteries. Data communication is provided by a General Packet Radio Service (GPRS) system, a continuous radio data-link to an Internet service provider. The GPRS system along with the solar power regulator is housed in a small enclosure mounted at the rear of the solar array. The enclosure serves as a junction box for all cabling that is routed through conduit inside and outside the equipment drums. The antenna for the GPRS is mounted on top of the enclosure. The enclosure permits quick access to check battery conditions and a connection directly to the RS-232 port of the SMA without removing the drum lids.


The SMA stations are three-component units consisting of vertical, north-south horizontal, and east-west horizontal seismometers manufactured by Kinemetrics, Inc., Pasadena, California, and known as the ETNA system (specifications summarized in Table 2.4). Each ETNA unit contains a digital recorder, a data storage unit, and a Global Positioning System (GPS) receiver (Figure 2.5) with the equipment housed in a watertight box.

**Table 2.4.** Instrument Parameters for the Kinometrics ETNA System in the Hanford SMA Network

Parameter	Value or Range
<b>Sensor</b>	
Type	Triaxial EpiSensor Accelerometer
Full-scale	$\pm 2 \text{ g}^{(a)}$
Frequency range	0–80 Hz
Damping	Approximately 70% critical <sup>(a)</sup>
<b>Data Acquisition</b>	
Number of channels	3
Sample rate	200 samples/sec
Resolution	18 bits
Digital output	Real-time, RS-232 output stream
<b>Seismic Trigger</b>	
Filter	0.1–12.5 Hz
Trigger level	$0.02\% \text{ g}^{(b)}$
Alarm (call-out) threshold	Not activated
Pre-event memory	10 s
Post-event time	40 s

(a) Setting is dependent on instrument calibration.

(b) See Section 2.2.4 for discussion of trigger thresholds.

**Figure 2.5.** Schematic Diagram of a Strong Motion Accelerometer Installation

The GPRS system provides the Internet address connection to access the system. Stations can be monitored from any computer with appropriate access, and data can be downloaded to a dedicated computer in the Seismic Assessment Laboratory. The data also can be downloaded directly at each site via a built-in cable connection at the enclosure in case of communication failure.



The GPS receiver is used principally to access the National Bureau of Standards timing system. The GPS receiver antenna is mounted on the enclosure at the rear of the solar array. The GPS receiver is activated internally approximately every 4 hr and checks the “location of the instrument” and the time. Any differences between the internal clock and the GPS time are recorded by the SMA. Any corrections to the internal timing are made automatically. Typically, the greatest correction recorded is approximately 4 milliseconds (ms).

### **2.2.3 Strong Motion Accelerometer Operations Center**

The combined operations, data recording, data interpretation, and maintenance facility is located in the Sigma V Building and is operated by the HSAP.

### **2.2.4 Strong Motion Operational Characteristics**

Signals from the three accelerometer channels use an 18-bit digitizer with data temporarily stored in a memory buffer. The digital sampling rate is 200 samples/s. The three channels are monitored for signals that exceed a programmable trigger threshold. When one accelerometer channel is triggered, the other channels automatically record. The nominal threshold used from 1998 to 2006 was 0.1% g (0.05% of the full-scale range of 2.0 g; g is the acceleration of gravity,  $9.8 \text{ m/s}^2$  or  $32 \text{ ft/s}^2$ ). Threshold trigger levels are set to trigger infrequently on noise sources (e.g., vehicles, sonic booms) near each site. In 2006, larger data storage capacities were installed that allowed the trigger thresholds to be reduced to 0.02% g (see Section 6). This permits the recording of ground motion data for smaller, non-damaging earthquakes that can be useful in estimating impacts of larger earthquakes. It also helps confirm the correct operation of the instruments by analyzing the smaller-amplitude triggers.

When one of the accelerometer channels exceeds the trigger threshold, the recorders save information within the data buffers. Data recording begins 10 s before the actual trigger time, continues until the trigger threshold is no longer exceeded, and ends with an additional 40 s of data. The saved files created by a triggered event are stored on memory cards to be retrieved and examined by HSAP staff.



## 3.0 Geology and Tectonic Analysis

The Hanford Site lies within the Columbia Basin, an intermontane basin between the Cascade Range and the Rocky Mountains filled with Cenozoic volcanic rocks and sediments. This basin forms the northern part of the Columbia Plateau physiographic province (Fenneman 1931) and the Columbia River flood-basalt province (Reidel et al. 1989). In the central and western parts of the Columbia Basin, the Columbia River Basalt Group (CRBG) overlies Tertiary continental sedimentary rocks and is overlain by late Tertiary and Quaternary fluvial and glaciofluvial deposits (Campbell 1989; Reidel et al. 1989, 1994; DOE 1988). In the eastern part, little or no sediment separates the basalt and underlying crystalline basement, and a thin (<10-m) veneer of eolian sediments overlies the basalt (Reidel et al. 1989, 1994).

The Columbia Basin has two structural subdivisions or subprovinces—the Yakima Fold Belt and the Palouse Slope. The Yakima Fold Belt includes the western and central parts of the Columbia Basin and is a series of anticlinal ridges and synclinal valleys with major thrust faults typically along the northern flanks (Figure 3.1) (Reidel and Fecht 1994a, 1994b). The Palouse Slope is the eastern part of the basin and is less deformed than the Yakima Fold Belt, with only a few faults and low-amplitude long-wave-length folds on an otherwise gently westward dipping paleoslope. Figure 3.2 shows north-south (B-B') and east-west (A-A') cross sections through the Columbia Basin based on surface mapping (Reidel and Fecht 1994a, 1994b), deep boreholes (Reidel et al. 1994), geophysical data (Rohay et al. 1985; DOE 1988), and magnetotelluric data obtained as part of BWIP (DOE 1988).


### 3.1 Earthquake Stratigraphy

Seismic studies at the Hanford Site have shown that the earthquake activity is related to crustal stratigraphy (large groupings of rock types) (Rohay et al. 1985; DOE 1988). The main geologic units important to earthquakes at the Hanford Site and the surrounding area are


- Miocene Columbia River Basalt Group
- Sub-basalt sediments of Paleocene, Eocene, Oligocene, and Early Miocene age
- Precambrian and Paleozoic cratonic basement
- Mesozoic accreted terranes forming the basement west of the craton margin.

### 3.2 Geologic Structure Beneath the Monitored Area

Between the late 1950s and the mid 1980s, deep boreholes were drilled for hydrocarbon exploration in the Columbia Basin. These boreholes provided accurate measurements of the physical properties of the CRBG and the pre-basalt sediments (Reidel et al. 1989, 1994), but the thickness of the sub-basalt sediments and nature of the basement are still poorly understood. Table 3.1, derived from Reidel et al. (1994), was developed for the geologic interpretation in this report. The thicknesses of these units are variable across the monitored area. Table 3.1 summarizes the approximate thickness at the borders of the monitored area.



**Figure 3.1.** Physical and Structural Geology of the Hanford Site, Washington



G01010082.2

**Figure 3.2.** Geologic Cross Sections through the Columbia Basin (Reidel et al. 1994)

**Table 3.1.** Thicknesses of Stratigraphic Units in the Monitoring Area (from Reidel et al. 1994)

Stratigraphy	North	South	East	West
Columbia River Basalt Group (includes suprabasalt sediments)	3.0 km	4.5 km	2.2 km	4.2 km
Pre-basalt sediments	3.0 km	>4.5 km	0	>6.0 km

The thickness of the basalt and the sub-basalt sediments varies as a result of different tectonic environments. The western edge of the North American craton (late Precambrian/Paleozoic continental margin and Precambrian craton) is located in the eastern portion of the monitored area (Reidel et al. 1994). The stratigraphy on the craton consists of CRBG overlying basement; the basement is continental crustal rock that underlies much of western North America. The stratigraphy west of the craton consists of 4 to 5 km of CRBG overlying up to 6 km of pre-basalt sediments. This in turn overlies accreted terranes of Mesozoic age. The area west of the craton was subsiding during the Eocene and Oligocene, accumulating great thickness of pre-CRBG sediments. Continued subsidence in this area during the Miocene resulted in thicker CRBG compared to that on the craton. Subsidence continues today but at a greatly reduced rate (Reidel et al. 1994).

### 3.3 Tectonic Pattern

Studies have concluded that earthquakes can occur in the following six different tectonic environments (earthquake sources) at the Hanford Site (Geomatrix 1996):

- **Major Geologic Structures.** Reverse/thrust faults in the CRBG associated with major anticlinal ridges such as Rattlesnake Mountain, Yakima Ridge, and Umtanum Ridge could produce some of the largest earthquakes.
- **Secondary Faults.** These faults are typically smaller (1 to 20 km in length) than the main reverse/thrust faults that occur along the major anticlinal ridges (up to 100 km in length). Secondary faults can be segment boundaries (tear faults) and small faults of any orientation that formed along with the main structure.
- **Swarm Areas.** Small geographic areas not known to contain any geologic structures produce clusters of events (swarms), usually located in synclinal valleys. These clusters consist of a series of small shocks with no outstanding principal event. Swarms occur over a period of days or months, and the events may number into the hundreds and then quit, only to start again at a later date. This differs from the sequence of foreshocks, mainshock, and trailing-off aftershocks that have the same epicenter or are associated with the same fault system. In the past, swarms were thought to occur only in the CRBG. Most swarm areas are in the basalt, but swarm events also appear to occur in all geologic layers. However, typically a swarm event at a specific time is usually restricted to one layer. Seven earthquake swarm areas are recognized in the HSN area, but this list will be updated as new swarm areas develop. The Saddle Mountains, Wooded Island, Wahluke, Coyote Rapids, and Horse Heaven Hills swarm areas are typically active at one time or another during the year (see Figure 5.1 for a map of these swarm areas). The other earthquake swarm areas are active less frequently.

- **Entire Columbia Basin.** The entire basin, including the Hanford Site, could produce a “floating” earthquake. A floating earthquake is one that, for seismic design purposes, can happen anywhere in a tectonic province and is not associated with any known geologic structure. Seismic interpretation classifies it as a random event for purposes of seismic design and vibratory ground motion studies.
- **Basement Source Structures.** Studies (Geomatrix 1996) suggest that major earthquakes can originate in tectonic structures in the basement. Because little is known about geologic structures in the basement beneath the Hanford Site, earthquakes cannot be directly tied to a mapped fault. Earthquakes occurring in the basement without known sources are treated as random events.
- **Cascadia Subduction Zone.** This source has been postulated to be capable of producing a magnitude 9 earthquake. Because this source is along the western boundary of Washington State and outside the HSN, the Cascadia subduction zone is not an earthquake source that is monitored at the Hanford Site, so subduction zone earthquakes are not reported here. Because any earthquake along the Cascadia subduction zone can have a significant impact on the Hanford Site or can be felt like the February 2001 Nisqually earthquake, UW monitors and reports on this earthquake source for the DOE. Ground motion from any moderate or larger Cascadia subduction zone earthquake is detected by Hanford SMAs and reported (see Section 5).



## 4.0 Earthquake Catalog Description

An interactive program called XPED, developed at the University of Washington, was used to determine earthquake locations and magnitudes. This program reads seismogram data recorded by the Earthworm system and lets the user measure arrival times and durations from earthquakes. Arrival and duration times are used as input to the hypocenter routine within XPED to estimate locations and magnitudes of the seismic events. XPED results for local earthquakes (46-47° north latitude, 119-120° west longitude) are reported in Table 4.1. Other seismic events located in southeastern Washington, the Pacific Northwest, or outside the region also are evaluated, with results stored on the computer system; these results are not reported in this document. These other results sometimes are used as a check to confirm that the HSN is functioning properly (e.g., quality checks on data recording).

### 4.1 Coda-Length Magnitude

Coda-length magnitude ( $M_c$ ), an estimate of local magnitude ( $M_L$ ) (Richter 1958), is calculated using a relationship developed for Washington State by Crosson (1972):

$$M_c = 2.82 \log (D) - 2.46$$

where  $D$  is the duration of the observed event. Many of the earthquakes have magnitude determinations that are very small ( $M_c < 0$ ) and highly uncertain. In Section 4, we define earthquakes as “minor” with magnitudes ( $M_c$ ) smaller than 1.0. Coda-length magnitudes for events classified as explosions are not reported because they are biased by a prominent surface wave that extends the apparent duration in a way inconsistent with coda-length measurement.

### 4.2 Velocity Model

XPED uses the crustal velocity model for eastern Washington given in Table 4.2. The model does not include a surficial layer for the Hanford or Ringold formations because most seismometer stations are sited on basalt. The crustal velocity model extends 38 km deep (to the mantle) and consists of six layers, each with uniform seismic velocity. The crustal velocity model was developed using available geologic information and calibrated from seismic data recorded from accurately located earthquake and blast events in eastern Washington. Time corrections (delays) are incorporated into the velocity model to account for significant deviations in station elevations or stations situated on sedimentary layers. Station delays also are determined empirically from accurately located earthquakes and blast events in the region.

**Table 4.1.** Local Seismic Data, October 1, 2009 – September 30, 2010

Type	Date	Time	Latitude	Longitude	Depth	Mag	NS/NP	Gap	Dmin	RMS	Q	Location
	09/10/01	03:27:57.51	46N24.03	119W17.35	1.78	0.2	7/09	141	1	0.04	AC	Wooded Island Swarm
	09/10/01	13:24:27.69	46N24.06	119W17.67	0.28	-0.3	5/06	126	2	0.04	AD	Wooded Island Swarm
	09/10/02	17:55 28.84	46N13.11	119W28.28	9.32	0.2	11/17	216	9	0.09	AD	16 km WSW of Richland
	09/10/04	14:09:03.03	46N24.93	119W16.58	0.97	0.5	9/9	114	1	0.04	AB	Wooded Island Swarm
	09/10/04	14:26:49.49	46N24.56	119W16.13	0.43	1.2	18/18	88	2	0.10	AA	Wooded Island Swarm
	09/10/04	14:27:35.35	46N24.77	119W16.53	1.29	-0.1	7/7	116	2	0.07	AB	Wooded Island Swarm
	09/10/04	14:28:26.26	46N24.90	119W16.51	0.95	0.6	9/9	116	1	0.08	AB	Wooded Island Swarm
	09/10/04	14:37:44.44	46N24.89	119W16.49	1.85	0.2	7/7	116	1	0.02	AB	Wooded Island Swarm
	09/10/04	15:29 32.01	46N15.15	119W27.71	8.77	0.5	13/13	282	5	0.06	AD	14 km WSW of Richland
	09/10/05	16:03:20.20	46N24.91	119W16.16	2.10	0.1	9/9	125	2	0.05	AB	Wooded Island Swarm
	09/10/06	00:12 06.08	46N31.95	119W42.46	18.42	0.1	5/8	130	8	0.02	AD	6 km WSW of 200 West
	09/10/06	07:25:39.39	46N24.37	119W16.03	0.41	-0.3	7/7	135	2	0.14	AB	Wooded Island Swarm
	09/10/06	07:39:24.24	46N24.39	119W15.85	0.42	-0.3	7/7	141	2	0.11	AC	Wooded Island Swarm
	09/10/06	08:05:31.31	46N23.52	119W16.89	0.04	0.6	17/17	91	0	0.06	AB	Wooded Island Swarm
	09/10/06	09:06:17.17	46N23.50	119W16.77	0.27	0.3	11/11	88	0	0.02	AA	Wooded Island Swarm
	09/10/06	12:32:12.12	46N25.00	119W16.43	1.39	0.0	10/13	117	1	0.05	AB	Wooded Island Swarm
	09/10/06	12:34:21.21	46N24.95	119W16.58	1.45	0.2	13/13	114	1	0.04	AB	Wooded Island Swarm
	09/10/06	12:40:27.27	46N24.99	119W16.49	1.83	0.6	14/14	116	1	0.03	AB	Wooded Island Swarm
	09/10/06	23:27:12.12	46N24.91	119W16.57	1.46	1.1	22/22	86	1	0.09	AA	Wooded Island Swarm
	09/10/06	23:44:57.57	46N24.97	119W16.54	1.58	0.2	8/11	115	1	0.06	AB	Wooded Island Swarm
	09/10/07	00:37:41.41	46N25.12	119W16.59	1.96	0.0	8/8	113	1	0.05	AB	Wooded Island Swarm
	09/10/09	01:37:54.54	46N24.57	119W15.89	0.41	0.4	14/15	89	2	0.08	AA	Wooded Island Swarm
	09/10/09	01:58:18.18	46N25.04	119W16.10	1.72	-0.1	9/9	125	1	0.03	AB	Wooded Island Swarm
	09/10/09	04:27:40.40	46N24.49	119W15.83	0.43	0.2	10/10	139	2	0.13	AC	Wooded Island Swarm
	09/10/09	10:53:39.39	46N24.51	119W15.79	0.41	0.2	12/12	140	2	0.09	AC	Wooded Island Swarm
	09/10/10	07:54 40.07	46N41.79	119W38.89	15.72	0.0	12/17	79	9	0.06	AA	7 km NNW of 100-K



**Table 4.1.** (contd)

Type	Date	Time	Latitude	Longitude	Depth	Mag	NS/NP	Gap	Dmin	RMS	Q	Location
	09/10/11	02:22:09.09	46N24.95	119W16.04	1.81	0.6	10/10	127	2	0.05	AB	Wooded Island Swarm
	09/10/11	05:56:15.15	46N24.42	119W15.87	2.52	-0.1	7/7	139	2	0.07	AC	Wooded Island Swarm
	09/10/13	02:21:34.34	46N23.37	119W16.08	0.02	-0.3	6/6	250	0	0.06	AD	Wooded Island Swarm
	09/10/14	01:52:30.30	46N24.03	119W16.30	0.86	-0.5	5/5	209	1	0.11	AD	Wooded Island Swarm
	09/10/14	04:26:40.40	46N24.63	119W16.07	1.60	0.1	9/9	130	2	0.12	AB	Wooded Island Swarm
	09/10/14	06:12:57.57	46N24.88	119W16.33	0.43	-0.1	9/9	120	2	0.06	AB	Wooded Island Swarm
	09/10/15	15:28:44.44	46N24.89	119W15.90	1.60	0.5	10/11	131	2	0.05	AB	Wooded Island Swarm
	09/10/15	17:18:38.38	46N24.87	119W16.14	1.16	0.4	7/7	125	2	0.10	AB	Wooded Island Swarm
	09/10/16	07:43 47.42	46N29.89	119W39.59	8.64	-0.5	8/11	128	3	0.06	AB	7 km SSW of 200 West
	09/10/16	07:44 24.91	46N29.77	119W39.48	7.89	-1.1	6/8	129	3	0.10	AC	7 km SSW of 200 West
	09/10/16	07:49 05.10	46N29.67	119W39.35	7.77	-1.1	7/10	130	3	0.08	AB	7 km S of 200 West
	09/10/16	10:05 08.55	46N29.87	119W39.61	8.77	-0.3	7/9	128	3	0.06	AB	7 km SSW of 200 West
	09/10/16	12:46:36.36	46N24.95	119W16.12	1.50	0.8	13/15	87	2	0.05	AA	Wooded Island Swarm
	09/10/16	13:42:29.29	46N23.55	119W16.90	1.61	1.4	17/19	82	0	0.06	AA	Wooded Island Swarm
	09/10/17	03:53:34.34	46N25.00	119W16.53	1.44	0.0	7/10	115	1	0.04	AB	Wooded Island Swarm
	09/10/17	17:58:29.29	46N24.88	119W16.11	2.00	-0.1	6/6	230	2	0.03	AD	Wooded Island Swarm
	09/10/18	05:49:48.48	46N24.82	119W16.38	0.42	0.1	7/7	215	2	0.10	AD	Wooded Island Swarm
	09/10/18	09:09:00.00	46N24.87	119W16.08	1.76	0.5	11/13	127	2	0.03	AB	Wooded Island Swarm
	09/10/18	10:39:44.44	46N24.88	119W16.14	2.03	0.2	8/8	125	2	0.03	AB	Wooded Island Swarm
	09/10/18	12:36:29.29	46N24.93	119W16.18	2.09	0.1	8/8	124	2	0.05	AB	Wooded Island Swarm
	09/10/20	06:02:42.42	46N24.48	119W15.91	1.81	0.1	7/10	137	2	0.07	AC	Wooded Island Swarm
	09/10/21	12:02:58.58	46N25.02	119W16.02	1.94	0.5	13/13	127	2	0.04	AB	Wooded Island Swarm
	09/10/21	12:03:33.33	46N24.90	119W15.94	3.11	0.1	8/8	130	2	0.10	AB	Wooded Island Swarm
	09/10/21	12:03:51.51	46N24.81	119W15.89	2.76	-0.5	4/4	132	2	0.00	AD	Wooded Island Swarm
	09/10/22	06:59:19.19	46N24.50	119W16.01	1.60	-0.1	7/7	134	2	0.05	AB	Wooded Island Swarm
	09/10/24	01:54:42.42	46N24.93	119W15.95	2.06	0.2	9/9	129	2	0.04	AB	Wooded Island Swarm
	09/10/26	13:12:53.53	46N24.70	119W15.84	2.05	0.5	14/14	135	2	0.04	AB	Wooded Island Swarm

**Table 4.1.** (contd)

Type	Date	Time	Latitude	Longitude	Depth	Mag	NS/NP	Gap	Dmin	RMS	Q	Location
	09/10/26	22:38:14.14	46N24.97	119W15.99	1.68	0.8	8/11	128	2	0.02	AB	Wooded Island Swarm
	09/10/27	11:19:48.48	46N24.63	119W15.78	1.46	0.2	11/11	138	2	0.04	AC	Wooded Island Swarm
	09/10/27	14:32:39.39	46N24.97	119W15.94	1.76	0.1	8/8	129	2	0.02	AB	Wooded Island Swarm
	09/11/05	02:10:39.39	46N24.99	119W16.10	1.93	0.7	16/17	125	2	0.05	AB	Wooded Island Swarm
	09/11/05	02:21:35.35	46N24.91	119W16.21	1.25	0.4	10/10	123	2	0.03	AB	Wooded Island Swarm
	09/11/05	02:31:45.45	46N24.95	119W16.08	2.16	0.1	9/9	126	2	0.07	AB	Wooded Island Swarm
	09/11/10	07:32:42.42	46N24.92	119W15.63	0.04	0.4	8/9	137	2	0.05	AC	Wooded Island Swarm
	09/11/10	08:47:54.54	46N24.90	119W15.64	0.18	0.2	7/8	137	2	0.05	AC	Wooded Island Swarm
	09/11/14	06:20 50.58	46N19.50	119W46.87	20.33	0.6	18/19	154	14	0.09	AC	13 km N of Prosser
	09/11/14	06:22 12.75	46N19.94	119W47.00	17.55	-0.2	11/12	271	15	0.06	AD	14 km N of Prosser
	09/11/14	06:23 11.36	46N19.30	119W47.72	20.01	0.9	15/15	137	14	0.12	AC	13 km N of Prosser
	09/11/14	13:58:38.38	46N25.05	119W15.63	0.03	-0.3	7/7	135	2	0.09	AB	Wooded Island Swarm
	09/11/14	17:19 15.18	46N10.44	119W34.02	17.41	0.8	15/16	168	10	0.07	AC	16 km ESE of Prosser
	09/11/14	19:28 47.08	46N10.73	119W34.08	16.86	0.8	14/15	248	9	0.05	AD	16 km ESE of Prosser
	09/11/14	22:38:03.03	46N24.65	119W16.06	1.37	-0.1	9/9	130	2	0.01	AB	Wooded Island Swarm
X	09/11/24	21:31 11.47	46N08.14	119W45.68	0.04		14/14	124	10	0.09	AC	8 km S of Prosser
	09/11/26	22:01 27.34	46N15.09	119W22.68	19.68	0.8	18/18	134	6	0.10	AB	8 km WSW of Richland
	09/11/27	19:41 49.91	46N36.18	119W51.02	7.41	0.2	10/10	169	4	0.04	AC	17 km WNW of 200 West
	09/11/28	14:29:25.25	46N24.87	119W16.13	1.94	1.5	22/22	87	2	0.09	AA	Wooded Island Swarm
	09/11/28	14:30:01.01	46N24.87	119W15.67	0.03	-0.8	4/5	137	2	0.02	AD	Wooded Island Swarm
	09/12/01	13:47:43.43	46N24.99	119W16.14	1.98	0.5	11/15	124	2	0.05	AB	Wooded Island Swarm
	09/12/05	03:46:15.15	46N25.11	119W15.68	1.74	-0.3	7/8	255	2	0.09	AD	Wooded Island Swarm
	09/12/06	00:45 10.88	46N35.92	119W51.27	7.45	0.1	8/12	174	4	0.04	AC	17 km WNW of 200 West
	09/12/15	16:17:01.01	46N24.67	119W15.65	2.32	0.0	9/11	141	2	0.09	AC	Wooded Island Swarm
X	09/12/15	21:54 29.29	46N14.46	119W43.92	0.33		9/9	143	4	0.08	AC	5 km NE of Prosser
	09/12/19	13:31:51.51	46N22.88	119W16.55	0.16	-0.1	9/10	179	0	0.09	AC	Wooded Island Swarm
	09/12/22	09:22:33.33	46N24.94	119W15.99	2.12	2.5	22/22	87	2	0.08	AA	Wooded Island Swarm

**Table 4.1.** (contd)

Type	Date	Time	Latitude	Longitude	Depth	Mag	NS/NP	Gap	Dmin	RMS	Q	Location
	09/12/22	15:25:12.12	46N24.75	119W16.01	2.21	0.2	10/11	130	2	0.12	AB	Wooded Island Swarm
	09/12/23	17:11:25.25	46N25.03	119W14.86	0.40	0.1	6/6	278	3	0.05	AD	Wooded Island Swarm
	09/12/23	17:38:39.39	46N25.16	119W16.14	2.08	0.2	7/7	123	1	0.04	AB	Wooded Island Swarm
	10/01/06	18:55:14.20	46N29.72	119W35.60	17.20	0.1	7/10	115	6	0.07	AB	8 km SSE of 200 West
	10/01/14	03:50:46.93	46N25.43	119W16.19	2.40	-0.1	5/06	157	1	0.14	BD	Wooded Island Swarm
	10/01/31	14:46:43.29	46N25.13	119W16.08	1.71	0.1	6/09	160	1	0.11	AC	Wooded Island Swarm
	10/02/02	06:26:07.10	46N42.94	119W30.05	4.05	0.9	12/17	71	5	0.10	AB	11 km NE of 100-K
	10/02/03	14:00:09.00	46N24.08	119W17.81	2.22	1.0	15/17	85	3	0.06	AA	Wooded Island Swarm
	10/02/03	15:45:49.55	46N24.57	119W18.15	2.45	-0.1	5/06	155	2	0.09	AD	Wooded Island Swarm
	10/02/04	03:58:01.29	46N24.35	119W18.24	1.11	-0.1	5/07	194	2	0.02	AD	Wooded Island Swarm
	10/02/04	05:54:36.13	46N23.83	119W17.64	2.38	3.0	20/22	86	3	0.06	AA	Wooded Island Swarm
	10/02/04	06:20:44.20	46N24.34	119W18.02	1.54	0.6	7/09	158	2	0.06	AC	Wooded Island Swarm
	10/02/04	19:42:53.43	46N24.25	119W18.05	0.40	0.5	5/06	201	2	0.06	AD	Wooded Island Swarm
	10/02/04	21:25:29.70	46N24.03	119W16.67	2.85	-0.1	3/04	240	3	0.00	AD	Wooded Island Swarm
	10/02/06	15:04:28.16	46N24.42	119W17.86	2.45	0.2	5/08	158	2	0.05	AD	Wooded Island Swarm
	10/02/06	15:16:58.73	46N24.39	119W17.93	2.38	0.6	7/09	158	2	0.03	AC	Wooded Island Swarm
	10/02/06	15:41:39.71	46N24.36	119W17.94	2.08	-0.1	4/05	202	2	0.00	AD	Wooded Island Swarm
	10/02/06	20:50:19.41	46N24.18	119W17.85	2.20	0.8	8/10	160	3	0.04	AC	Wooded Island Swarm
	10/02/08	09:49:42.10	46N24.00	119W17.58	2.22	0.9	9/10	163	3	0.02	AC	Wooded Island Swarm
	10/02/08	09:51:16.04	46N24.40	119W17.84	2.66	0.2	5/06	158	2	0.02	AD	Wooded Island Swarm
	10/02/08	09:54:40.69	46N24.14	119W17.89	0.35	0.1	5/06	160	3	0.02	AD	Wooded Island Swarm
	10/02/08	10:11:42.88	46N24.13	119W17.75	2.27	0.5	5/07	211	3	0.08	AD	Wooded Island Swarm
	10/02/08	11:13:47.35	46N24.01	119W17.78	1.78	0.2	5/07	162	3	0.01	AD	Wooded Island Swarm
	10/02/08	11:52:02.06	46N24.12	119W17.99	1.62	0.2	4/07	160	3	0.02	AD	Wooded Island Swarm
	10/02/08	12:21:31.35	46N24.09	119W17.90	0.19	0.8	10/14	161	3	0.08	AC	Wooded Island Swarm
	10/02/08	12:26:04.74	46N24.82	119W16.20	2.01	0.5	5/07	162	2	0.06	AD	Wooded Island Swarm
	10/02/08	13:42:12.78	46N23.92	119W17.61	0.45	0.5	6/09	164	3	0.08	AC	Wooded Island Swarm

**Table 4.1.** (contd)

Type	Date	Time	Latitude	Longitude	Depth	Mag	NS/NP	Gap	Dmin	RMS	Q	Location
	10/02/12	09:28:06.80	46N24.03	119W17.74	1.91	-0.3	6/07	104	2	0.12	AC	Wooded Island Swarm
	10/02/12	20:20:20.26	46N08.32	119W10.24	0.41	1.5	17/18	261	18	0.13	AD	8 km SSW of Kennewick
	10/02/14	08:09:45.71	46N23.94	119W17.78	1.24	1.4	14/18	77	2	0.05	AA	Wooded Island Swarm
	10/02/14	08:38:20.46	46N23.92	119W17.60	1.66	0.8	12/16	92	1	0.06	AB	Wooded Island Swarm
	10/02/14	08:38:29.10	46N24.11	119W17.48	1.48	0.7	5/06	137	2	0.05	AD	Wooded Island Swarm
	10/02/14	09:07:08.39	46N23.95	119W17.63	1.55	0.6	6/09	99	2	0.02	AC	Wooded Island Swarm
	10/02/14	09:37:42.19	46N24.01	119W17.61	1.39	0.9	13/16	88	2	0.07	AA	Wooded Island Swarm
	10/02/14	09:40:13.92	46N24.07	119W17.62	1.20	0.1	7/11	100	2	0.09	AB	Wooded Island Swarm
	10/02/14	23:51:51.07	46N24.05	119W17.76	1.01	0.2	7/10	103	2	0.08	AB	Wooded Island Swarm
	10/02/14	23:51:51.07	46N24.05	119W17.76	1.01	0.2	7/10	103	2	0.08	AB	Wooded Island Swarm
	10/02/16	04:03:28.93	46N23.94	119W17.93	1.45	0.6	8/11	97	2	0.02	AB	Wooded Island Swarm
	10/02/16	04:11:35.39	46N23.88	119W17.72	0.96	0.8	13/15	96	2	0.06	AB	Wooded Island Swarm
	10/02/16	04:11:56.02	46N24.04	119W16.87	2.04	0.1	4/05	171	1	0.04	AD	Wooded Island Swarm
	10/02/16	04:12:15.32	46N23.95	119W17.69	1.83	0.5	9/10	92	2	0.03	AB	Wooded Island Swarm
	10/02/16	04:26:27.08	46N24.04	119W17.86	1.70	0.7	8/10	92	2	0.04	AB	Wooded Island Swarm
	10/02/16	04:43:34.09	46N23.63	119W17.62	0.49	0.1	6/07	108	1	0.05	AC	Wooded Island Swarm
	10/02/16	13:22:48.53	46N24.48	119W18.31	3.00	-0.3	6/09	125	2	0.05	AC	Wooded Island Swarm
	10/02/19	05:48:39.10	46N23.96	119W17.54	1.51	0.6	12/14	88	1	0.03	AA	Wooded Island Swarm
	10/02/23	17:29:26.49	46N24.00	119W17.31	1.23	0.2	6/11	93	1	0.06	AC	Wooded Island Swarm
	10/02/24	00:05:51.74	46N23.86	119W17.63	1.77	0.8	8/12	95	1	0.04	AB	Wooded Island Swarm
	10/02/24	00:06:21.52	46N24.33	119W17.22	3.22	-0.8	4/05	156	2	0.00	AD	Wooded Island Swarm
	10/02/24	02:57:43.79	46N23.93	119W17.31	1.67	0.4	7/10	92	1	0.09	AB	Wooded Island Swarm
	10/02/24	02:58:44.09	46N24.06	119W16.98	1.15	-0.5	5/08	164	1	0.04	AD	Wooded Island Swarm
	10/02/24	03:05:17.55	46N23.95	119W17.28	1.77	0.6	10/13	83	1	0.07	AA	Wooded Island Swarm
	10/02/24	03:24:10.92	46N23.94	119W17.35	1.73	-0.4	6/08	94	1	0.05	AC	Wooded Island Swarm
	10/02/24	03:29:56.95	46N23.97	119W17.46	1.58	0.4	9/11	86	1	0.06	AA	Wooded Island Swarm
	10/02/24	03:30:35.47	46N24.27	119W17.05	3.45	-0.8	4/06	165	2	0.04	AD	Wooded Island Swarm

**Table 4.1.** (contd)

Type	Date	Time	Latitude	Longitude	Depth	Mag	NS/NP	Gap	Dmin	RMS	Q	Location
	10/02/24	04:01:08.02	46N24.00	119W17.82	1.13	1.8	19/21	77	2	0.05	AA	Wooded Island Swarm
	10/02/24	04:01:28.57	46N24.16	119W16.82	0.41	0.5	4/06	176	1	0.07	AD	Wooded Island Swarm
	10/02/24	04:14:51.78	46N23.92	119W17.70	1.47	1.2	13/15	77	2	0.08	AA	Wooded Island Swarm
	10/02/24	04:15:49.95	46N23.93	119W17.88	0.34	-0.5	5/07	113	2	0.11	AD	Wooded Island Swarm
	10/02/24	04:17:08.62	46N23.91	119W17.90	0.42	-0.1	5/07	111	2	0.06	AD	Wooded Island Swarm
	10/02/24	04:17:17.37	46N24.14	119W17.37	0.26	-0.1	3/04	143	2	0.06	AD	Wooded Island Swarm
	10/02/24	04:17:46.27	46N24.02	119W17.06	1.64	-0.3	5/06	157	1	0.10	BD	Wooded Island Swarm
	10/02/24	04:18:20.44	46N23.88	119W17.72	1.50	0.9	14/16	77	2	0.06	AA	Wooded Island Swarm
	10/02/24	04:20:29.63	46N24.22	119W15.83	1.57	-0.5	4/06	236	2	0.09	BD	Wooded Island Swarm
	10/02/24	06:01:21.39	46N24.01	119W17.60	1.66	0.3	11/13	88	2	0.04	AA	Wooded Island Swarm
	10/02/24	06:03:23.03	46N24.03	119W17.53	1.83	0.4	7/09	132	2	0.02	AB	Wooded Island Swarm
	10/02/24	06:07:58.60	46N23.92	119W17.84	1.95	0.6	7/10	114	2	0.06	AB	Wooded Island Swarm
	10/02/24	06:30:01.70	46N23.86	119W17.39	1.59	0.4	7/11	89	1	0.04	AB	Wooded Island Swarm
	10/02/24	06:30:27.13	46N23.72	119W17.00	1.67	-0.5	4/06	149	1	0.09	BD	Wooded Island Swarm
	10/02/24	06:43:56.82	46N24.04	119W17.84	0.82	0.1	5/07	118	2	0.02	AD	Wooded Island Swarm
	10/02/24	11:23:15.55	46N24.09	119W16.89	0.42	-0.4	6/07	102	1	0.06	AC	Wooded Island Swarm
	10/02/24	12:13:43.44	46N23.86	119W17.96	0.40	0.9	8/10	101	2	0.04	AB	Wooded Island Swarm
	10/02/24	12:49:32.54	46N24.07	119W17.61	1.29	0.2	6/08	101	2	0.04	AC	Wooded Island Swarm
	10/02/24	13:43:59.99	46N24.07	119W17.28	0.41	-0.1	5/08	146	1	0.14	AD	Wooded Island Swarm
	10/02/24	16:30:56.04	46N24.10	119W17.36	1.70	-0.1	5/07	96	1	0.07	AD	Wooded Island Swarm
	10/02/25	14:06:00.27	46N24.15	119W17.69	3.14	0.5	7/10	84	2	0.13	AB	Wooded Island Swarm
	10/02/27	04:05:59.86	46N24.81	119W15.98	1.94	-0.1	4/07	130	2	0.03	AD	Wooded Island Swarm
	10/03/06	11:54:51.40	46N24.06	119W17.67	1.64	0.4	9/12	87	2	0.02	AA	Wooded Island Swarm
	10/03/06	23:08:32.11	46N28.00	119W31.65	14.33	-0.2	8/10	94	8	0.04	AB	10 km S of 200 East
	10/03/07	04:53:18.26	46N43.28	119W29.50	4.57	-0.2	6/10	91	4	0.07	AC	12 km NE of 100-K
	10/03/10	00:32:18.12	46N24.19	119W17.79	1.15	0.1	7/08	85	2	0.06	AB	Wooded Island Swarm
	10/03/13	01:46:35.14	46N23.93	119W17.15	1.60	0.2	7/11	87	1	0.07	AB	Wooded Island Swarm

**Table 4.1.** (contd)

Type	Date	Time	Latitude	Longitude	Depth	Mag	NS/NP	Gap	Dmin	RMS	Q	Location
	10/03/24	23:51:11.33	46N35.90	119W25.12	13.78	-0.3	9/14	138	3	0.05	AC	10 km ENE of 200 East
	10/03/25	07:59:06.68	46N23.97	119W17.61	2.31	0.2	7/09	90	2	0.04	AB	Wooded Island Swarm
	10/03/25	15:07:07.02	46N50.03	119W28.42	13.70	0.0	11/16	106	6	0.06	AB	23 km NNE of 100-K
	10/03/27	13:27:58.16	46N32.06	119W57.67	4.06	-0.1	7/08	139	5	0.09	AC	25 km W of 200 West
	10/03/28	18:02:47.69	46N24.96	119W16.19	1.99	0.2	6/09	123	2	0.07	AC	Wooded Island Swarm
	10/03/28	18:07:50.17	46N24.93	119W16.27	1.74	0.3	9/12	122	1	0.05	AB	Wooded Island Swarm
	10/03/28	18:12:33.36	46N24.83	119W16.30	1.76	0.3	7/10	122	2	0.03	AB	Wooded Island Swarm
	10/03/28	18:29:24.30	46N24.93	119W16.38	2.21	-0.5	4/06	119	1	0.05	AD	Wooded Island Swarm
	10/03/28	18:29:45.31	46N24.84	119W16.22	1.86	0.2	6/10	124	2	0.07	AC	Wooded Island Swarm
	10/03/28	18:31:08.11	46N24.87	119W16.23	2.19	0.2	8/11	123	2	0.04	AB	Wooded Island Swarm
	10/03/28	18:31:45.11	46N25.21	119W17.11	0.05	-0.5	5/07	102	1	0.08	AD	Wooded Island Swarm
	10/03/28	18:33:19.98	46N24.85	119W16.29	2.27	0.3	6/09	122	2	0.06	AC	Wooded Island Swarm
	10/03/30	17:19:10.28	46N44.36	119W27.36	0.02	1.8	19/21	71	3	0.07	AA	15 km NE of 100-K
	10/04/07	22:53:42.07	46N38.90	119W07.77	3.67	0.5	12/15	120	8	0.07	AB	20 km S of Othello
	10/04/09	08:15:13.83	46N36.24	119W51.51	7.54	1.1	14/18	127	5	0.10	AB	18 km WNW of 200 West Area
	10/04/09	08:19:26.39	46N38.76	119W08.01	2.00	0.8	18/22	118	8	0.08	AB	20 km S of Othello
	10/04/09	18:22:28.25	46N35.85	119W51.56	7.10	1.5	12/14	128	7	0.08	AB	18 km WNW of 200 West Area
	10/04/09	18:22:28.30	46N35.92	119W51.26	7.27	1.5	17/18	88	4	0.08	AA	17 km WNW of 200 West Area
P	10/04/13	21:55:39.18	46N15.11	119W27.66	3.36		6/06	212	5	0.04	AD	14 km WSW of Richland
P	10/04/23	02:17:29.69	46N26.16	119W01.59	0.43		13/13	149	6	0.12	AC	21 km ENE of 300 Area
	10/04/23	10:18:26.80	46N42.09	119W28.54	0.42	1.2	19/24	45	3	0.08	AA	11 km NE of 100-K Area
	10/05/01	00:47:30.61	46N24.04	119W18.12	0.44	0.5	10/10	107	2	0.03	AB	Wooded Island Swarm
	10/05/08	11:57:43.25	46N49.89	119W41.41	3.02	3.0	35/35	34	12	0.13	AC	22 km NNW of 100-K Area
	10/05/11	02:09:31.77	46N25.13	119W16.92	0.40	0.1	6/07	190	1	0.09	AD	Wooded Island Swarm
	10/05/13	20:32:47.24	46N42.96	119W33.28	4.51	0.4	12/13	115	4	0.07	AB	9 km NNE of 100-K Area
	10/05/19	02:54:28.00	46N24.25	119W17.75	0.43	0.2	5/06	126	2	0.02	AD	Wooded Island Swarm
	10/05/24	00:35:43.93	46N49.71	119W41.63	2.84	0.9	17/17	145	12	0.13	AC	22 km NNW of 100-K Area

**Table 4.1.** (contd)

Type	Date	Time	Latitude	Longitude	Depth	Mag	NS/NP	Gap	Dmin	RMS	Q	Location
	10/05/24	17:58:10.77	46N00.17	119W29.42	0.55	1.0	8/08	154	24	0.15	BC	31 km SE of Prosser
P	10/05/28	19:49:27.29	46N26.81	119W03.31	0.05		7/07	140	4	0.14	AC	20 km ENE of 300 Area
P	10/06/02	20:08:49.68	46N26.06	119W01.23	3.40		5/05	115	6	0.07	AD	22 km ENE of 300 Area
	10/06/05	14:09:24.79	46N25.00	119W16.50	1.65	0.9	16/18	115	1	0.06	AB	Wooded Island Swarm
	10/06/08	05:50:53.33	46N38.90	119W45.34	16.69	-0.8	6/06	256	3	0.01	AD	12 km W of 100-K Area
	10/06/08	17:11:20.36	46N24.93	119W16.61	1.66	0.1	6/09	113	1	0.02	AC	Wooded Island Swarm
	10/06/10	11:42:38.33	46N40.26	119W13.73	0.99	0.7	16/17	104	0	0.07	AB	18 km SSW of Othello
P	10/06/15	19:43:12.92	46N23.00	119W56.43	0.79		10/10	130	22	0.08	AC	24 km NW of Prosser
	10/06/22	20:24:36.33	46N38.39	119W39.29	4.06	0.9	13/14	70	4	0.09	AA	4 km W of 100-K Area
	10/06/23	05:37:18.84	46N44.26	119W26.98	1.25	0.7	13/14	88	2	0.04	AA	16 km NE of 100-K Area
	10/06/23	08:17:07.86	46N44.34	119W26.99	0.95	-0.1	8/09	105	2	0.06	AB	16 km NE of 100-K Area
	10/06/25	13:10:18.13	46N36.73	119W46.44	15.92	-0.4	7/08	218	1	0.05	AD	12 km WNW of 200 West Area
	10/06/25	18:54:53.29	46N24.72	119W16.18	1.54	0.3	12/12	126	2	0.03	AB	Wooded Island Swarm
	10/06/28	10:51:48.05	46N49.56	119W41.48	5.63	2.6	13/14	102	12	0.07	AC	22 km NNW of 100-K Area
	10/07/04	22:03:49.43	46N24.70	119W23.71	4.13	-0.3	5/06	173	2	0.02	AD	4 km SW of 400 Area
	10/07/06	11:17:59.93	46N38.84	119W48.47	17.44	0.9	8/13	266	5	0.07	AD	16 km W of 100-K Area
	10/07/08	15:39:31.33	46N50.10	119W41.50	0.02	0.3	11/11	149	13	0.10	AC	23 km NNW of 100-K Area
	10/07/19	06:42:50.77	46N29.15	119W29.06	19.47	1.4	19/33	40	11	0.09	AA	9 km SSE of 200-E Area
	10/07/21	11:29:59.29	46N24.45	119W15.86	1.33	0.3	7/09	139	2	0.05	AC	Wooded Island Swarm
X	10/07/23	20:34:32.06	46N38.82	119W35.90	2.93		5/05	118	4	0.04	AD	0 km N of 100-K Area
	10/07/30	03:02:34.40	46N22.88	119W52.30	12.82	0.3	10/14	234	14	0.08	AD	21 km NNW of Prosser
	10/07/30	03:32:36.48	46N23.04	119W52.20	12.58	0.1	7/12	232	14	0.06	AD	21 km NNW of Prosser
	10/07/30	03:36:29.27	46N23.04	119W52.22	8.22	1.1	12/19	214	14	0.08	AD	21 km NNW of Prosser
	10/08/18	12:52:34.55	46N43.76	119W34.58	2.69	0.4	11/15	145	2	0.14	AC	10 km N of 100-K Area
	10/08/27	03:56:58.02	46N43.15	119W34.02	0.04	0.7	17/18	129	4	0.06	AB	9 km N of 100-K Area
	10/08/27	10:27:28.58	46N43.15	119W33.91	0.34	0.2	11/14	127	4	0.08	AB	9 km N of 100-K Area
	10/08/29	05:22:43.62	46N24.06	119W17.95	0.38	-0.5	3/08	153	2	0.05	AD	Wooded Island Swarm

**Table 4.1.** (contd)

Type	Date	Time	Latitude	Longitude	Depth	Mag	NS/NP	Gap	Dmin	RMS	Q	Location
	10/09/04	06:58:13.03	46N45.51	119W26.32	0.02	0.6	6/06	120	4	0.05	AC	18 km NE of 100-K Area
	10/09/05	13:28:20.92	46N24.82	119W15.95	0.03	-0.6	5/09	131	2	0.11	AD	Wooded Island Swarm
	10/09/05	20:29:19.14	46N33.94	119W40.70	7.04	-0.1	10/10	94	5	0.11	AB	3 km WNW of 200-W Area
	10/09/29	19:57:16.45	46N23.07	119W56.70	0.02	1.5	9/11	101	11	0.13	AC	24 km NW of Prosser



### Explanation of Table 4.1

<b>Type:</b>	P is Probable Blast; X is Confirmed Blast; blank is local earthquake.
<b>Date:</b>	The year and date in Universal Time Coordinated (UTC). UTC is used throughout this report unless otherwise indicated.
<b>Time:</b>	The origin time of the earthquake given in Coordinated Universal Time (UTC). To covert UTC to Pacific Standard Time, subtract eight hours; to Pacific Daylight Time, subtract seven hours.
<b>Latitude:</b>	North latitude, in degrees and minutes, of the earthquake epicenter.
<b>Longitude:</b>	West longitude, in degrees and minutes, of the earthquake epicenter.
<b>Depth:</b>	The depth of the earthquake in kilometers (km).
<b>Mag:</b>	The magnitude is expressed as coda-length magnitude $M_c$ , an estimate of local magnitude $M_L$ (Richter 1958). If magnitude is blank, a determination was not made.
<b>NS/NP:</b>	Number of stations/number of phases used in the solutions.
<b>Gap:</b>	Azimuthal gap; the largest angle (relative to the epicenter) containing no stations.
<b>DMIN:</b>	The distance from the earthquake epicenter to the closest station.
<b>RMS:</b>	The root-mean-square residual (observed arrival times minus the predicted arrival times) at all stations used to locate the earthquake. It is useful as a measure of quality of the solution only when five or more well-distributed stations are used in the solution. Good solutions are normally characterized by RMS values of less than about 0.3 s.
<b>Q:</b>	Quality factors; indicate the general reliability of the solution/location (A is best quality, D is worst). See Section 3.3 of this report, "Quality Factors."

## 4.3 Quality Factors (Q)

XPED assigns a two-letter **Quality factor** (Table 4.1) that indicates the general reliability of the solution (**A** is best quality, **D** is worst). Similar quality factors are used by the USGS for events located with the computer program HYPO71. The first letter of the quality code is a measure of the hypocenter quality based primarily on arrival time residuals. For example: Quality **A** requires a root-mean-square residual (**RMS**) less than 0.15 s, while a **RMS** of 0.5 s or more is **D** quality (other estimates of the location uncertainty also affect this quality parameter). The second letter of the quality code is related to the spatial distribution of stations that contribute to the event location, including the number of stations (**NS**), the number of p-wave and s-wave phases (**NP**), the largest gap in event-station azimuth distribution (**GAP**), and the closest distance from the epicenter to a station (**DMIN**). Quality **A** requires a solution with **NP** >8, **GAP** <90°, and **DMIN** <5 km (or the hypocenter depth if it is greater than 5 km). If **NP** ≤5, **GAP** >180°, or **DMIN** >50 km, the solution is assigned Quality **D**.

Uncertainties associated with estimated depths depend upon the number of stations and number of phase measurements (**NS/NP**) utilized in the XPED calculation. Generally speaking, if the number of phases exceeds 10 measurements, the depth estimate is considered to be reliable. In this case, the second letter in the quality evaluation is either "A" or "B" (cf. Table 4.1). For example, the number of phase measurements from earthquakes ultimately classified as "deep" events typically falls within the 10-20 measurement range; these depth estimates are considered reliable. However, the number of phase

measurements from earthquakes classified as “shallow” or “intermediate” may be less than 10 readings; in this case the depth estimate is less certain and the event could be classified as occurring in the CRBG or pre-basalt layers.

**Table 4.2.** Crustal Velocity Model for Eastern Washington (from Rohay et al. 1985)

Depth to Top of Velocity Layer (km)	Layer	Velocity (km/s)
0.0	Saddle Mountains and Wanapum Basalts and intercalated Ellensburg Formation	3.7
0.4	Grande Ronde Basalt and pre-basalt sediments	5.2
8.5	Basement, Layer 1	6.1
13.0	Basement, Layer 2	6.4
23.0	Sub-basement	7.1
38.0	Mantle	7.9

## 5.0 Seismic Activity – FY 2010

### 5.1 Summary

During FY 2010, the Hanford Seismic Network recorded 873 triggers on the seismometer system, which included 259 seismic events in the southeast Washington area and an additional 324 regional and teleseismic events. There were 210 events determined to be local earthquakes relevant to the Hanford Site.

One hundred and fifty-five earthquakes were detected in the vicinity of Wooded Island, located about eight miles north of Richland just west of the Columbia River. The Wooded Island events recorded this fiscal year were a continuation of the swarm events observed during fiscal year 2009 and reported in previous quarterly and annual reports (Rohay et al. 2009a, 2009b, 2009c, 2010a, 2010b, and 2010c). Most events were considered minor (coda-length magnitude [ $M_c$ ] less than 1.0) with the largest event recorded on February 4, 2010 ( $3.0M_c$ ). The estimated depths of the Wooded Island events are shallow (averaging approximately 1.5 km deep) with a maximum depth at 3.5 km. This places the Wooded Island events within the Columbia River Basalt Group (CRBG). Based upon the last two quarters (Q3 and Q4) data, activity at the Wooded Island area swarm has largely subsided. Pacific Northwest National Laboratory (PNNL) will continue to monitor for activity at this location.

The highest-magnitude events ( $3.0M_c$ ) were recorded on February 4, 2010 within the Wooded Island swarm (depth 2.4 km) and August 5, 2010 on or near the Saddle Mountain anticline (depth 3.0 km). This latter event is not considered unusual in that earthquakes have been previously recorded at this location, for example, in October 2006 (Rohay et al. 2007).

With regard to the depth distribution, 173 earthquakes were located at shallow depths (less than 4 km, most likely in the Columbia River basalts), 18 earthquakes were located at intermediate depths (between 4 and 9 km, most likely in the pre-basalt sediments), and 19 earthquakes were located at depths greater than 9 km, within the crystalline basement. Geographically, 178 earthquakes were located in known swarm areas, 4 earthquakes occurred on or near a geologic structure (Saddle Mountain anticline), and 28 earthquakes were classified as random events.

**Table 5.1.** Depth Distribution of Earthquakes for FY 2010

Category	First Quarter	Second Quarter	Third Quarter	Fourth Quarter	FY 2010
Shallow (0-4 km deep)	65	83	16	9	173 (82%)
Intermediate (4-9 km deep)	8	3	5	2	18 (9%)
Deep (greater than 9 km deep)	8	4	2	5	19 (9%)
Total	81	90	23	16	210

**Table 5.2.** Earthquake Locations for FY 2010

Seismic Sources	First Quarter 10/01–12/31	Second Quarter 1/01–3/31	Third Quarter 4/01–6/30	Fourth Quarter 7/01–9/30	FY 2010
Geologic Structure	0	0	3	1	4 (2%)
Frenchman Hills					
Saddle Mountains/ Royal Slope		1			1
Wahluke Slope			1	1	2
Coyote Rapids		3	5	3	11
Swarm Areas					
Wye	65	81	6	3	155
Cold Creek	5	2			7
Rattlesnake Mountain.					
Horse Heaven Hills	2				2
Total for swarm areas	72	87	12	7	178 (85%)
Random Events	9	3	8	8	28 (13%)
Total for all earthquakes	81	90	23	16	210


Table 4.1 includes all events that were recorded within the Hanford monitored area during FY 2010. Figure 5.1 shows the locations of those events.

## 5.2 First Quarter FY 2010 Earthquakes

For the Hanford Seismic Network, 81 local earthquakes were recorded during the first quarter of FY 2010 with 65 events recorded within the Wooded Island swarm. In the following discussions, minor earthquakes refer to seismic events for which the magnitude is less than 1.0  $M_c$ . A detailed discussion of first quarter FY 2010 seismic events may be found in Rohay et al. (2010a).

### 5.2.1 Location and Depth of Earthquakes

With regard to the depth distribution, 65 earthquakes occurred at shallow depths (less than 4 km, most likely in the Columbia River Basalt Group – CRBG), 8 earthquakes at intermediate depths (between 4 and 9 km, most likely in the pre-basalt sediments), and 8 earthquakes were located at depths greater than 9 km, within the basement. Geographically, 72 earthquakes occurred in swarm areas and 9 earthquakes were classified as random events.



**Figure 5.1.** FY 2010 Earthquakes Recorded Within the Hanford Monitoring Area (October 2009 through September 2010)

## 5.2.2 Major Anticlinal Ridges

No earthquakes were associated with the major geologic structures in the area surrounding the Hanford Site for the first quarter of FY 2010.

### **5.2.3 Earthquake Swarm Areas**

Seventy-two (72) earthquakes were characterized as swarm events during the first quarter of FY 2010. The only swarm events with magnitude greater than 1.0  $M_c$  were recorded near Wooded Island, within the Wye Swarm Area.

#### **Wye Swarm Area**

During the first quarter FY 2010, 65 earthquakes were recorded in the vicinity of Wooded Island. The Wooded Island swarm is located about eight miles north of Richland, west of the Columbia River about halfway between Hanford's 300 Area and Energy Northwest. All but five of the events were considered minor; the maximum magnitude event (2.5  $M_c$ ) occurred on December 22 at depth 2.1 km. The average depth of the Wooded Island events during the quarter was 1.4 km with a maximum depth at 3.1 km. This placed the Wooded Island events within the Columbia River Basalt Group (CRBG).

### **5.2.4 Random or Floating Events**

Nine earthquakes were classified as random events during the first quarter of FY 2010, all having magnitudes less than 1.0  $M_c$ .

## **5.3 Second Quarter FY 2010 Earthquakes**

The Hanford Seismic Network recorded 90 local earthquakes during the second quarter of FY 2010 with 81 events recorded within the Wooded Island swarm. A detailed discussion of second quarter FY 2010 seismic events may be found in Rohay et al. (2010b).

### **5.3.1 Location and Depth of Earthquakes**

With regard to the depth distribution, 83 earthquakes occurred at shallow depths (less than 4 km, most likely in the Columbia River basalts), 3 earthquakes at intermediate depths (between 4 and 9 km, most likely in the pre-basalt sediments), and 4 earthquakes were located at depths greater than 9 km, within the crystalline basement. Geographically, 87 earthquakes occurred in swarm areas and 3 earthquakes were classified as random events.

### **5.3.2 Major Anticlinal Ridges**

No earthquakes were associated with the major geologic structures in the area surrounding the Hanford Site for the second quarter of FY 2010.

### **5.3.3 Earthquake Swarm Areas**

Eighty-seven (87) earthquakes were characterized as swarm events in the second quarter of FY 2010. The only swarm events with magnitude greater than 1.0  $M_c$  were recorded within the Wye and Coyote Rapids Swarm Areas.

## **Coyote Rapids**

An event measuring 1.8  $M_c$  was recorded March 30, 2010 at depth 0.02 km within the CRBG.

## **Wye Swarm Area**

During the second quarter FY 2010, 81 Wooded Island events were recorded. Seventy-six of the events were considered minor and the maximum magnitude event (3.0  $M_c$ ) occurred February 4, 2010, at depth 2.4 km. The average depth of the Wooded Island events during the quarter was 1.6 km with a maximum depth at 3.5 km. This placed the Wooded Island events within the CRBG.

### **5.3.4 Random or Floating Events**

Three events were classified as random events during the second quarter of FY 2010. An event measuring 1.5  $M_c$  was recorded February 12, 2010, with epicenter located approximately 8 km south of Kennewick and 30 km east of the Horse Heavens Hills swarm area at depth 0.4 km, within the CRBG.

## **5.4 Third Quarter FY 2010 Earthquakes**

The Hanford Seismic Network recorded 23 local earthquakes during the third quarter of FY 2010 with only 6 events recorded in the vicinity of Wooded Island. All but seven of the events were considered minor. A detailed discussion of third quarter FY 2010 seismic events may be found in Rohay et al. (2010c).

### **5.4.1 Location and Depth of Earthquakes**

With regard to the depth distribution, 16 earthquakes were located at shallow depths (less than 4 km, most likely in the Columbia River basalts), 5 earthquakes were located at intermediate depths (between 4 and 9 km, most likely in the pre-basalt sediments), and 2 earthquakes were located at depths greater than 9 km, within the crystalline basement. Geographically, 12 earthquakes were located in swarm areas, 3 located at or near geologic structures, and 8 earthquakes were classified as random events.

### **5.4.2 Major Anticlinal Ridges**

The highest magnitude event (3.0  $M_c$ ) for the quarter was recorded on May 8, 2010 at depth 3.0 km with epicenter located on the northern limb of the Saddle Mountains east of Sentinel Gap. This placed the event on the Saddle Mountain anticline. An event measuring 2.6  $M_c$  was also recorded on June 28, 2010, with epicenter at nearly the same location at depth 5.6 km. Due to the high degree of precision in estimating the location of the May 8 event, both events are considered to originate from nearly the same location, approximately 3.0 km deep in the CRBG. These events are not considered unusual in that earthquakes have been previously recorded at this location, for example in October 2006 (Rohay et al. 2007).

### **5.4.3 Earthquake Swarm Areas**

Twelve earthquakes were characterized as swarm events in the third quarter of FY 2010. The only swarm event with magnitude greater than 1.0  $M_c$  was recorded within the Coyote Rapids Swarm Area.

## **Coyote Rapids Swarm Area**

An event measuring 1.2  $M_c$  was recorded on April 23, 2010, at depth 0.4 km within the CRBG.

### **5.4.4 Random or Floating Events**

Eight events were classified as random events during the third quarter of fiscal year 2010. Three events measuring 1.1-1.5  $M_c$  were recorded on April 9, 2010, with the epicenters located approximately 3 km west of the Hanford Site at depths 7.1-7.5 km within the pre-basalt sediments; these events are considered to be from the same location. An event measuring 1.0  $M_c$  was recorded May 24, 2010, with epicenter approximately 45 km south of the Hanford Site at depth 0.6 km, within the CRBG.

## **5.5 Fourth Quarter FY 2010 Earthquakes**

During the fourth quarter of FY 2010, the HSN recorded 281 triggers, which included 29 events located in the southeast Washington area and an additional 89 regional and teleseismic events. Sixteen earthquakes were located in the Hanford monitoring area plus one blast event.

### **5.5.1 Location and Depth of Earthquakes**

The depth distribution and geographic pattern of the sixteen earthquakes that occurred in the Hanford monitoring area are classified in Table 5.1 and Table 5.2. Epicenters of these events are shown in Figure 5.2. A three-dimensional depiction of these events is shown in Figure 5.3.

With regard to the depth distribution, nine earthquakes were located at shallow depths (less than 4 km, most likely in the Columbia River basalts), two earthquakes were located at intermediate depths (between 4 and 9 km, most likely in the pre-basalt sediments), and five earthquakes were located at depths greater than 9 km, within the crystalline basement. Geographically, seven earthquakes were located in swarm areas, one earthquake in the vicinity of a geologic structure, and eight earthquakes were classified as random events.

### **5.5.2 Major Anticlinal Ridges**

A minor event was recorded on July 8, 2010, at depth 0.02 km with epicenter located on the northern limb of the Saddle Mountains east of Sentinel Gap. This places the event on the Saddle Mountain anticline within the CRBG. The epicenter of this event is located at or near the locations of the previous May 8, 2010 and June 28, 2010 events, however the depth is estimated to be more shallow.

### **5.5.3 Earthquake Swarm Areas**

Seven earthquakes were characterized as swarm events in the fourth quarter of FY 2010.

## **Coyote Rapids Swarm Area**

A minor event was recorded on August 18, 2010, at depth 2.7 km within the CRBG. Two minor events were recorded on August 27, 2010 at shallow depths 0.04 and 0.3 km, respectively within the CRBG.



### **Wahluke Slope Swarm Area**

A minor event was recorded on September 4, 2010 at depth 0.02 km, within the CRBG.

### **Wye Swarm Area**


Three minor events were recorded in the vicinity of Wooded Island on July 21, 2010, August 29, 2010, and September 5, 2010, at depths between 0.03 and 1.3 km.

### **5.5.4 Random or Floating Events**


Eight earthquake events were recorded and classified as random events during the fourth quarter of FY 2010. A minor event with epicenter located approximately 5 km north of the southern Hanford Site boundary between the Cold Creek Mountain and Wye swarm areas at depth 4.1 km (intermediate zone) was recorded on July 4, 2010. A minor event with epicenter off-site near the western Hanford Site boundary and north of the Columbia River at depth 17.4 km (crystalline basement) was recorded on July 6, 2010. An event measuring 1.4  $M_c$  with epicenter located onsite between the Cold Creek Mountain and Wye swarm areas at depth 19.5 km (crystalline basement) was recorded on July 19.

Two minor events (located at depths 12.6 and 12.8 km) and a third event measuring 1.1  $M_c$  (located at depth 8.2 km) had nearly identical epicenters located offsite west of the Rattlesnake Mountain swarm area. These events were recorded within a 34-minute period on July 30. A common depth of 12.6 km (within the crystalline basement) was preferred for this interpretation.

A minor event with epicenter located approximately 1 km west of the 200-West Area at depth 7.0 km (intermediate zone) was recorded on September 5, 2010. A magnitude 1.5 ( $M_c$ ) event with epicenter located offsite west of the Rattlesnake Mountain swarm area at depth 0.02 km (CRBG) was recorded on September 29, 2010.



**Figure 5.2.** Earthquakes Occurring in the Hanford Monitoring Area Between July 1, 2010 and September 30, 2010 (Fourth Quarter)



**Figure 5.3.** Three Dimensional Depiction of Earthquakes Occurring in the Hanford Monitoring Area between July 1, 2010 and September 30, 2010 (Fourth Quarter)



## 6.0 Strong Motion Accelerometer Operations


The Hanford SMA network has been in continuous operation since November 20, 1998. Initially, the threshold used in the SMA network was 0.1% g. In 2006, the trigger threshold was reduced to 0.02% g when new instruments with greater storage capacity were installed. The lower trigger threshold saves the ground motion recordings from smaller, non-damaging earthquakes that can be useful in estimating the ground motion expected from larger earthquakes, and to confirm correct operation of the instruments by analyzing the smaller-amplitude triggers (see Section 2.2).

### 6.1 Fourth Quarter FY 2010 Triggers of the Hanford SMA Network

The Hanford SMA network was not triggered during the fourth quarter of FY 2010.


### 6.2 FY 2010 Triggers of the Hanford SMA Network

The Hanford SMA network was triggered numerous times by the Wooded Island swarm events during FY 2010. The largest magnitude event (3.0  $M_c$ ) for the Wooded Island swarm was recorded on February 4, 2010, and triggered the 300 Area SMA (located 4.5 km south of the epicenter) and the 400 Area SMA (located 6.5 km northwest of the epicenter). The peak horizontal and vertical accelerations were 1.1% g and 0.5% g at the 300 Area SMA (Figure 6.1) and 0.4% g and 0.2% g at the 400 Area SMA. The maximum horizontal acceleration was approximately one-half of the reportable action level for Hanford facilities (2% g) and no action was required.



**Figure 6.1.** Acceleration Records from the February 4, 2010 Wooded Island Swarm Event (Magnitude 3.0) at the 300 Area SMA

The 200-East Area SMA station (H2E) triggered and recorded the May 8, 2010 (3.0  $M_c$ ) and the June 28, 2010 (2.6  $M_c$ ) earthquakes located on or near the Saddle Mountains anticline. The vertical and horizontal accelerations were less than 0.04% g, only slightly larger than the trigger threshold of 0.02% g. The acceleration records from the May 8 earthquake are shown in Figure 6.2.



**Figure 6.2.** Acceleration Records from the 200 East SMA May 8, 2010 Event (Magnitude 3.0) Located On or Near the Saddle Mountain Anticline

## 7.0 Capabilities in the Event of a Significant Earthquake

The SMA network was designed to provide ground motion data in areas at the Hanford Site that have high densities of people and/or facilities containing hazardous materials, to ensure that the Hanford Site is in compliance with DOE Order 420.1B, Chapter IV, Section 3.d, “Seismic Detection.” The network also allows the HSAP to support Hanford Site emergency services organizations in complying with DOE Order G 420.1-1, Section 4.7, “Emergency Preparedness and Emergency Communications,” by providing area ground motion data in the event of an earthquake on the Hanford Site. This section summarizes the capabilities of the HSAP in the event of an earthquake at Hanford.

Historically, only a few facilities at the Hanford Site had instruments to provide data on peak ground accelerations or any type of ground motion. The current SMA instruments were located so that if an earthquake occurred, ground motion data would be readily available to assess the damage at the 100-K Area, the 200 East and West Areas, and the 300 and 400 Area facilities, which have the greatest concentration of people and also contain hazardous materials (Moore and Reidel 1996).

Many facilities at the Hanford Site have undergone various degrees of seismic analysis, either during design or during requalification. Although the seismic design of a building may be known, when an earthquake is “felt” in a facility on the Hanford Site, a determination must be made as to the extent of damage before it can be reoccupied and the systems restarted. A “felt” earthquake may not cause any significant damage to a building but, without adequate characterization of the ground motion, initial determination of the building’s possibility of having damage may be impossible.

In the event of a major regional earthquake such as the 2001 Nisqually event, building managers, emergency directors, and engineers can obtain ground motion data recorded by the SMA network from the HSAP in the Sigma V Building. This is done through the Hanford Site Emergency Services organization. Normal hours of operation for the HSAP are between 6 a.m. and 4:30 p.m., Monday through Friday. If a SMA is triggered, the HSAP will download events that were recorded and determine the peak ground accelerations. This information is then passed on to Hanford Site Emergency Services personnel where the facility engineers can use the data to determine if the ground motion exceeded, is equal to, or is less than the building design. This, along with assessments from trained engineers, allows the facility manager to make a rapid and cost-effective determination on whether a building is safe to re-occupy or should not be used until it has been inspected in more detail. Buildings that have designs exceeding the recorded ground motion could be put back into service very quickly; buildings with designs that are very close to or less than measured ground motion could be given priority for onsite damage inspections.





## 8.0 References

Campbell NP. 1989. "Structural and Stratigraphic Interpretation of Rocks under the Yakima Fold Belt, Columbia Basin, Based on Recent Surface Mapping and Well Data." In *Volcanism and Tectonism in the Columbia River Flood-Basalt Province*, SP Reidel and PR Hooper (eds.), Special Paper 239, pp. 209–222. Geological Society of America, Boulder, Colorado.

Crosson RS. 1972. "Small Earthquakes, Structure and Tectonics of the Puget Sound Region." *Bulletin of the Seismological Society of America* 62(5):1133–1171.

DOE. 1988. Site Characterization Plan for the Reference Location, Hanford, Washington – Consultation Draft. DOE/RW-0164, Vol. 1, U.S. Department of Energy, Washington, D.C.

DOE Order 420.1B, Chapter IV, Section 3.d. "Seismic Detection." U.S. Department of Energy, Washington, D.C.

DOE Order G 420.1-1, Section 4.7. "Emergency Preparedness and Emergency Communications." U.S. Department of Energy, Washington, D.C.

Fenneman NM. 1931. *Physiography of Western United States*. McGraw-Hill Book Company, Inc., New York.

Geomatrix. 1996. Probabilistic Seismic Hazard Analysis, DOE Hanford Site, Washington. WHC-SD-W236A-TI-002, Rev. 1, Westinghouse Hanford Company, Richland, Washington.

Moore C and SP Reidel. 1996. Hanford Site Seismic Monitoring Instrumentation Plan. WHC-SD-GN-ER-30036, Westinghouse Hanford Company, Richland, Washington.

Reidel SP and KR Fecht. 1994a. Geologic Map of the Richland 1:100,000 Quadrangle, Washington. Open File Report 94-8, Division of Geology and Earth Resources, Washington State Department of Natural Resources, Olympia.

Reidel SP and KR Fecht. 1994b. Geologic Map of the Priest Rapids 1:100,000 Quadrangle, Washington. Open File Report 94-13, Division of Geology and Earth Resources, Washington State Department of Natural Resources, Olympia.

Reidel SP, KR Fecht, MC Hagood, and TL Tolan. 1989. "Geologic Development of the Central Columbia Plateau." In *Volcanism and Tectonism in the Columbia River Flood-Basalt Province*, SP Reidel and PR Hooper (eds.), Special Paper 239, pp. 247-264. Geological Society of America, Boulder, Colorado.

Reidel SP, NP Campbell, KR Fecht, and KA Lindsey. 1994. "Late Cenozoic Structure and Stratigraphy of South-Central Washington." In *Regional Geology of Washington State*, E Cheney and R Lasmanis (eds.), Bulletin 80, pp. 159-180. Division of Geology and Earth Resources, Washington State Department of Natural Resources, Olympia.

Richter CF. 1958. Elementary Seismology. W. H. Freeman & Company, San Francisco, California.

Rohay AC, DW Glover, and SD Malone. 1985. Time-Term Analysis of Upper Crustal Structure in the Columbia Basin, Washington. RHO-BW-SA-435 P, Rockwell Hanford Operations, Richland, Washington.

Rohay AC, MD Sweeney, DC Hartshorn, RE Clayton, and JL Devary. 2007. Annual Hanford Seismic Report for Fiscal Year 2007. PNNL-17183, Pacific Northwest National Laboratory, Richland, Washington.

Rohay AC, MD Sweeney, DC Hartshorn, RE Clayton, and JL Devary. 2009a. Second Quarter Seismic Report for Fiscal Year 2009. PNNL-18633, Pacific Northwest National Laboratory, Richland, Washington.

Rohay AC, MD Sweeney, DC Hartshorn, RE Clayton, and JL Devary. 2009b. Third Quarter Seismic Report for Fiscal Year 2009. PNNL-18188, Pacific Northwest National Laboratory, Richland, Washington.

Rohay AC, MD Sweeney, DC Hartshorn, RE Clayton, and JL Devary. 2009c. Annual Hanford Seismic Report for Fiscal Year 2009. PNNL-19071, Pacific Northwest National Laboratory, Richland, Washington.

Rohay AC, MD Sweeney, DC Hartshorn, RE Clayton, and JL Devary. 2010a. First Quarter Seismic Report for Fiscal Year 2010. PNNL-19283, Pacific Northwest National Laboratory, Richland, Washington.

Rohay AC, MD Sweeney, DC Hartshorn, RE Clayton, and JL Devary. 2010b. Second Quarter Seismic Report for Fiscal Year 2010. PNNL-19513, Pacific Northwest National Laboratory, Richland, Washington.

Rohay AC, MD Sweeney, DC Hartshorn, RE Clayton, and JL Devary. 2010c. Third Quarter Seismic Report for Fiscal Year 2010. PNNL-19801, Pacific Northwest National Laboratory, Richland, Washington.

Wicks, CW, JS Gomberg, and CS Weaver. 2009. InSAR Measurement of Surface Deformation at the Hanford Reservation associated with the 2009 Wooded Island Earthquake Swarm. (abstract) American Geophysical Union Fall Meeting, December, 2009.

## Distribution

Distribution of the Annual Hanford Seismic Monitoring Report for FY 2010 will be accomplished via download of electronic PDF files from an external PNNL website. The following organizations will be notified by letter with instructions to download the electronic document.

### **No. of Copies**

### **No. of Copies**

#### **OFFSITE**

N. Rasmussen  
3140 Ravenshoe Drive  
Las Vegas, NV 89134

P. Rizzo  
105 Mall Boulevard, Suite 270E  
Expo Mart  
Monroeville, PA 15146

J. Litehiser  
Bechtel National, Inc.  
P.O. Box 193965  
San Francisco, CA 94119-3965

J. Zollweg  
Boise State University  
Department of Geosciences  
Boise, ID 83725

J. Kimball  
Defense Nuclear Facilities Safety Board  
625 Indiana Avenue NW, Suite 700  
Washington, DC 20004

- 3 Energy Northwest  
P.O. Box 968  
PE-24  
Richland, WA 99352  
S.M. Black  
D.W. Gregoire  
G.R. Lisle

Administrator  
Kennewick General Hospital  
P.O. Box 6128  
Kennewick, WA 99336

M. Stickney  
Montana Bureau of Mines and Geology  
Montana Tech University  
Earthquake Studies Office  
1300 West Park Street  
Butte, MT 59701-8997

S. Lilligren  
Nez Perce Tribe  
P.O. Box 365  
Lapwai, ID 83540

- 2 Oregon Department of Geology and  
Mineral Industries  
Suite 965, 800 NE Oregon Street #28  
Portland, OR 97232  
Library  
I. Madin

T. Conrads  
Parsons Constructors, Inc.  
3005 E. Ainsworth Street  
Pasco, WA 99301

R. Whale  
Shell E&P  
200 N. Dairy Ashford  
Houston, TX 77079

Bill Phillips  
Idaho Geological Survey  
P.O. Box 443014  
University of Idaho  
Moscow, ID 83844-3014

**No. of  
Copies**

3 University of Washington  
Seismology Laboratory  
Department of Earth and Space Sciences  
Box 351310  
Seattle, WA 98195-1310  
P. Bodin  
R. Steele  
J. Vidale

2 University of Washington  
U.S. Geological Survey  
Department of Earth and Space Sciences  
Box 351310  
Seattle, WA 98195-1310  
C. Weaver  
T. Yelin

I.G. Wong  
URS Corporation  
1333 Broadway, Suite 800  
Oakland, CA 94612

U.S. Fish and Wildlife Service  
64 Maple Street  
Burbank, WA 99323-8521

T. Brocher  
U.S. Geological Survey  
Mail Stop 977  
345 Middlefield Road  
Menlo Park, CA 94025

2 Washington Division of Geology and Earth  
Resources  
P.O. Box 4707  
Olympia, WA 98504-7007  
Library  
T. Walsh

**No. of  
Copies**

J. Schelling  
Washington Emergency Management  
Division  
Building 20, M/S: TA-20  
Camp Murray, WA 98430-5122

Washington State University  
Department of Geology  
P.O. Box 643420  
Pullman, WA 99164-2812

S.P. Reidel  
WSU Tri-Cities  
2710 University Drive  
Richland, WA 99354

R. Carson  
Department of Geology  
Whitman College  
345 Boyer Avenue  
Walla Walla, WA 99362

**ONSITE**

**9 DOE Richland Operations Office**

B.L. Charboneau	A6-33
O.A. Farabee	A5-11
K.L. Flynn	A2-15
T.Y. Hale	A7-89
R.D. Hildebrand	A6-38
J.G. Morse	A5-11
D.J. Thelen	A3-05
S.M. McDuffie	A5-19
B.E. Hill	A5-17
S.R. Weil	A5-15
D.C. Ward	A5-15

**4 DOE Office of River Protection**

R.A. Gilbert	H6-60
M.R. Moreno	H6-60
R.M. Venkata	H6-60
R.K. Nielsen	H6-60

**No. of  
Copies**

**2 DOE Pacific Northwest Science Office**

T.L. Davis	K9-42
G. Sauve	K9-42

**9 CH2M HILL Plateau Remediation  
Company**

M.E. Brown	G5-51
S.A. Fargo	H3-23
S.M. Faulk	A3-05
B.H. Ford	H8-43
M.I. Wood	H8-51
G.D. Cummins	H8-15
S.W. Petersen	R3-50
S.F. Miller	H8-15

**Stoller**

R.G. McCain	B2-62
-------------	-------

**No. of  
Copies**

**2 Washington River Protection Solutions**

D.A. Myers	S7-66
S.J. Eberlein	E6-31

**Washington State Department of Ecology**

J. Caggiano	H0-57
-------------	-------

**8 Pacific Northwest National Laboratory**

R.E. Clayton	K6-75
J.L. Devary	K6-96
B.E. Opitz	K6-75
A.C. Rohay	K6-75
M.D. Sweeney	K6-75
S.D. Unwin	K9-69
Hanford Technical Library	P8-55





*Proudly Operated by **Battelle** Since 1965*

902 Battelle Boulevard  
P.O. Box 999  
Richland, WA 99352  
1-888-375-PNNL (7665)

[www.pnl.gov](http://www.pnl.gov)



U.S. DEPARTMENT OF  
**ENERGY**