



US 20210163966A1

(19) **United States**(12) **Patent Application Publication****Deng et al.**(10) **Pub. No.: US 2021/0163966 A1**(43) **Pub. Date: Jun. 3, 2021**

(54) **PRODUCTION OF ORGANIC ACIDS FROM
ASPERGILLUS CIS-ACONITIC ACID
DECARBOXYLASE (CADA) DELETION
STRAINS**

C12P 7/42 (2006.01)*C12N 9/04* (2006.01)*C12P 7/48* (2006.01)(52) **U.S. Cl.**

CPC *C12N 15/80* (2013.01); *C12N 9/88*
(2013.01); *C12Y 401/01011* (2013.01); *C12N*
9/1096 (2013.01); *C12Y 206/01018* (2013.01);
C12N 9/0006 (2013.01); *C12Y 101/01059*
(2013.01); *C12P 7/48* (2013.01); *C12P 7/42*
(2013.01)

(71) Applicant: **Battelle Memorial Institute**, Richland,
WA (US)

(72) Inventors: **Shuang Deng**, Richland, WA (US);
Ziyu Dai, Richland, WA (US); **Jon K.**
Magnuson, Richland, WA (US)

(73) Assignee: **Battelle Memorial Institute**, Richland,
WA (US)

(21) Appl. No.: **17/172,660**

(22) Filed: **Feb. 10, 2021**

Related U.S. Application Data

(62) Division of application No. 16/393,149, filed on Apr.
24, 2019, now Pat. No. 10,947,548.

(60) Provisional application No. 62/661,804, filed on Apr.
24, 2018.

Publication Classification(51) **Int. Cl.***C12N 15/80* (2006.01)*C12N 9/88* (2006.01)*C12N 9/10* (2006.01)(57) **ABSTRACT**

This application provides recombinant *Aspergillus* fungi having an endogenous cis-aconitic acid decarboxylase (cadA) gene genetically inactivated, which allows aconitic acid production by the recombinant fungi. Such recombinant fungi can further include an exogenous nucleic acid molecule encoding aspartate decarboxylase (panD), an exogenous nucleic acid molecule encoding β -alanine-pyruvate aminotransferase (BAPAT), and an exogenous nucleic acid molecule encoding 3-hydroxypropionate dehydrogenase (HPDH). Kits including these fungi, and methods of using these fungi to produce aconitic acid and 3-hydroxypropionic acid (3-HP) are also provided.

Specification includes a Sequence Listing.

FIG. 1




FIG. 2




FIG. 3B




FIG. 3A




FIG. 4




FIG. 5




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 7




FIG. 8




FIG. 9A





FIG. 9B



PRODUCTION OF ORGANIC ACIDS FROM ASPERGILLUS CIS-ACONITIC ACID DECARBOXYLASE (CAD A) DELETION STRAINS

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application is a divisional of U.S. application Ser. No. 16/393,149 filed Apr. 24, 2019, which claims priority to U.S. Provisional Application No. 62/661,804 filed Apr. 24, 2018, both herein incorporated by reference in their entireties.

ACKNOWLEDGMENT OF GOVERNMENT SUPPORT

[0002] This disclosure was made with Government support under Contract DE-AC05-76RL0 1830 awarded by the U.S. Department of Energy. The Government has certain rights in the invention.

FIELD

[0003] Recombinant *Aspergillus* genetically inactivated for an endogenous cis-aconitic acid decarboxylase (cadA) gene are provided, as are methods of using such recombinant fungi to produce aconitic acid and 3-hydroxypropionic acid (3-HP).

BACKGROUND

[0004] Itaconic acid (IA) is utilized as a monomer or co-monomer to form polymers that are used as raw material for plastics, resins, synthetic fibers and elastomers, detergents and cleaners. *Aspergillus terreus* Thom, produces an appreciable amount of itaconic acid when grown in a glucose medium. Cell-free extracts of *Aspergillus terreus* contain cis-aconitic decarboxylase (cadA), which can decarboxylate cis-aconitic acid into equal moles of itaconic acid and carbon dioxide.

[0005] The itaconic acid gene cluster (IA cluster) includes four genes, including cis-aconitic acid decarboxylase (cadA), a predicted transcription factor (tf), mitochondrial organic acid transporter (mttA), and MFS (Major Facilitator Superfamily) type transporter (mfsA) located in plasma membranes. Expression of one or more genes of the IA gene cluster in heterologous hosts, including *E. coli*, *A. niger*, and *S. cerevisiae*, can result in the production of itaconic acid in non itaconic acid host microorganisms.

[0006] Characterization and regulation of genes in the IA biosynthesis cluster through gene deletion had not been previously investigated. The inventors used protoplast transformation to delete each gene in the IA cluster in *Aspergillus terreus*/*Aspergillus pseudoterreus*, which allowed for the effect on cell growth and IA production to be investigated.

SUMMARY

[0007] The role of cis-aconitic acid decarboxylase (cadA), a predicted transcription factor (tf), mitochondrial organic acid transporter (mttA), and MFS (Major Facilitator Superfamily) type transporter (mfsA) in IA biosynthesis in *A. pseudoterreus* ATCC 32359 is shown herein. Expressed Sequence Tag (EST) analysis showed a similar expression pattern among those four genes distinct from neighboring genes. Systematic gene deletion analysis demonstrated that

tf, cadA, mttA and mfsA genes in the cluster are essential for IA production. Interestingly, significant amounts of aconitic acid production was detected in the cadA deletion strain but not in the other deletion strains.

[0008] Based on these observations, a novel recombinant Δ cadA *Aspergillus* strain is provided, which can be used for aconitic acid and other organic acid production. Provided herein are isolated recombinant fungi (such as *Aspergillus filamentous* fungi) having a gene inactivation (also referred to herein as a gene deletion or functional deletion) of a cis-aconitic acid decarboxylase (cadA) gene (referred to herein as Δ cadA strains). In some examples, the *Aspergillus* fungi is *Aspergillus terreus* or *Aspergillus pseudoterreus*, or particular strains thereof (for example *A. pseudoterreus* ATCC32359 and *A. terreus* NRRL 1960). In particular examples, a Δ cadA strain exhibits one or more of the following characteristics: produces at least 2-fold, at least 3-fold, at least 3.5 fold, at least 5-fold, at least 8-fold, or at least 10-fold more total aconitic acid than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus* (for example at day 3, 4, 5, 6, 7, 8, 9 or 10 of production); produces at least 2-fold more cis-aconitic acid at day 5, 6, 7, 8, 9, or 10 of culturing in Riscaldati medium than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus*; produces at least 2-fold, at least 3-fold, at least 5-fold, or at least 10-fold more trans-aconitic acid at day 10 of culturing in Riscaldati medium than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus*; or combinations thereof. In some examples, such increases are relative to *Aspergillus terreus* strain ATCC 32359 grown under the same conditions.

[0009] In particular examples, a Δ cadA fungi further includes an exogenous nucleic acid molecule encoding aspartate 1-decarboxylase (panD), an exogenous nucleic acid molecule encoding β -alanine-pyruvate aminotransferase (BAPAT), and an exogenous nucleic acid molecule encoding 3-hydroxypropionate dehydrogenase (HPDH). The Δ cadA fungi expressing panD, BAPAT, and HPDH can be used to produce 3-HP. Such exogenous nucleic acid molecules can be part of one or more exogenous nucleic acid molecules, such as 1, 2 or 3 exogenous nucleic acid molecules. In one example, the exogenous nucleic acid molecule encoding panD has at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 53 or 65 and/or encodes a panD protein having at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 54. In one example, the exogenous nucleic acid molecule encoding BAPAT has at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 55, and/or encodes a BAPAT protein having at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 56. In one example, the exogenous nucleic acid molecule encoding HPDH has at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 57, and/or encodes a HPDH protein having at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 58. Such panD, BAPAT, and HPDH nucleic acid molecules can be part of a vector. In addition, expression of the panD, BAPAT, and HPDH can be driven by one or more promoters.

[0010] The endogenous cadA gene is genetically inactivated in some examples by a deletion mutation (complete or partial) or by insertional mutation (e.g., by insertion of an antibiotic resistance gene, such as hygromycin). In some

examples, prior to its genetic inactivation, the *cadA* gene encodes a protein having at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 50 or 52. In some examples, prior to its genetic inactivation, the *cadA* gene (or a *cadA* coding sequence) comprises at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 49, 51, 59 or 92.

[0011] Also provided herein are compositions (such as a culture media or fermentation broth) and kits that include a *Aspergillus* Δ *cadA* strain. Also provided herein are compositions (such as a culture media or fermentation broth) and kits that include an *Aspergillus* *AcadA* strain that also express panD, BAPAT, and HPDH, in some examples such genes are exogenous to the fungi. In some examples, the composition or kit includes Riscaldati medium (such as modified Riscaldati medium with 20 \times trace elements).

[0012] Also provided herein are methods of making aconitic acid (such as cis-aconitic acid, trans-aconitic acid, or both) using the disclosed *Aspergillus* Δ *cadA* strains. For example, such a method can include culturing an isolated Δ *cadA* *Aspergillus* under conditions that permit the fungus to make aconitic acid, thereby producing aconitic acid. For example, the *AcadA* fungus can be cultured in Riscaldati medium. In some examples, the method further includes isolating the aconitic acid produced, for example isolating it from the culture media or from the fungus.

[0013] Also provided herein are methods of making 3-hydroxypropionic acid (3-HP using the disclosed *Aspergillus* Δ *cadA* strains that also expresses panD, BAPAT, and HPDH (which can be exogenous). For example, such a method can include culturing an isolated Δ *cadA* *Aspergillus* that also expresses panD, BAPAT, and HPDH under conditions that permit the fungus to make 3-HP, thereby producing 3-HP. For example, the Δ *cadA* fungus that also expresses panD, BAPAT, and HPDH can be cultured in Riscaldati medium (such as one including 20 \times trace elements). In some examples, the method further includes isolating the 3-HP produced, for example isolating it from the culture media or from the fungus.

[0014] The foregoing and other objects and features of the disclosure will become more apparent from the following detailed description, which proceeds with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] FIG. 1. Hypothesized itaconic acid (IA) production and transport pathway in *Aspergillus pseudoterreus* and *Aspergillus terreus*. Glucose is utilized by *A. terreus* and *A. pseudoterreus* to form pyruvate and is subsequently converted to citric acid for tricarboxylic acid (TCA) cycle in the mitochondria. Citric acid is dehydrated to cis-aconitic acid, which is then transported from mitochondria to cytosol through transporter mttA. In the cytosol, cis-aconitic acid is decarboxylated into itaconic acid and CO₂ by cis-aconitic decarboxylase. Finally, itaconic acid secreted outside of cell through transporters, for example mfsA.

[0016] FIG. 2. *Aspergillus pseudoterreus* ATCC 32359 fermentation data for collecting samples for EST sequencing. A 20 L volume of Riscaldati production medium (see Riscaldati et al., *J Biotechnol* 2000, 83:219-230) in a 30 L working volume Sartorius fermenter was inoculated with 10⁶ *A. pseudoterreus* spores per ml. The three samples referred to as “preproduction, production onset and produc-

tion” were collected at 40, 50 and 62 hours, respectively. Itaconic acid and glucose data are shown on the left y-axis and fungal ash free dry weight (AFDW) is shown on the right y-axis.

[0017] FIGS. 3A-3B. *Aspergillus pseudoterreus* IA Cluster Analysis after five days growth in Riscaldati medium. Spores 0.5 \times 10⁸ were inoculated into 50 ml of production media for IA production as described in Riscaldati et al. (*J Biotechnol* 2000, 83:219-230). The cultivation was performed at 30° C. on a rotary shaker at 150 rpm. At the end of five days, samples were obtained for HPLC analysis and biomass measurement. (A) Dry mass measurement of wild type and mutant strains (B) Itaconic acid production of wild type and mutant strains. The average obtained from three independent experiments are shown. Error bars represent standard deviations from the means.

[0018] FIG. 4. Kinetics of itaconic acid production by wild type *A. pseudoterreus* and Δ t strains grown in production media at 30° C. Spores 0.5 \times 10⁸ were inoculated into 50 ml of production media for itaconic acid production as described in Riscaldati et al. (*J Biotechnol* 2000, 83:219-230). The cultivation was performed at 30° C. on a rotary shaker at 150 rpm. All experiments were done in three replicates. At day 2, 4, 6, and 7, HPLC analysis was performed to determine amount of IA produced. Each sample was measured in five replicates. Error bars represent standard deviation from the means.

[0019] FIG. 5. Real-time (RT)-PCR analysis of the relative levels of mttA, *cadA*, mfsA mRNAs in wild type and Δ t strains. Spores 0.5 \times 10⁸ were inoculated into 50 ml of production media for itaconic acid production as described in Riscaldati et al. (*J Biotechnol* 2000, 83:219-230). The cultivation was performed at 30° C. on a rotary shaker at 150 rpm. All experiments were done in three biological replicates. At day 3, samples were collected and RNA was extracted for RT-PCR. The average of results obtained from five independent RNA preparations is shown. All transcript levels were measured in triplicate for each RNA preparation. Error bars represent standard deviations from the means. Compared to wild type, expression level of mttA, *cadA* and mfsA were decreased 57, 37 and 23 fold in the Δ t strain.

[0020] FIGS. 6A-6C. Aconitic acid production in Δ *cadA* strain. The cultivation was performed at 30° C. on a rotary shaker at 150 rpm. All experiments were done in three biological replicates. (A) at day 5, only Δ *cadA* produced cis-aconitic and trans-aconitic acid, while wild type and other mutants did not. (B) Time course of cis- and trans-aconitic acid production in Δ *cadA* strain over 10 days. (C) Comparison of total aconitic acid production between wild type and Δ *cadA* mutant strains.

[0021] FIG. 7. Arrangement of transgene expression cassette for 3-HP Production in *A. pseudoterreus* with a synthetic beta-alanine pathway. A description of each Fragment is described in Example 8. The relevant fragments were cloned into pBlueScript SK(−) vector linearized with restriction enzyme H3/PstI. The whole expression cassette was linearized with restriction enzyme XhoI for the protoplast transformation for homologous recombination at *cadA* locus.

[0022] FIG. 8. Southern blot confirmation of *cadA* gene interruption by 3HP transgene expression cassette (FIG. 7). The *cadA* gene in the transgenic strains #2 (3HP-2) and #6 (3HP-6) was disrupted by the homologous recombination,

while the random integration occurred in the strains #4 (3HP-4) and #5 (3HP-5). No insertion was observed in strains #1 and #3.

[0023] FIGS. 9A-9B. 3-HP production. *A. pseudoterreus* having a genetically inactivated cadA locus alone (cad1Δ), or additionally expressing panD, BAPAT, and HPDH (3HP-2, 3HP-4, 3HP-5, and 3-HP6), were grown at 30° C. on a rotary shaker at 200 rpm for (A) 7 days, or (B) over 8 days, in the Riscaldati media with 20× TE, and 3-HP present in the supernatant measured using HPLC.

SEQUENCE LISTING

[0024] The nucleic acid sequences listed in the accompanying sequence listing are shown using standard abbreviations for nucleotide bases and amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. The sequence listing submitted herewith, generated on Feb. 5, 2021, 80 kb, is herein incorporated by reference. In the accompanying sequence listing:

[0025] SEQ ID NOS: 1-8 are primers used to delete the tf gene in *A. pseudoterreus*.

[0026] SEQ ID NOS: 9-16 are primers used to delete the mttA gene in *A. pseudoterreus*.

[0027] SEQ ID NOS: 17-24 are primers used to delete the cadA gene in *A. pseudoterreus*.

[0028] SEQ ID NOS: 25-32 are primers used to delete the mfsA gene in *A. pseudoterreus*.

[0029] SEQ ID NOS: 33-40 are primers used to delete the p450 gene in *A. pseudoterreus*.

[0030] SEQ ID NOS: 41-42 are primers used to amplify mttA in *A. pseudoterreus*.

[0031] SEQ ID NOS: 43-44 are primers used to amplify cadA in *A. pseudoterreus*.

[0032] SEQ ID NOS: 45-46 are primers used to amplify mfsA in *A. pseudoterreus*.

[0033] SEQ ID NOS: 47-48 are primers used to amplify benA in *A. pseudoterreus*.

[0034] SEQ ID NOS: 49 and 50 are exemplary cadA nucleic acid and protein sequences, respectively, from *A. terreus* (GenBank Accession Nos. AB326105.1 and BAG49047.1).

[0035] SEQ ID NOS: 51 and 52 are exemplary cadA nucleic acid and protein sequences, respectively, from *A. vadensis* CBS 113365 (GenBank Accession Nos. XM_025706777.1 and XP_025563141.1).

[0036] SEQ ID NOS: 53 and 54 are exemplary aspartate 1-decarboxylase (panD) nucleic acid and protein sequences, respectively, from *Tribolium castaneum* (GenBank Accession Nos. NM_001102585.1 and NP_001096055.1). Coding sequence nt 41-1663.

[0037] SEQ ID NOS: 55 and 56 are exemplary β-alanine-pyruvate aminotransferase (BAPAT) nucleic acid and protein sequences, respectively, from *Bacillus cereus* AH1272 (GenBank Accession Nos. ACMS01000158.1 (complement (10606 . . . 11961)) and EEL86940.1).

[0038] SEQ ID NOS: 57 and 58 are exemplary 3-hydroxypropionate dehydrogenase (HPDH) nucleic acid and protein sequences (GenBank Accession No. WP_000636571), respectively.

[0039] SEQ ID NO: 59 is an *A. pseudoterreus* 5'-cadA nucleic acid sequence.

[0040] SEQ ID NOS: 60-61 are primers used to isolate an *A. pseudoterreus* 5'-cadA gene.

[0041] SEQ ID NO: 62 is an *A. niger* gpdA promoter nucleic acid sequence.

[0042] SEQ ID NOS: 63-64 are primers used to isolate an *A. niger* gpdA promoter.

[0043] SEQ ID NO: 65 is panD cDNA of *Tribolium castaneum* with codon optimization for *A. pseudoterreus*.

[0044] SEQ ID NOS: 66-67 are primers used to isolate panD cDNA of *Tribolium castaneum* with codon optimization for *A. pseudoterreus*.

[0045] SEQ ID NO: 68 is a bidirectional terminator from *A. niger* elf3/multifunctional chaperone.

[0046] SEQ ID NOS: 69-70 are primers used to isolate bidirectional terminator from *A. niger* elf3/multifunctional chaperone.

[0047] SEQ ID NO: 71 is codon optimized synthetic cDNA of β-alanine-pyruvate aminotransferase (BAPAT) of *Bacillus cereus*.

[0048] SEQ ID NOS: 72-73 are primers used to isolate a codon optimized synthetic cDNA of BAPAT of *Bacillus cereus*.

[0049] SEQ ID NO: 74 is an *A. niger* eno1 promoter.

[0050] SEQ ID NOS: 75-76 are primers used to isolate an *A. niger* eno1 promoter.

[0051] SEQ ID NO: 77 is an *A. nidulans* gpdA promoter.

[0052] SEQ ID NOS: 78-79 are primers used to isolate an *A. nidulans* gpdA promoter.

[0053] SEQ ID NO: 80 is the codon optimized synthetic cDNA of *E. coli* 3-hydroxypropionate dehydrogenase (HPDH).

[0054] SEQ ID NOS: 81-82 are primers used to isolate a codon optimized synthetic cDNA of *E. coli* HPDH.

[0055] SEQ ID NO: 83 is a trpC terminator of *A. nidulans*.

[0056] SEQ ID NOS: 84-85 are primers used to isolate the trpC terminator of *A. nidulans*.

[0057] SEQ ID NO: 86 is a trpC terminator of *A. nidulans*.

[0058] SEQ ID NOS: 87-88 are primers used to isolate a trpC terminator of *A. nidulans*.

[0059] SEQ ID NO: 89 is an *A. oryzae* ptrA selection marker gene.

[0060] SEQ ID NOS: 90-91 are primers used to isolate the *A. oryzae* ptrA selection marker gene.

[0061] SEQ ID NO: 92 is an *A. pseudoterreus* 3'-cadA gene.

[0062] SEQ ID NOS: 93-94 are primers used to isolate an *A. pseudoterreus* 3'-cadA gene fragment.

[0063] SEQ ID NO: 95 is a combination of Fragments 7 to 9 (SEQ ID NOS: 77, 80, and 83, respectively).

[0064] SEQ ID NO: 96 is a primer used to isolate Fragments 7 to 9 (in combination with SEQ ID NO: 88).

[0065] SEQ ID NO: 97 is a combination of Fragments 11 and 12 (SEQ ID NOS: 89 and 92, respectively).

[0066] SEQ ID NO: 98 is a primer used to isolate Fragments 11 to 12 (in combination with SEQ ID NO: 90).

DETAILED DESCRIPTION

[0067] Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew et al. (eds.), *The Encyclopedia of*

Molecular Biology, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).


[0068] The singular terms “a,” “an,” and “the” include plural referents unless context clearly indicates otherwise. Similarly, the word “or” is intended to include “and” unless the context clearly indicates otherwise. Hence “comprising A or B” means including A, or B, or A and B. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. All publications, references and Genbank® Accession numbers (the sequence available on Apr. 24, 2019) mentioned herein are incorporated by reference in their entireties. The materials, methods, and examples are illustrative only and not intended to be limiting.

[0069] In order to facilitate review of the various embodiments of the disclosure, the following explanations of specific terms are provided:

[0070] 3-hydroxypropionate dehydrogenase (HPDH): EC 1.1.1.59 An enzyme that catalyzes the chemical reaction: 3-hydroxypropionate+NAD⁺↔3-oxopropionate+NADH+H⁺. The term HPDH includes any HPDH gene (such as a bacterial or fungal panD sequence), cDNA, mRNA, or protein, that is a HPDH that can convert 3-hydroxypropionate and NAD⁺ into 3-oxopropionate, NADH, and H⁺ and vice versa. Expression or increased expression of HPDH, for example in an *Aspergillus* also expressing BAPAT and panD and having a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

[0071] HPDH sequences are publicly available. For example, SEQ ID NO: 57 discloses an HPDH coding sequence and GenBank® Accession No: WP_000636571 discloses an HPDH protein sequence (SEQ ID NO: 58); GenBank® Accession Nos. FR729477.2 (nt 1005136 . . . 1005885) and CBY27203.1 disclose exemplary *Yersinia enterocolitica* subsp. *paleartica* Y11 HPDH nucleic acid and protein sequences, respectively; and GenBank® Accession Nos: CP004083.1 (complement(1399227 . . . 1399973)) and AJQ99264.1 disclose exemplary *Enterobacteriaceae bacterium* bta3-1 HPDH nucleic acid and protein sequences, respectively. However, one skilled in the art will appreciate that in some examples, a HPDH sequence can include variant sequences (such as allelic variants and homologs) that retain HPDH activity and when expressed in an *Aspergillus* also expressing BAPAT and panD and with a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

[0072] Aconitic acid: An organic acid with two isomers, cis- and trans-aconitic acid. The Δ cadA fungi provided herein can be used to produce cis- and trans-aconitic acid.



[0073] Aspartate 1-decarboxylase (panD): EC 4.1.1.11. An enzyme that catalyzes the chemical reaction: L-aspartate↔beta-alanine+CO₂. The term panD includes any panD gene (such as a bacterial or fungal panD sequence), cDNA, mRNA, or protein, that is a panD that can convert L-aspartate into beta-alanine+CO₂ and vice versa. Expression or increased expression of panD, for example in an *Aspergillus* also expressing BAPAT and HPDH and having a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

[0074] panD sequences are publicly available. For example, GenBank® Accession Nos: NM_001102585.1 and NP_001096055.1 disclose *Tribolium castaneum* panD nucleic acid and protein sequences, respectively (SEQ ID NOS: 55 and 56); GenBank® Accession Nos. CP002745.1 (complement(4249351 . . . 4249824)) and AEK63458.1 disclose exemplary *Collimonas fungivorans* Ter331 panD nucleic acid and protein sequences, respectively; and GenBank® Accession Nos: CP029034.1 (nt 1201611 . . . 1201994) and AWE15802.1 disclose exemplary *Bacillus velezensis* panD nucleic acid and protein sequences, respectively. However, one skilled in the art will appreciate that in some examples, a panD sequence can include variant sequences (such as allelic variants and homologs) that retain panD activity and when expressed in an *Aspergillus* also expressing BAPAT and HPDH and with a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

[0075] β -alanine-pyruvate aminotransferase (BAPAT): EC 2.6.1.18. An enzyme that can catalyze the reaction L-alanine+3-oxopropionate↔beta-alanine+pyruvate. The term BAPAT includes any BAPAT gene (such as a bacterial or fungal panD sequence), cDNA, mRNA, or protein, that is a BAPAT that can convert beta-alanine and pyruvate to L-alanine and 3-oxopropionate [or malonic semialdehyde], and vice versa. Expression or increased expression of BAPAT, for example in an *Aspergillus* also expressing HPDH and panD and having a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at

least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

[0076] BAPAT sequences are publicly available. For example, GenBank® Accession Nos: ACMS01000158.1 (complement(10606 . . . 11961)) and EEL86940.1 disclose *Bacillus cereus* AH1272 BAPAT nucleic acid and protein sequences, respectively (SEQ ID NOS: 55 and 56); GenBank® Accession Nos. DF820429.1 (complement (241627 . . . 242967)) and GAK28710.1 disclose exemplary *Serratia liquefaciens* FK01 BAPAT nucleic acid and protein sequences, respectively; and GenBank Accession Nos: LGUJ01000001.1 complement (92812 . . . 94140) and KOY12524.1 disclose exemplary *Bradyrhizobium diazoefficiens* BAPAT nucleic acid and protein sequences, respectively. However, one skilled in the art will appreciate that in some examples, a BAPAT sequence can include variant sequences (such as allelic variants and homologs) that retain BAPAT activity and when expressed in an *Aspergillus* also expressing HPDH and panD and with a genetically inactivated cadA gene (AcadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

[0077] cadA (cis-aconitic acid decarboxylase): The cadA gene encodes an enzyme (EC 4.1.1.6) that catalyzes the chemical reaction $\text{cis-aconitate} \rightleftharpoons \text{itaconate} + \text{CO}_2$. The term cadA (or cadA) includes any cadA gene (such as a fungal cadA sequence), cDNA, mRNA, or protein, that is a cadA that can catalyze the decarboxylation of cis-aconitate to itaconate and CO_2 and vice versa, and when genetically inactivated results in a fungus that produces more aconitic acid than the parent strain without a genetically inactivated cadA gene (such as at least 20%, at least 30%, at least 50%, at least 60%, at least 75%, at least 100%, at least 200%, at least 500%, or 1000% more than a parent strain under the same growing conditions, for example at day 5 of production). In some examples, a parental strain containing a functional native cadA sequence does not produce detectable aconitic acid. In some examples, genetic inactivation of cadA results in a fungus that produces more trans-aconitic acid than cis-aconitic acid at day 10 of production, (such as at least 2-fold, at least 3-fold, at least 4-fold, or at least 5-fold more at day 10 of production).

[0078] cadA sequences are publicly available for many species of *Aspergillus*. For example, GenBank® Accession Nos: AB326105.1 and BAG49047.1 disclose *Aspergillus terreus* cadA nucleic acid and protein sequences, respectively (SEQ ID NOS: 49 and 50); GenBank® Accession Nos: XM_025706777.1 and XP_025563141.1 disclose *Aspergillus vadensis* CBS 113365 cadA nucleic acid and protein sequences, respectively (SEQ ID NOS: 51 and 52); and GenBank® Accession Nos: XM_025663103.1 and XP_025520527.1 disclose *Aspergillus piperis* CBS 112811 cadA nucleic acid and protein sequences, respectively. However, one skilled in the art will appreciate that in some examples, a cadA sequence can include variant sequences (such as allelic variants and homologs) that retain cadA activity but when genetically inactivated in *Aspergillus* results in a fungus that has an ability to produce more aconitic acid than the parent strain without a genetically inactivated cadA gene (such as at least 20%, at least 30%, at

least 50%, at least 60%, at least 75%, at least 100%, at least 200%, at least 500%, or 1000% more than a parent strain under the same growing conditions, for example at day 5 of production).

[0079] Detectable: Capable of having an existence or presence ascertained. For example, production of aconitic acid or 3-HP is detectable if the signal generated is strong enough to be measurable.

[0080] Exogenous: The term “exogenous” as used herein with reference to nucleic acid and a particular cell refers to any nucleic acid that does not originate from that particular cell as found in nature. Thus, a non-naturally-occurring nucleic acid is considered to be exogenous to a cell once introduced into the cell. A nucleic acid that is naturally-occurring also can be exogenous to a particular cell. For example, an entire chromosome isolated from cell X is an exogenous nucleic acid with respect to cell Y once that chromosome is introduced into cell Y, even if X and Y are the same cell type.

[0081] In some examples, the panD, BAPAT, and HPDH nucleic acid or protein expressed in an *Aspergillus terreus* or *Aspergillus pseudoterreus* fungi does not naturally occur in the *Aspergillus terreus* or *Aspergillus pseudoterreus* fungi and is therefore exogenous to that fungi. For example, the panD, BAPAT, and HPDH nucleic acid molecule introduced into an *Aspergillus terreus* or *Aspergillus pseudoterreus* fungi can be from another organism, such as a bacterial panD, BAPAT, and HPDH sequence.

[0082] Genetic enhancement or up-regulation: When used in reference to the expression of a nucleic acid molecule, such as a gene, refers to any process which results in an increase in production of a gene product. A gene product can be RNA (such as mRNA, rRNA, tRNA, and structural RNA) or protein. Examples of processes that increase transcription include those that facilitate formation of a transcription initiation complex, those that increase transcription initiation rate, those that increase transcription elongation rate, those that increase processivity of transcription and those that relieve transcriptional repression (for example by blocking the binding of a transcriptional repressor). Gene up-regulation can include inhibition of repression as well as expression above an existing level. Examples of processes that increase translation include those that increase translational initiation, those that increase translational elongation and those that increase mRNA stability. In one example, additional copies of genes are introduced into a cell in order to increase expression of that gene in the resulting transgenic cell.

[0083] Gene up-regulation includes any detectable increase in the production of a gene product. In certain examples, production of a gene product increases by at least 1.5-fold, at least 2-fold, or at least 5-fold, such as aspartate decarboxylase (panD), β -alanine-pyruvate aminotransferase (BAPAT), and 3-hydroxypropionate dehydrogenase (HPDH). For example, expression of a panD, BAPAT, and HPDH genes in *Aspergillus* (e.g., *A. terreus*) results in an *Aspergillus* strain having increased levels of the panD, BAPAT, and HPDH proteins, respectively, relative to the parent strain, which can permit the recombinant fungus to produce 3-HP. Genetic enhancement is also referred to herein as “enhancing or increasing expression.”

[0084] Genetic inactivation or down-regulation: When used in reference to the expression of a nucleic acid molecule, such as a gene, refers to any process which results in

a decrease in production of a gene product. A gene product can be RNA (such as mRNA, rRNA, tRNA, and structural RNA) or protein. Therefore, gene down-regulation or deactivation includes processes that decrease transcription of a gene or translation of mRNA.

[0085] For example, a mutation, such as a substitution, partial or complete deletion, insertion, or other variation, can be made to a gene sequence that significantly reduces (and in some cases eliminates) production of the gene product or renders the gene product substantially or completely non-functional. For example, a genetic inactivation of the *cadA* gene in *Aspergillus* (e.g., *A. pseudoterreus*) results in *Aspergillus* having a non-functional or non-existent *cadA* protein, which results in the recombinant fungus to produce more aconitic acid. Genetic inactivation is also referred to herein as “functional deletion”.

[0086] Isolated: To be significantly separated from other agents. An “isolated” biological component (such as a nucleic acid molecule or protein) has been substantially separated, produced apart from, or purified away from other biological components in the cell of the organism in which the component occurs, for example, other chromosomal and extra-chromosomal DNA and RNA, and proteins. Nucleic acid molecules and proteins which have been “isolated” include nucleic acid molecules and proteins purified by standard purification methods. The term also embraces nucleic acid molecules and proteins prepared by recombinant expression in a host cell as well as chemically synthesized proteins and nucleic acids. Samples of isolated biological components include samples of the biological component wherein the biological component represents greater than 90% (for example, greater than 95%, such as greater than 98%) of the sample.

[0087] An “isolated” microorganism (such as a Δ *cadA* strain of *Aspergillus*) has been substantially separated or purified away from microorganisms of different types, strains, or species. Microorganisms can be isolated by a variety of techniques, including serial dilution and culturing and resistance to certain chemicals, such as antibiotics. In some examples, an isolated *AcadA* strain of *Aspergillus* is at least 90% (for example, at least 95%, as at least 98%, at least 99%, or at least 99.99%) pure.

[0088] Mutation: A change in a nucleic acid sequence (such as a gene sequence) or amino acid sequence, for example as compared to a nucleic acid or amino acid sequence present in a wild-type or native organism. In particular examples, a mutation is introduced into a *cadA* gene in *Aspergillus*. Mutations can occur spontaneously, or can be introduced, for example using molecular biology methods (e.g., thereby generating a recombinant or transformed cell or microorganism). In particular examples, a mutation includes one or more nucleotide substitutions, deletions, insertions, or combinations thereof. In particular examples, the presence of one or more mutations in a gene can significantly inactivate and reduce expression of that gene.

[0089] Promoter: An array of nucleic acid control sequences which direct transcription of a nucleic acid. A promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements which can be located as much as several thousand base pairs from the start site of transcription. In some examples, a promoter

is bi-directional. Native and non-native promoters can be used to drive expression of a gene, such as *panD*, *BAPAT*, and *HPDH*. Exemplary promoters that can be used include but are not limited to: *enl* promoter from *A. niger*, and *dthl* from *A. nidulans* or *A. niger*.

[0090] Examples of promoters include, but are not limited to the SV40 promoter, the CMV enhancer-promoter, and the CMV enhancer/ β -actin promoter. Both constitutive and inducible promoters can be used in the methods provided herein (see e.g., Bitter et al., *Methods in Enzymology* 153: 516-544, 1987). Also included are those promoter elements which are sufficient to render promoter-dependent gene expression controllable for cell-type specific, tissue-specific, or inducible by external signals or agents; such elements may be located in the 5' or 3' regions of the gene. Promoters produced by recombinant DNA or synthetic techniques can also be used to provide for transcription of the nucleic acid sequences.

[0091] Recombinant: A recombinant nucleic acid molecule or protein is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. In particular examples, this artificial combination is accomplished by chemical synthesis or by the artificial manipulation of isolated segments of nucleic acids, for example, by genetic engineering techniques such as those described in Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 3d ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001. The term recombinant includes nucleic acid molecules that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid molecule. A recombinant or transformed organism or cell, such as a recombinant *Aspergillus*, is one that includes at least one exogenous nucleic acid molecule, such as one used to genetically inactivate an endogenous *cadA* gene, and one used to express a non-native protein, such as exogenous *panD*, *BAPAT*, and *HPDH* nucleic acid coding sequences.

[0092] Sequence identity/similarity: The identity/similarity between two or more nucleic acid sequences, or two or more amino acid sequences, is expressed in terms of the identity or similarity between the sequences. Sequence identity can be measured in terms of percentage identity; the higher the percentage, the more identical the sequences are. Sequence similarity can be measured in terms of percentage similarity (which takes into account conservative amino acid substitutions); the higher the percentage, the more similar the sequences are.

[0093] Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith & Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman & Wunsch, *J. Mol. Biol.* 48:443, 1970; Pearson & Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444, 1988; Higgins & Sharp, *Gene*, 73:237-44, 1988; Higgins & Sharp, *CABIOS* 5:151-3, 1989; Corpet et al., *Nuc. Acids Res.* 16:10881-90, 1988; Huang et al. *Computer Appls. in the Biosciences* 8, 155-65, 1992; and Pearson et al., *Meth. Mol. Bio.* 24:307-31, 1994. Altschul et al., *J. Mol. Biol.* 215:403-10, 1990, presents a detailed consideration of sequence alignment methods and homology calculations.

[0094] The NCBI Basic Local Alignment Search Tool (BLAST) (Altschul et al., *J. Mol. Biol.* 215:403-10, 1990) is available from several sources, including the National Cen-

ter for Biological Information (NCBI, National Library of Medicine, Building 38A, Room 8N805, Bethesda, Md. 20894) and on the Internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx. Additional information can be found at the NCBI web site.

[0095] BLASTN is used to compare nucleic acid sequences, while BLASTP is used to compare amino acid sequences. To compare two nucleic acid sequences, the options can be set as follows: -i is set to a file containing the first nucleic acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second nucleic acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastn; -o is set to any desired file name (e.g., C:\output.txt); -q is set to -1; -r is set to 2; and all other options are left at their default setting. For example, the following command can be used to generate an output file containing a comparison between two sequences: C:\B12seq -i c:\seq1.txt -j c:\seq2.txt -p blastn -o c:\output.txt -q -1 -r 2.

[0096] To compare two amino acid sequences, the options of B12seq can be set as follows: -i is set to a file containing the first amino acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second amino acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastp; -o is set to any desired file name (e.g., C:\output.txt); and all other options are left at their default setting. For example, the following command can be used to generate an output file containing a comparison between two amino acid sequences: C:\B12seq -i c:\seq1.txt -j c:\seq2.txt -p blastp -o c:\output.txt. If the two compared sequences share homology, then the designated output file will present those regions of homology as aligned sequences. If the two compared sequences do not share homology, then the designated output file will not present aligned sequences.

[0097] Once aligned, the number of matches is determined by counting the number of positions where an identical nucleotide or amino acid residue is presented in both sequences. The percent sequence identity is determined by dividing the number of matches either by the length of the sequence set forth in the identified sequence, or by an articulated length (e.g., 100 consecutive nucleotides or amino acid residues from a sequence set forth in an identified sequence), followed by multiplying the resulting value by 100. For example, a nucleic acid sequence that has 1166 matches when aligned with a test sequence having 1554 nucleotides is 75.0 percent identical to the test sequence (i.e., $1166 \div 1554 \times 100 = 75.0$). The percent sequence identity value is rounded to the nearest tenth. For example, 75.11, 75.12, 75.13, and 75.14 are rounded down to 75.1, while 75.15, 75.16, 75.17, 75.18, and 75.19 are rounded up to 75.2. The length value will always be an integer. In another example, a target sequence containing a 20-nucleotide region that aligns with 20 consecutive nucleotides from an identified sequence as follows contains a region that shares 75 percent sequence identity to that identified sequence (i.e., $15 \div 20 \times 100 = 75$).

[0098] For comparisons of amino acid sequences of greater than about 30 amino acids, the Blast 2 sequences function is employed using the default BLOSUM62 matrix set to default parameters, (gap existence cost of 11, and a per residue gap cost of 1). Homologs are typically characterized by possession of at least 70% sequence identity counted over the full-length alignment with an amino acid sequence using the NCBI Basic Blast 2.0, gapped blastp with databases such

as the nr or swissprot database. Queries searched with the blastn program are filtered with DUST (Hancock and Armstrong, 1994, Comput. Appl. Biosci. 10:67-70). Other programs use SEG. In addition, a manual alignment can be performed. Proteins with even greater similarity will show increasing percentage identities when assessed by this method, such as at least 75%, 80%, 85%, 90%, 95%, or 99% sequence identity.

[0099] Nucleic acid sequences that do not show a high degree of identity may nevertheless encode identical or similar (conserved) amino acid sequences, due to the degeneracy of the genetic code. Changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid molecules that all encode substantially the same protein. Such homologous nucleic acid sequences can, for example, possess at least 60%, 70%, 80%, 90%, 95%, 98%, or 99% sequence identity determined by this method.

[0100] One of skill in the art will appreciate that these sequence identity ranges are provided for guidance only; it is possible that strongly significant homologs could be obtained that fall outside the ranges provided. Thus, a variant cadA, panD, BAPAT, or HPDH protein or nucleic acid molecule that can be used with the organisms and methods of the present disclosure can have at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to the SEQ ID NOs: and GenBank® Accession Nos. provided herein.

[0101] Transformed: A cell, such as a fungal cell, into which a nucleic acid molecule has been introduced, for example by molecular biology methods. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including, but not limited to chemical methods (e.g., calcium-phosphate transfection), physical methods (e.g., electroporation, microinjection, particle bombardment), fusion (e.g., liposomes), receptor-mediated endocytosis (e.g., DNA-protein complexes, viral envelope/capsid-DNA complexes) and by biological infection by viruses such as recombinant viruses. In one example, the protoplast transformation provide herein, such as in Example 1, is used.

[0102] Vector: A nucleic acid molecule as introduced into a host cell, thereby producing a transformed or recombinant host cell. A vector may include nucleic acid sequences that permit it to replicate in the host cell, such as an origin of replication. A vector may also include a panD, BAPAT, or HPDH coding sequence, or a sequence used to genetically inactivate cadA for example in combination with a promoter, and/or selectable marker genes, and other genetic elements. A vector can transduce, transform or infect a cell, thereby causing the cell to express nucleic acids and/or proteins other than those native to the cell. A vector optionally includes materials to aid in achieving entry of the nucleic acid into the cell, such as a viral particle, liposome, protein coating or the like. In one example, a vector is a plasmid.

Overview

[0103] The filamentous fungus *Aspergillus pseudoterreus* has been used for industrial production of itaconic acid. cis-aconitic acid decarboxylase (cadA) is the key enzyme in itaconic acid production. The itaconic acid biosynthesis cluster is composed of genes tf, mttA, cadA and mfsA. As shown in FIG. 1, itaconic acid (IA) is produced from

glucose. Glucose is utilized in the cell mainly by the glycolytic pathway and metabolized to pyruvate, which forms citric acid. cis-aconitic acid is derived from citric acid as a primary precursor of IA. cis-aconitic acid decarboxylase (cadA) removes carbon dioxide from cis-aconitic acid and forms itaconic acid. However, cadA is localized in the cytosol, while cis-aconitic acid is formed from the TCA cycle in the mitochondria. mttA is localized on the mitochondrial membrane and functioned to transport cis-aconitic acid from mitochondria to cytosol. Another transporter, mfsA is also an organic acid transporter that may be involved in exporting itaconic acid out of cells.

[0104] The first demonstration of genetically inactivating the cadA gene in *Aspergillus pseudoterreus* is shown herein. In the cadA deletion strain (Δ cadA), no more itaconic acid is produced. At the same time significant amount of cis-aconitic acid and trans-aconitic acid are detected. Blocking the itaconic acid production pathway permits the carbon to be diverted towards other organic acid production. The Δ cadA *Aspergillus* can be used as a host for chemical platform, and provides a new way to produce aconitic acid and other organic acids (for example by expressing other genes needed for production of those acids, such as panD, BAPAT, and HPDH for 3-HP production). This strain works as biocatalyst that converts biomass into aconitic acid through bioproduction method at room temperature (such as about 20-35° C.) and ordinary pressure (such as about 1 atm). Current processes of aconitic acid production include chemical synthesis that require high temperatures and harmful reagents.

[0105] The EST data provided herein demonstrated that four genes, tf, cadA, mttA and mfsA show high transcription frequency after IA production starts, but not before IA production begins. The high expression of these genes persists through the production process. Genes upstream and downstream of the cluster did not show expression differences before and after production. One gene downstream next to mfsA, a p450 enzyme, also showed high expression after IA production started, however, deletion of this gene did not effect IA yield.

[0106] Correlations between the IA gene cluster and IA production were further investigated by constructing deletion strains. In a Δ cad strain, no IA was detected, while trace amounts of IA were detected in an mttA knockout. IA production in an mfsA deletion strain decreased one third compared with wild type. This indicates mfsA can transport IA across the cell membrane. In the Δ tf strain, IA production decreased eight fold and slowed the production rate compared to wild type. Also in the tf deletion strain, expression of cadA, mttA and mfsA significantly decreased. RT-PCR results indicated that the expression level of genes in the IA cluster was regulated by tf, which is turned on by IA production conditions.

[0107] The Δ cadA strain produced aconitic acid. During the production, cis-aconitic acid was detected first, followed by the appearance of trans-aconitic acid. cis-aconitic acid levels remained consistent from day 5 forward. The trans-aconitic acid levels continued to increase from days 4 to 10. By day 10, more than 10g/L trans-aconitic acid was detected in the supernatant. In the Δ cadA strain, cis-aconitic acid decarboxylase is not produced, and the cis-aconitic acid cannot be converted to itaconic acid by decarboxylation and accumulates in the cell. cis-aconitic acid was transported

outside the cell. cis-aconitic acid is not stable in the acid solution and is rapidly converted into trans-aconitic acid.

[0108] Aconitic acid is an unsaturated tricarboxylic acid and is noted as a top 30 potential building block by United States Department of Energy (DOE). Trans-aconitic acid can be used to make polymers. Currently, trans-aconitic acid is produced by chemical synthesis and requires high temperature and harmful solvents. Generation of trans-aconitic acid has been achieved by metabolic engineering aconitase isomerase from *Pseudomonas* sp. WU-0701 into *E. coli*. However, the substrate for the recombinant *E. coli* to produce trans-aconitic acid is citric acid, which has to be generated first from fermentation. In contrast, the disclosed Δ cadA fungi can produce trans-aconitic acid directly from renewable biomass substrates. Also since the cadA is not functional and precursors from TAC cycle accumulate in the cell, the carbon can be rerouted to generate other organic acid since *A. pseudoterreus* is industrial filamentous fungi and tolerant to low pH.

[0109] Based on these observations, provided herein are isolated recombinant (i.e., transformed) *Aspergillus* fungi that include a genetic inactivation (also referred to as a functional deletion) of an endogenous cis-aconitic acid decarboxylase (cadA) gene. Such fungi are referred to herein as Δ cadA fungi. Exemplary *Aspergillus* species that can be used include *Aspergillus pseudoterreus* and *Aspergillus terreus*. In some examples, the endogenous cadA gene is genetically inactivated by mutation (such as a complete or partial deletion of the cadA gene) or by insertional mutation (such as by insertion of another nucleic acid molecule into the cadA gene, such as an antibiotic resistance marker).

[0110] In some examples, the cadA gene prior to its genetic inactivation encodes a protein having at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 50 or 52. In some examples, the cadA gene (or its coding sequence) prior to its genetic inactivation comprises at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 49, 51, 59 or 92.

[0111] The disclosed Δ cadA fungi can include other exogenous genes to express proteins needed to permit the fungi to produce other organic acids. For example, the disclosed Δ cadA fungi can further include an exogenous nucleic acid molecule encoding aspartate 1-decarboxylase (panD), an exogenous nucleic acid molecule encoding β -alanine-pyruvate aminotransferase (BAPAT), and an exogenous nucleic acid molecule encoding 3-hydroxypropionate dehydrogenase (HPDH). panD, BAPAT, and HPDH coding sequences can be part of a one or more nucleic acid molecules, such as a vector. In addition, expression of the panD, BAPAT, and HPDH coding sequences can be driven by one or more promoters, such as a bi-directional promoter. In some examples, the promoter is native to the gene it is expressing. In some examples, the promoter is from *A. niger*. In some examples, the panD, BAPAT, and/or HPDH coding sequences are inserted into the cadA gene, genetically inactivating cadA. In some examples, the exogenous nucleic acid molecule encoding panD has at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 53 or 65, and/or encodes a panD protein comprising at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 54. In some examples, the exogenous nucleic acid molecule encoding BAPAT has at least 80%, at least

90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 55, and/or encodes a BAPAT protein having at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 56. In some examples, the exogenous nucleic acid molecule encoding HPDH has at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 57, and/or encodes a HPDH protein having at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 58.

[0112] The disclosure also provides compositions that include the Δ cadA fungi, and the Δ cadA fungi expressing other genes (such as panD, BAPAT, and HPDH). Such a composition can include a solid or liquid culture or growth media, such as complete media, minimal media, or Riscaldati medium (such as modified Riscaldati medium with 20× trace elements).

[0113] The disclosure also provides kits that include the Δ cadA fungi, and the Δ cadA fungi expressing other genes (such as panD, BAPAT, and HPDH). Such a kits can include a solid or liquid culture or growth media, such as complete media, minimal media, or Riscaldati medium (such as modified Riscaldati medium with 20× trace elements).

[0114] Also provided are methods of using the disclosed Δ cadA fungi to make aconitic acid. Such a method can include culturing the recombinant *Aspergillus* Δ cadA fungi under conditions that permit the fungus to make aconitic acid, such as growth in Riscaldati medium, thereby making aconitic acid. In some examples the aconitic acid generated is cis-aconitic acid, trans-aconitic acid, or both. In some examples, the fungi are cultured at room temperature (e.g., 20-35° C.) at normal atmospheric pressure (e.g., 1 atm). In some examples, the method includes purifying or isolating the aconitic acid, for example from the culture media or from the cultured fungus. In some examples, the aconitic acid is isolated at least 2 days, at least 3 days, at least 5 days, at least 8 days or at least 10 days after the start of culturing, such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 days after the start of culturing.

[0115] Also provided are methods of using the disclosed Δ cadA fungi expressing panD, BAPAT, and HPDH to make 3-HP. Such a method can include culturing the recombinant *Aspergillus* Δ cadA fungi expressing panD, BAPAT, and HPDH under conditions that permit the fungus to make 3-HP, such as growth in Riscaldati medium (such as modified Riscaldati medium with 20× trace elements), thereby making 3-HP. In some examples, the fungi are cultured at room temperature (e.g., 20-35° C.) at normal atmospheric pressure (e.g., 1 atm). In some examples, the method includes purifying or isolating the 3-HP, for example from the culture media or from the cultured fungus. In some examples, the 3-HP is isolated at least 2 days, at least 3 days, at least 5 days, at least 8 days or at least 10 days after the start of culturing, such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 days after the start of culturing.

Recombinant Δ cadA Fungi

[0116] The present disclosure provides isolated recombinant *Aspergillus* fungi having its endogenous cadA gene genetically inactivated (e.g., functional deletion) of. Such fungi are referred to herein as Δ cadA fungal strains. It is shown herein that Δ cadA *Aspergillus* strains have increased

aconitic acid production as compared to *Aspergillus* having native levels of cadA expression.

[0117] Any variety or strain of *Aspergillus* can be used. In particular examples, the *Aspergillus* fungus is *A. terreus* or *A. pseudoterreus*, as well as particular strains thereof (for example *A. terreus* NRRL 1960, *A. pseudoterreus* ATCC 32359).

[0118] In addition, any method for genetic inactivation can be used, as long as the expression of the cadA gene is significantly reduced or eliminated, or the function of the cadA protein is significantly reduced or eliminated. In particular examples, the cadA gene is genetically inactivated by complete or partial deletion mutation or by insertional mutation. In some examples genetic inactivation need not be 100%. In some embodiments, genetic inactivation refers to at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95% gene or protein inactivation. The term “reduced” or “decreased” as used herein with respect to a cell and a particular gene or protein activity refers to a lower level of activity than that measured in a comparable cell of the same species. For example, a particular *A. terreus* or *A. pseudoterreus* lacking cadA activity has reduced cadA activity if a comparable *A. terreus* or *A. pseudoterreus* not having an cadA genetic inactivation has detectable cadA activity.

[0119] cadA sequences are disclosed herein and others are publicly available, for example from GenBank or EMBL. In some examples, the cadA gene functionally deleted encoded a protein having at least 80%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 50 or 52 prior to is genetic inactivation. In some examples, the endogenous cadA gene functionally deleted comprises at least 80%, at least 90%, at least 95%, at least 97%, or at least 98% sequence identity to SEQ ID NO: 49, 51, 59, or 92 prior to is genetic inactivation.

[0120] The genetic inactivation of cadA results in many phenotypes in the recombinant Δ cadA *Aspergillus*, such as *A. terreus* or *A. pseudoterreus*. For example, Δ cadA mutants can have one or more of the following phenotypes: produces at least 2-fold, at least 3-fold, at least 3.5 fold, at least 5-fold, at least 8-fold, or at least 10-fold more total aconitic acid than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus* (for example at day 3, 4, 5, 6, 7, 8, 9 or 10 of production); produces at least 2-fold more cis-aconitic acid at day 5, 6, 7, 8, 9, or 10 of culturing in Riscaldati medium than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus*; produces at least 2-fold, at least 3-fold, at least 5-fold, or at least 10-fold more trans-aconitic acid at day 10 of culturing in Riscaldati medium than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus*; or combinations thereof. In some examples, such increases are relative to *Aspergillus terreus* strain ATCC 32359 grown under the same conditions as the Δ cadA mutant. In some examples, an increased total aconitic acid production by Δ cadA fungi occurs at least 3 days (such as at least 4, 5, 6, 7, 8, 9, or 10 days) after inoculation in Riscaldati medium (such as at least 0.5 g/L aconitic acid or at least 1 g/L aconitic acid), as compared to no detectable aconitic acid produced by *Aspergillus terreus* strain ATCC 32359 at the same time point.

[0121] Additional genes can also be inactivated in the Δ cadA fungi, wherein the additional genes may or may not provide additional enhancement of aconitic acid production to the fungus. In one example, the Δ cadA fungi includes overexpressed or upregulated aconitic acid transporters.

[0122] In some examples, Δ cadA fungi include one or more additional exogenous nucleic acid molecules, for example to permit production of other organic acids by the recombinant fungi. In one example, the Δ cadA fungi includes an exogenous nucleic acid molecule encoding aspartate decarboxylase (panD), an exogenous nucleic acid molecule encoding β -alanine-pyruvate aminotransferase (BAPAT), and an exogenous nucleic acid molecule encoding 3-hydroxypropionate dehydrogenase (HPDH). Such exogenous nucleic acid molecules can be part of one or more exogenous nucleic acid molecules (such as 1, 2 or 3 exogenous nucleic acid molecules). In some examples, exogenous nucleic acid molecules can be part of a vector, such as a plasmid or viral vector. In some examples, expression of the exogenous nucleic acid molecules is driven by one or more promoters, such as a constitutive or inducible promoter, or a bi-directional promoter. In some examples, the promoter used to drive expression of panD, BAPAT, and HPDH is a native promoter (e.g., native to the panD, BAPAT, and HPDH gene expressed). In other examples, the promoter used to drive expression of panD, BAPAT, and HPDH is a non-native promoter (e.g., exogenous to the panD, BAPAT, and HPDH gene expressed). In some examples, such a Δ cadA fungi expressing panD, BAPAT, and HPDH are used to produce 3-HP.

[0123] A. Methods of Functionally Deleting cadA

[0124] As used herein, an “inactivated” or “functionally deleted” cadA gene means that the cadA gene has been mutated, for example by insertion, deletion, or substitution (or combinations thereof) of one or more nucleotides such that the mutation substantially reduces (and in some cases abolishes) expression or biological activity of the encoded cadA gene product. The mutation can act through affecting transcription or translation of the cadA gene or its mRNA, or the mutation can affect the cadA polypeptide product itself in such a way as to render it substantially inactive.

[0125] In one example, a strain of *Aspergillus* is transformed with a vector which has the effect of down-regulating or otherwise inactivating a cadA gene. This can be done by mutating control elements such as promoters and the like which control gene expression, by mutating the coding region of the gene so that any protein expressed is substantially inactive, or by deleting the cadA gene entirely. For example, a cadA gene can be functionally deleted by complete or partial deletion mutation (for example by deleting a portion of the coding region of the gene) or by insertional mutation (for example by inserting a sequence of nucleotides into the coding region of the gene, such as a sequence of about 1-5000 nucleotides). In one example, the cadA gene is genetically inactivated by inserting coding sequences for panD, BAPAT, and/or HPDH. Thus, the disclosure provides transformed fungi that include at least one exogenous nucleic acid molecule which genetically inactivates a cadA gene. In one example, such a transformed cell produces more aconitic acid, for example relative to a comparable fungus with a native or wild-type cadA sequence.

[0126] In particular examples, an insertional mutation includes introduction of a sequence that is in multiples of three bases (e.g., a sequence of 3, 9, 12, or 15 nucleotides) to reduce the possibility that the insertion will be polar on downstream genes. For example, insertion or deletion of even a single nucleotide that causes a frame shift in the open reading frame, which in turn can cause premature termination of the encoded cadA polypeptide or expression of a

substantially inactive polypeptide. Mutations can also be generated through insertion of foreign gene sequences, for example the insertion of a gene encoding antibiotic resistance (such as hygromycin or bleomycin), or panD, BAPAT, and/or HPDH coding sequences.

[0127] In one example, genetic inactivation is achieved by deletion of a portion of the coding region of the cadA gene. For example, some, most (such as at least 50%) or virtually the entire coding region can be deleted. In particular examples, about 5% to about 100% of the gene is deleted, such as at least 20% of the gene, at least 40% of the gene, at least 75% of the gene, or at least 90% of the cadA gene.

[0128] Deletion mutants can be constructed using any of a number of techniques. In one example, homologous double crossover with fusion PCR products is employed to genetically inactivate one or more genes in *Aspergillus*. A specific example of such a method is described in Example 1 below.

[0129] In one example, a strategy using counterselectable markers can be employed which has been utilized to delete genes. For a review, see Reyrat et al. (*Infect. Immun.* 66:4011-4017, 1998). In this technique, a double selection strategy is employed wherein a plasmid is constructed encoding both a selectable and counterselectable marker, with flanking DNA sequences derived from both sides of the desired deletion. The selectable marker is used to select for fungi in which the plasmid has integrated into the genome in the appropriate location and manner. The counterselectable marker is used to select for the very small percentage of fungi that have spontaneously eliminated the integrated plasmid. A fraction of these fungi will then contain only the desired deletion with no other foreign DNA present.

[0130] In another technique, the cre-lox system is used for site specific recombination of DNA (for example see Steiger et al., *Appl. Environ. Microbiol.* 77(1):114, 2011). The system includes 34 base pair lox sequences that are recognized by the bacterial cre recombinase gene. If the lox sites are present in the DNA in an appropriate orientation, DNA flanked by the lox sites will be excised by the cre recombinase, resulting in the deletion of all sequences except for one remaining copy of the lox sequence. Using standard recombination techniques, the targeted gene of interest (e.g., cadA) can be deleted in the *Aspergillus* genome and to replace it with a selectable marker (for example a gene coding for kanamycin resistance) that is flanked by the lox sites. Transient expression (by electroporation of a suicide plasmid containing the cre gene under control of a promoter that functions in *Aspergillus*) of the cre recombinase should result in efficient elimination of the lox flanked marker. This process will produce a mutant containing the desired deletion mutation and one copy of the lox sequence.

[0131] In another method, a cadA gene sequence in the *Aspergillus* genome is replaced with a marker gene, such as green fluorescent protein, β -galactosidase, or luciferase. In this technique, DNA segments flanking a desired deletion are prepared by PCR and cloned into a suicide (non-replicating) vector for *Aspergillus*. An expression cassette, containing a promoter active in *Aspergillus* and the appropriate marker gene, is cloned between the flanking sequences. The plasmid is introduced into wild-type *Aspergillus*. Fungi that incorporate and express the marker gene are isolated and examined for the appropriate recombination event (replacement of the wild type cadA gene with the marker gene).

[0132] Thus, for example, a fungal cell can be engineered to have a disrupted *cadA* gene using common mutagenesis or knock-out technology. (Methods in Yeast Genetics (1997 edition), Adams, Gottschling, Kaiser, and Sterns, Cold Spring Harbor Press, 1998; Datsenko and Wanner, *Proc. Natl. Acad. Sci. USA* 97: 6640-5, 2000; and Dai et al., *Appl. Environ. Microbiol.* 70(4):2474-85, 2004). Alternatively, antisense technology can be used to reduce or eliminate the activity of *cadA*. For example, a fungal cell can be engineered to contain a cDNA that encodes an antisense molecule that prevents *cadA* from being translated. The term "antisense molecule" encompasses any nucleic acid molecule or nucleic acid analog (e.g., peptide nucleic acids) that contains a sequence that corresponds to the coding strand of an endogenous *cadA* gene. An antisense molecule also can have flanking sequences (e.g., regulatory sequences).

[0133] Thus, antisense molecules can be ribozymes or antisense oligonucleotides. A ribozyme can have any general structure including, without limitation, hairpin, hammerhead, or axehead structures, provided the molecule cleaves RNA. Further, gene silencing can be used to reduce the activity of *cadA*.

[0134] In one example, to genetically inactivate *cadA* in *A. pseudoterreus* or *A. terreus*, protoplast transformation is used, for example as described in Example 1. For example, conidia of *A. pseudoterreus* or *A. terreus* are grown in liquid complete medium at room temperature (e.g., about 20-35° C., such as 30° C.) and grown for at least 12 hours (such as at least 16 hours, or at least 18 hours, such as 12-24 hours, or 16-18 hours), at least 100 rpm, such as at least 150 rpm, for example 100 to 200 rpm. The resulting mycelia are subsequently harvested, for example by filtration. Protoplasts are prepared, for example by treating the harvested mycelia with a lysing enzyme (for example in an osmotic wash buffer for at least 30 min, at least 60 min, at least 120 min, or at leave 240 min, such as 2 h). The resulting protoplasts are collected (e.g., by filtering). Protoplasts can be washed, for example with a Washing Solution (0.6M KCl, 0.1M Tris/HCl, pH 7.0) and Conditioning Solution (0.6M KCl, 50 mM CaCl₂, 10 mM Tris/HCl, pH 7.5). The protoplasts are transformed, for example in the conditioning solution. In some examples, at least 0.5 ug, at least 1 ug, or at least 2 ug of DNA (such as 1-2 ug DNA) is added to at least 10⁶ protoplasts (such as at least 10⁷ or 2×10⁷ protoplasts). Polyethylene glycol (PEG), such as PEG8000 is added (such as 25% PEG8000, 0.6M KCl, 50 mM CaCl₂, 10 mM Tris/HCl, and pH 7.5) and the reaction incubated for at least 5 min (such as at least 10 min, at least 20 min, or at least 30 min, such as 10-30 min, 15-20 min, or 20 min) on ice. Additional PEG solution can be added and the reaction incubated for at least 1 min, at least 3 min, or at least 5 min, on ice. Conditioning Solution is added to the reaction, and the protoplast suspension mixed with warm selection agar (Minimal media+0.6M KCl+1.5% Agar+100 ug/ml hygromycin) (such as at 50° C.), and poured directly onto petri dish plates and allowed to solidify. Solidified plates can be inverted and incubated overnight at room temperature (e.g., about 20-35° C., such as 30° C.). The following day, the plates can be overlaid with Minimal Medium containing a selection antibiotic, such as hygromycin. Colonies appear after 3-4 days. Transformants can be excised and transferred to MM plate containing the selection antibiotic.

[0135] B. Measuring Gene Inactivation

[0136] A fungus having an inactivated *cadA* gene can be identified using known methods. For example, PCR and nucleic acid hybridization techniques, such as Northern and Southern analysis, can be used to confirm that a fungus has a genetically inactivated *cadA* gene. In one example, real-time reverse transcription PCR (qRT-PCR) is used for detection and quantification of targeted messenger RNA, such as mRNA of *cadA* gene in the parent and mutant strains as grown at the same culture conditions. Immunohisto-chemical and biochemical techniques can also be used to determine if a cell expresses *cadA* by detecting the expression of the *cadA* peptide encoded by *cadA*. For example, an antibody having specificity for *cadA* can be used to determine whether or not a particular fungus contains a functional nucleic acid encoding *cadA* protein. Further, biochemical techniques can be used to determine if a cell contains a *cadA* gene inactivation by detecting a product produced as a result of the lack of expression of the peptide. For example, production of aconitic acid by *A. terreus* or *A. pseudoterreus* can indicate that such a fungus contains an inactivated *cadA* gene.

[0137] C. Measuring Aconitic Acid Production

[0138] Methods of determining whether a genetic inactivation of *cadA* in *Aspergillus*, such as *A. terreus* or *A. pseudoterreus*, increases aconitic acid production, for example relative to the same strain of *A. terreus* or *A. pseudoterreus* with a native *cadA* sequence (such as a parental strain), are provided herein. Although particular examples are disclosed herein, the methods are not limiting.

[0139] For example, production of aconitic acid by *Aspergillus* (such as a $\Delta cadA$ strain) can be measured using a spectrophotometric assay, by liquid chromatography (LC), or high-pressure liquid chromatography (HPLC) methods. In some examples, the supernatant of the fungus is analyzed for the presence of aconitic acid. In some examples, the culture media containing the $\Delta cadA$ strain is filtered prior to measuring aconitic acid in the culture media (supernatant).

[0140] D. *cadA* Sequences

[0141] *cadA* protein and nucleic acid sequences are publicly available and specific examples are provided herein. In addition, *cadA* sequences can be identified using molecular biology methods.

[0142] Examples of *cadA* nucleic acid sequences are shown in SEQ ID NOS: 49, 51, 59 and 92. However, the disclosure also encompasses variants of SEQ ID NOS: 49, 51, 59 and 92 which encode a functional *cadA* protein. One skilled in the art will understand variants of the *cadA* nucleic acid sequences provided herein can be genetically inactivated. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). In addition, the degeneracy of the code permits multiple nucleic acid sequences to encode the same protein. Such variant *cadA* nucleic acid molecules can share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to any *cadA* nucleic acid sequence, such as SEQ ID NO: 49, 51, 59 or 92.

[0143] Examples of *cadA* protein sequences are shown in SEQ ID NOS: 50 and 52. However, the disclosure also encompasses variants SEQ ID NOS: 50 and 52 which retain *cadA* activity. One skilled in the art will understand that variants of these *cadA* enzyme sequences can be inactivated. Variant sequences can be identified, for example by aligning

known *cadA* sequences. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). Such *cadA* peptides share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to a *cadA* protein sequence, such as SEQ ID NO: 50 or 52.

[0144] In some examples, a *cadA* sequence that is to be genetically inactivated encodes or includes one or more conservative amino acid substitutions. A conservative amino acid substitution is a substitution of one amino acid (such as one found in a native sequence) for another amino acid having similar biochemical properties. Typically, conservative substitutions have little to no impact on the activity of a resulting peptide. In one example, a *cadA* sequence (such as SEQ ID NO: 50 or 52) includes one or more amino acid substitutions, such as conservative substitutions (for example at 1, 2, 5 or 10 residues). Examples of amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative substitutions include: Ser for Ala; Lys for Arg; Gln or His for Asn; Glu for Asp; Ser for Cys; Asn for Gln; Asp for Glu; Pro for Gly; Asn or Gln for His; Leu or Val for Ile; Ile or Val for Leu; Arg or Gln for Lys; Leu or Ile for Met; Met, Leu or Tyr for Phe; Thr for Ser; Ser for Thr; Tyr for Trp; Trp or Phe for Tyr; and Ile or Leu for Val. Further information about conservative substitutions can be found in, among other locations in, Ben-Bassat et al., (*J. Bacteriol.* 169:751-7, 1987), O'Regan et al., (*Gene* 77:237-51, 1989), Sahin-Toth et al., (*Protein Sci.* 3:240-7, 1994), Hochuli et al., (*Bio/Technology* 6:1321-5, 1988), WO 00/67796 (Curd et al.) and in standard textbooks of genetics and molecular biology.

[0145] The *cadA* gene inactivated in a fungus, in particular examples, includes a sequence that encodes a *cadA* protein having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a *cadA* protein sequence, such as SEQ ID NO: 50 or 52, wherein the protein can catalyze the decarboxylation of cis-aconitate to itaconate and CO₂ and vice versa. In a specific example, the *cadA* gene inactivated in a fungus encodes a *cadA* protein shown in SEQ ID NO: 50 or 52.

[0146] The *cadA* gene that is to be inactivated in a fungus, in particular examples, includes a sequence (such as a coding sequence) having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a *cadA* nucleic acid sequence, such as SEQ ID NO: 49, 51, 59, or 92, and encodes a *cadA* protein that can catalyze the decarboxylation of cis-aconitate to itaconate and CO₂ and vice versa. In a specific example, *cadA* gene inactivated in a fungus is the sequence of SEQ ID NO: 2 or 4.

[0147] One skilled in the art will appreciate that additional *cadA* sequences can be identified. For example, *cadA* nucleic acid molecules that encode a *cadA* protein can be identified and obtained using molecular cloning or chemical nucleic acid synthesis procedures and techniques, including PCR. In addition, nucleic acid sequencing techniques and software programs that translate nucleic acid sequences into amino acid sequences based on the genetic code can be used to determine whether or not a particular nucleic acid has any sequence homology with known *cadA* sequences. Sequence

alignment software such as MEGALIGN (DNASTAR, Madison, Wis., 1997) can be used to compare various sequences.

[0148] In addition, nucleic acid hybridization techniques can be used to identify and obtain a nucleic acid molecule that encodes a *cadA* protein. Briefly, any known *cadA* nucleic acid molecule, or fragment thereof, can be used as a probe to identify similar nucleic acid molecules by hybridization under conditions of moderate to high stringency. Such similar nucleic acid molecules then can be isolated, sequenced, and analyzed to determine whether the encoded protein is a *cadA* protein.

[0149] E. *panD*, BAPAT, and HPDH Sequences

[0150] *panD*, BAPAT, and HPDH protein and nucleic acid sequences are publicly available and specific examples are provided herein. In addition, *panD*, BAPAT, and HPDH sequences can be identified using molecular biology methods.

[0151] Exemplary of *panD* coding sequences are shown in SEQ ID NO: 53 and 65. However, the disclosure also encompasses variants of SEQ ID NO: 53 and 65 which encode a functional *panD* protein. Exemplary of BAPAT coding sequences are shown in SEQ ID NO: 55 and 71. However, the disclosure also encompasses variants of SEQ ID NO: 55 and 71 which encode a functional BAPAT protein. Exemplary of HPDH coding sequences are shown in SEQ ID NO: 57 and 80. However, the disclosure also encompasses variants of SEQ ID NO: 57 and 80 which encode a functional HPDH protein.

[0152] One skilled in the art will understand variants of the *panD*, BAPAT, and HPDH nucleic acid sequences provided herein can be introduced into an *Aspergillus* fungus, such as one that is Δ *cadA*, such as inserting *panD*, BAPAT, and HPDH expression sequences into the native *cadA* gene to inactivate it. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). In addition, the degeneracy of the code permits multiple nucleic acid sequences to encode the same protein. In some examples, a *panD*, BAPAT, and HPDH sequence that is to be expressed in an *Aspergillus* fungus is codon optimized for expression in *Aspergillus*, such as *Aspergillus terreus* or *pseudoterreus*. Such variant *panD*, BAPAT, and HPDH nucleic acid molecules in some examples share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to any *panD*, BAPAT, and HPDH nucleic acid sequence, such as SEQ ID NO: 53, 55, or 57, respectively, or SEQ ID NO: 65, 71, or 80, respectively.

[0153] Exemplary *panD*, BAPAT, and HPDH protein sequences are shown in SEQ ID NOS: 54, 56, and 58, respectively. However, the disclosure also encompasses variants SEQ ID NOS: 54, 56, and 58 which retain *panD*, BAPAT, and HPDH activity, respectively. One skilled in the art will understand that variants of these *panD*, BAPAT, and HPDH sequences can be expressed in an *Aspergillus* fungus, such as one that is Δ *cadA*. Variant sequences can be identified, for example by aligning known *panD*, BAPAT, and HPDH sequences. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). Such *panD*, BAPAT, and HPDH pep-

tides expressed in a Δ cadA fungus in some examples share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to a panD, BAPAT, and HPDH protein sequence, such as SEQ ID NO: 54, 56, or 58, respectively.

[0154] In some examples, a panD, BAPAT, and HPDH sequence that is to be expressed in an *Aspergillus* fungus encodes or includes one or more conservative amino acid substitutions. In one example, a panD, BAPAT, or HPDH sequence (such as SEQ ID NO: 54, 56, or 58, respectively) includes one or more amino acid substitutions, such as conservative substitutions (for example at 1, 2, 5, or 10 residues). Examples of conservative substitutions are provided elsewhere herein.

[0155] The panD, BAPAT, and HPDH gene expressed in a fungus, in particular examples, includes a sequence that encodes a panD, BAPAT, and HPDH protein having at least at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a panD, BAPAT, and HPDH protein sequence, such as SEQ ID NO: 54, 56, or 58, respectively, wherein the variant protein has the biological activity of panD, BAPAT, or HPDH, respectively. In a specific example, the panD, BAPAT, and HPDH gene expressed in a Δ cadA fungus encodes the protein shown in SEQ ID NO: 54, 56, and 58, respectively.

[0156] One skilled in the art will appreciate that additional panD, BAPAT, and HPDH sequences can be identified. For example, panD, BAPAT, and HPDH nucleic acid molecules that encode a panD, BAPAT, and HPDH protein, respectively can be identified and obtained using molecular cloning or chemical nucleic acid synthesis procedures and techniques, including PCR. In addition, nucleic acid sequencing techniques and software programs that translate nucleic acid sequences into amino acid sequences based on the genetic code can be used to determine whether or not a particular nucleic acid has any sequence homology with panD, BAPAT, or HPDH sequences. Sequence alignment software such as MEGALIGN (DNASTAR, Madison, Wis., 1997) can be used to compare various sequences.

[0157] In addition, nucleic acid hybridization techniques can be used to identify and obtain a nucleic acid molecule that encodes a panD, BAPAT, or HPDH protein. Briefly, any known panD, BAPAT, or HPDH nucleic acid molecule, or fragment thereof, can be used as a probe to identify similar nucleic acid molecules by hybridization under conditions of moderate to high stringency. Such similar nucleic acid molecules then can be isolated, sequenced, and analyzed to determine whether the encoded protein is a panD, BAPAT, or HPDH protein.

[0158] In one example, exogenous panD, BAPAT, and/or HPDH nucleic acid sequences are introduced into *A. pseudoterreus* or *A. terreus* using protoplast transformation, for example as described in Example 1 (and described above).

[0159] F. Methods of Increasing panD, BAPAT, and HPDH Expression

[0160] In some examples, a native *A. pseudoterreus* or *A. terreus* fungi does not have or express panD, BAPAT, and/or HPDH nucleic acid sequences. Thus, in some examples, expression of these genes is increased by introducing panD, BAPAT, and/or HPDH nucleic acid coding sequences (such as may be codon optimized) into the *A. pseudoterreus* or *A. terreus* fungi.

[0161] In some examples, a native *A. pseudoterreus* or *A. terreus* fungi does express native panD, BAPAT, and/or HPDH nucleic acid sequences. Thus, in some examples, expression of these genes is upregulated by introducing additional copies of panD, BAPAT, and/or HPDH nucleic acid coding sequences (such as may be codon optimized) into the *A. pseudoterreus* or *A. terreus* fungi. As used herein, “up-regulated” gene means that expression of the gene or gene product (e.g., protein) has been up-regulated, for example by introduction of additional copies of the appropriate gene or coding sequence into the fungus (or other molecular biology methods), such that the introduced nucleic acid sequence is expressed, resulting in increased expression or biological activity of the encoded gene product. In some embodiments, introduction of one or more transgenes including panD, BAPAT, and/or HPDH coding sequences into a native *A. pseudoterreus* or *A. terreus* fungi increases expression of panD, BAPAT, and/or HPDH by at least 20%, at least 40%, at least 50%, at least 100%, at least 150%, at least 200%, at least 300%, or at least 500%, for example relative to the parental fungal strain without the introduced panD, BAPAT, and/or HPDH coding sequences. The term “increased” or “up-regulated” as used herein with respect to a cell and a particular gene or protein activity refers to a higher level of activity than that measured in a comparable cell of the same species. For example, a particular fungi having increased or up-regulated panD, BAPAT, and/or HPDH activity has increased panD, BAPAT, and/or HPDH activity if a comparable fungi having native panD, BAPAT, and/or HPDH activity has less detectable panD, BAPAT, and/or HPDH activity (for example as measured by gene or protein expression).

[0162] In one example, a strain of *Aspergillus* is transformed with a vector which has the effect of up-regulating a panD, BAPAT, and/or HPDH gene (such as a native or non-native panD, BAPAT, and/or HPDH gene). This can be done by introducing one or more panD, BAPAT, and/or HPDH coding sequences (such as a gene sequence), whose expression is controlled by elements such as promoters and the like which control gene expression, by introducing a nucleic acid sequence which itself (or its encoded protein) can increase panD, BAPAT, and/or HPDH protein activity in the fungus, or by introducing another molecule (such as a protein or antibody) increases panD, BAPAT, and/or HPDH protein activity in the fungus. For example, a panD, BAPAT, and/or HPDH gene can be up-regulated by introduction of a vector that includes one or more panD, BAPAT, and/or HPDH sequences (such as 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 panD, BAPAT, and/or HPDH sequences or copies of such sequences) into the desired fungus. In some examples, such panD, BAPAT, and/or HPDH sequences are from different fungal species, can be multiple copies from a single species, or combinations thereof, such as panD, BAPAT, and/or HPDH sequences from at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 different fungal species. In some examples, the panD, BAPAT, and/or HPDH sequence(s) introduced into the fungus is optimized for codon usage. Thus, the disclosure in some examples provides transformed fungi that include at least one exogenous nucleic acid molecule which includes a panD, BAPAT, and/or HPDH gene or coding sequence (such as a nucleic acid sequence encoding SEQ ID NO: 54, 56, or 58, respectively), for example in combination with Δ cadA.

In one example, such transformed cells produce more 3HP, for example relative to a comparable fungus with a native *cadA*.

[0163] In one example, the cre-lox system is used for site specific recombination of DNA (for example see Steiger et al., *Appl. Environ. Microbiol.* 77(1):114, 2011). Using recombination techniques, the targeted gene of interest (e.g., *cadA*) can be deleted in the *Aspergillus* genome and replaced with one or more copies of a non-native *panD*, *BAPAT*, and/or *HPDH* sequence (for example in *A. terreus*, replacing one or both *A. terreus cadA* sequences with *panD*, *BAPAT*, and/or *HPDH* sequences from *A. nidulans* or *A. flavus*) flanked by the lox sites. Transient expression (by electroporation of a suicide plasmid containing the cre gene under control of a promoter that functions in *Aspergillus*) of the cre recombinase should result in efficient elimination of the lox flanked marker. This process will produce a fungus containing the desired insertion mutation and one copy of the lox sequence.

[0164] In one example, a transgene is generated and expressed in the desired fungal cell, such as an Δ *cadA* fungal cell, to increase *panD*, *BAPAT*, and *HPDH* expression. For example, one or more transgenes can include a *panD*, *BAPAT*, and *HPDH* genomic or cDNA sequence (such as one having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to any *panD*, *BAPAT*, and *HPDH* sequence provided herein), for example operably linked to one or more promoters, such as *gpdA* and *eno1*. In one example, the promoter has at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 74 and/or 77. In some examples, the transgene further includes a *trpC* transcriptional terminator sequence of *A. nidulans*, for example downstream of the *panD*, *BAPAT*, and/or *HPDH* sequence. As an alternative to *trpC*, other transcriptional terminators can be used, such as promoters which include a transcriptional terminators (e.g., *ArsA7*, *ArsA37*, polyubiquitin (*ubi4*)). In one example, the *trpC* transcriptional terminator has at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 83 or 86. In one example, the *trpC* transcriptional terminator comprises or consists of the sequence shown in SEQ ID NO: 83 or 86. In some examples, the transgene further includes a *ptrA* sequence, for example downstream of the *trpC* transcriptional terminator sequence. As an alternative to *ptrA*, the bleomycin gene or *bar* gene can be used. In one example, the *ptrA* sequence has at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 89. In one example, the *ptrA* sequence comprises or consists of the sequence shown in SEQ ID NO: 89.

[0165] In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 59, 62, 65, 68, 71, and/or 74. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 59, 62, 65, 68, 71, and/or 74.

[0166] In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 77, 80, and/or 83. In one

example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 77, 80, and/or 83.

[0167] In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 86, 89, and/or 92. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 86, 89, and/or 92.

[0168] In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 89 and/or 92. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 89 and/or 92.

[0169] In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 95 and/or 97. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 95 and/or 97.

[0170] In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 59, 62, 65, 68, 71, 74, 77, 80, 83, 86, 89, and/or 92. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 59, 62, 65, 68, 71, 74, 77, 80, 83, 86, 89, and/or 92.

[0171] In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 59, 62, 65, 68, 71, 74, 77, 80, 83, 89, and/or 92. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 59, 62, 65, 68, 71, 74, 77, 80, 83, 89, and/or 92.

[0172] G. Measuring Gene Expression

[0173] A Δ *cadA* fungus expressing *panD*, *BAPAT*, and/or *HPDH* can be identified using known methods. For example, PCR and nucleic acid hybridization techniques, such as Northern, RT-PCR, and Southern analysis, can be used to confirm that a fungus expresses *panD*, *BAPAT*, and/or *HPDH* such as an increase in the *panD*, *BAPAT*, and/or *HPDH* copy number. Immunohisto-chemical and biochemical techniques can also be used to determine if a cell expresses *panD*, *BAPAT*, and/or *HPDH* by detecting the expression of the *panD*, *BAPAT*, and/or *HPDH* peptide encoded by *panD*, *BAPAT*, and/or *HPDH*. For example, an antibody having specificity for *panD*, *BAPAT*, and/or *HPDH* can be used to determine whether or not a particular fungus has increased *panD*, *BAPAT*, and/or *HPDH* protein expression, respectively. Further, biochemical techniques can be used to determine if a cell has increased *panD*, *BAPAT*, and/or *HPDH* expression by detecting a product produced as a result of the expression of the peptide. For example, production of 3-HP by Δ *cadA A. terreus* or *A. pseudoterreus* can indicate that such a fungus expresses *panD*, *BAPAT*, and *HPDH*.

[0174] H. Measuring 3-HP Production

[0175] Methods of determining whether a genetic inactivation of *cadA* in combination with expression of *panD*, *BAPAT*, and *HPDH* in *Aspergillus* increases 3-HP production, for example relative to the same strain with a native *cadA* sequence, (such as a parental strain) include HPLC.

[0176] Methods of Producing Aconitic Acid

[0177] The recombinant Δ cadA fungi can be used to produce aconitic acid (for example for as a building block for other materials, such as polymers). Such fungi can be from any *Aspergillus* species, such as *Aspergillus terreus* or *pseudoterreus*. For example, the disclosure provides methods of making aconitic acid (such as cis-aconitic acid, trans-aconitic acid, or both), which can include culturing Δ cadA fungi under conditions that permit the fungus to make aconitic acid, for example in Riscaldati medium.

[0178] In some examples, the fungi are cultured at room temperature (e.g., 20-35° C.) at normal atmospheric pressure (e.g., 1 atm). In some examples, the method includes purifying or isolating the aconitic acid, for example from the culture media or from the cultured fungus. In some examples, the aconitic acid is isolated at least 2 days, at least 3 days, at least 5 days, at least 8 days or at least 10 days after the start of culturing, such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 days after the start of culturing.

[0179] Methods of making aconitic acid include culturing Δ cadA fungi under conditions that permit the fungus to make aconitic acid. In general, the culture media and/or culture conditions can be such that the fungi grow to an adequate density and produce aconitic acid efficiently. In one example the Δ cadA fungi are cultured or grown in an acidic liquid medium, such as Riscaldati medium (100 g Glucose, 0.11 g KH_2PO_4 , 2.36 g $(\text{NH}_4)_2\text{SO}_4$, 2.08 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.074 g NaCl, 0.13 g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1 ml of 1000x trace elements in 1000 ml DI water, adjust pH to 3.4 with H_2SO_4 , 1000x trace elements contains 1.3g/L $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 5.5g/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.7 g/L $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$). In one example the Δ cadA fungi are cultured or grown in a liquid medium having an initial pH of less than 4, such as less than 3.5, for example about pH 3 to 4, 3.5 to 4, 3.3 to 3.5, for example pH 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9 or 4. In some examples the Δ cadA fungi are cultured or grown in a liquid Riscaldati medium at about 20 to 35° C. (such as 20° C. to 30° C., 25° C. to 30° C., 28 to 32° C., or 30° C.) with rotation (such as at least 100 rpm, at least 120 rpm, such as 150 rpm) at normal pressure.

[0180] In one example, the fungi are grown in culture containers (such as baffled flasks, and in some examples are silanized (5% solution of dichlorodimethylsilane in heptane (Sigma, St. Louis, Mo.)). Each culture container is inoculated with spores (such as at least 10^6 spores/ml [agree?]) and incubated for at least 3 days, at least 4 days, at least 5 days, or at least 10 days at 30° C. and 100 to 200 rpm to obtain aconitic acid.


[0181] In one example, the Δ cadA fungi produce more aconitic acid than a corresponding fungus with wild-type cadA. In specific examples, the Δ cadA fungi produce at least 1 g/l of total aconitic acid after 4 days, for example at least 2 g/l, at least 3 g/l, at least 4 g/l, at least 5 g/l, at least 6 g/l, at least 7 g/l, at least 8 g/l, at least 9 g/l or at least 10 g/l after at least 5 days, at least 6 days, at least 7 days, at least 8 days, or at least 10 days, such as after 4 to 6 days, 8 to 10 days, or 4 to 5 days) when grown in Riscaldati medium at 30° C. with 150 rpm shaking. In specific examples, the Δ cadA fungi produce at least 1 g/l of cis-aconitic acid after 4 days, for example at least 2 g/l after at least 5 days, at least 6 days, at least 7 days, at least 8 days, or at least 10 days, such as after 4 to 6 days, 8 to 10 days, or 4 to 5 days when grown in Riscaldati medium at 30° C. with 150 rpm shaking. In

specific examples, the Δ cadA fungi produce at least 1 g/l of trans-aconitic acid after 6 days, for example at least 2 g/l, at least 3 g/l, at least 4 g/l, at least 5 g/l, at least 6 g/l, at least 7 g/l, at least 8 g/l, at least 9 g/l or at least 10 g/l after at least 7 days, at least 8 days, or at least 10 days, such as after 6 to 12 days, 5 to 10 days, or 6 to 10 days) when grown in Riscaldati medium at 30° C. with 150 rpm shaking.

[0182] In some examples, the method further includes isolating the aconitic acid made by the Δ cadA fungi. Once produced, any method can be used to isolate the aconitic acid. For example, separation techniques (such as filtration) can be used to remove the fungal biomass from the culture medium, and isolation procedures (e.g., filtration, distillation, precipitation, electrodialysis, and ion-exchange procedures) can be used to obtain the aconitic acid from the broth (such as a fungi-free broth). In addition, the aconitic acid can be isolated from the culture medium after the aconitic acid production phase has been terminated.

Methods of Producing 3-HP

[0183] The recombinant Δ cadA fungi that also express panD, BAPAT, and HPDH can be used to produce 3-HP



[0184] (for example for as a building block for other materials, such as acrylonitrile, acrylic acid by dehydration, malonic acid by oxidation, esters by esterification reactions with alcohols, and reduction to 1,3 propanediol). Such fungi can be from any *Aspergillus* species, such as *Aspergillus terreus* or *pseudoterreus*. For example, the disclosure provides methods of making 3-HP, which can include culturing Δ cadA fungi that also express panD, BAPAT, and HPDH under conditions that permit the fungus to make 3-HP, for example in Riscaldati medium (such as modified Riscaldati medium with 20x trace elements).

[0185] In some examples, the Δ cadA fungi that also express panD, BAPAT, and HPDH are cultured at room temperature (e.g., 20-35° C.) at normal atmospheric pressure (e.g., 1 atm). In some examples, the method includes purifying or isolating the 3-HP, for example from the culture media or from the cultured fungus. In some examples, the 3-HP is isolated at least 2 days, at least 3 days, at least 5 days, at least 7 days, at least 8 days or at least 10 days after the start of culturing, such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 days after the start of culturing.

[0186] Methods of making 3-HP include culturing Δ cadA fungi that also express panD, BAPAT, and HPDH under conditions that permit the fungus to make 3-HP. In general, the culture media and/or culture conditions can be such that the fungi grow to an adequate density and produce 3-HP efficiently. In one example the Δ cadA fungi that also express panD, BAPAT, and HPDH are cultured or grown in an acidic liquid medium, such as Riscaldati medium (100 g Glucose, 0.11 g KH_2PO_4 , 2.36 g $(\text{NH}_4)_2\text{SO}_4$, 2.08g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.074g NaCl, 0.13g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1 ml of 1000x trace elements in 1000 ml DI water, adjust pH to 3.4 with H_2SO_4 , 1000x trace elements contains 1.3 g/L $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 5.5

g/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.7 g/L $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, which may include 20× trace elements). In one example the ΔcadA fungi are cultured or grown in a liquid medium having an initial pH of less than 4, such as less than 3.5, for example about pH 3 to 4, 3.5 to 4, 3.3 to 3.5, for example pH 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9 or 4. In some examples the ΔcadA fungi that also express panD, BAPAT, and HPDH are cultured or grown in a liquid modified Riscaldati medium with 20× trace elements at about 20 to 35° C. (such as 20° C. to 30° C., 25° C. to 30° C., 28 to 32° C., or 30° C.) with rotation (such as at least 100 rpm, at least 120 rpm, such as 150 or 200 rpm) at normal pressure.

[0187] In one example, the fungi are grown in culture containers (such as baffled flasks, and in some examples are silanized (5% solution of dichlorodimethylsilane in heptane (Sigma, St. Louis, Mo.)). Each culture container is inoculated with spores (such as at least 10⁶ spores/ml) and incubated for at least 3 days, at least 4 days, at least 5 days, or at least 10 days at 30° C. and 100 to 300 rpm (such as 150 or 200 rpm) to obtain 3-HP.

[0188] In one example, the cadA fungi that also express panD, BAPAT, and HPDH produce more 3-HP than a corresponding fungus with wild-type cadA (either with or without panD, BAPAT, and HPDH expression). In specific examples, the ΔcadA fungi that also express panD, BAPAT, and HPDH produce at least 0.1 g/l of 3-HP after at least 4 days, for example at least 0.2 g/l at least 0.25 g/l at least 0.3 g/l, at least 0.4 g/l, at least 0.5 g/l at least 0.6 g/l at least 0.7 g/l at least 0.8 g/l, at least 0.9 g/l, at least 1.1 g/l, at least 1.2 g/l, at least 1.5 g/l, or at least 1.6 g/l after at least 5 days, at least 6 days, at least 7 days, at least 8 days, or at least 10 days, such as after 4 to 6 days, 8 to 10 days, or 4 to 5 days, when grown in Riscaldati medium (such as modified Riscaldati medium with 20× trace elements) at 30° C. with 150 rpm shaking.

[0189] In some examples, the method further includes isolating the 3-HP made by the ΔcadA fungi. Once produced, any method can be used to isolate the 3-HP. For example, separation techniques (such as filtration) can be used to remove the fungal biomass from the culture medium, and isolation procedures (e.g., filtration, distillation, precipitation, electrodialysis, and ion-exchange procedures) can be used to obtain the 3-HP from the broth (such as a fungi-free broth). In addition, the 3-HP can be isolated from the culture medium after the 3-HP production phase has been terminated.

Compositions and Kits

[0190] Also provided by the present disclosure are compositions that include isolated ΔcadA fungi (which in some examples also express panD, BAPAT, and HPDH, such as exogenous panD, BAPAT, and HPDH proteins), such as a medium for culturing, storing, or growing the fungus. In some examples, the ΔcadA fungi and ΔcadA fungi which express panD, BAPAT, and HPDH in the composition are freeze dried or lyophilized.

[0191] Also provided by the present disclosure are kits that include isolated ΔcadA fungi (which in some examples also express panD, BAPAT, and HPDH, such as exogenous panD, BAPAT, and HPDH proteins), such as a kit that includes a medium for culturing, storing, or growing the

fungus. In some examples, the ΔcadA fungi and ΔcadA fungi which express panD, BAPAT, and HPDH in the kit are freeze dried or lyophilized.

[0192] Exemplary mediums include that can be in the disclosed compositions and kits include solid medium (such as those containing agar, for example complete medium (CM) or minimal medium (MM)) and liquid media (such as a fermentation broth, such as CM, MM, or CAP medium). In one example, the kit or composition includes Riscaldati medium (100 g Glucose, 0.11 g KH_2PO_4 , 2.36 g $(\text{NH}_4)_2\text{SO}_4$, 2.08 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.074 g NaCl, 0.13 g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1 ml of 1000× trace elements in 1000 ml DI water, adjust pH to 3.4 with H_2SO_4 , 1000× trace elements contains 1.3 g/L $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 5.5 g/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.7 g/L $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$), for example

	Conc. (g/L)	Amount	Notes
Glucose	100	100 g	
KH_2PO_4	0.11	0.11 g	
$(\text{NH}_4)_2\text{SO}_4$	2.36	2.36 g	
$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	2.08	2.08 g	
NaCl	0.074	0.074 g	
$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	0.13	0.13 g	
$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$	0.0013	0.0013 g	Use 1000 X soln.
$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	0.0055	0.0055 g	Use 1000 X soln.
$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	0.0002	0.0002 g	Use 1000 X soln.
$\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$	0.0007	0.0007 g	Use 1000 X soln.
DI Water (L)		1 L	
Autoclave Time	15 min for small flasks 30 min for large flasks	30-60	
Comments:	Adjust to pH = 3.4 with H_2SO_4		

[0193] In one example, the kit or composition includes a modified Riscaldati medium with 20× trace elements, for example 20 times of the following

$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$	0.0013	0.0013 g	Use 1000 X soln.
$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	0.0055	0.0055 g	Use 1000 X soln.
$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	0.0002	0.0002 g	Use 1000 X soln.
$\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$	0.0007	0.0007 g	Use 1000 X soln.

EXAMPLE 1

Materials and Methods

[0194] This example describes methods used in the experiments described in Examples 2-6 below.

[0195] Strains and vector. The parental wild type *A. pseudoterreus* strain ATCC 32359 was from ATCC. The hygromycin phosphotransferase (hph) marker cassette was amplified from vector pCB1003.

[0196] Growth conditions. All strains were maintained on complete medium (CM) agar and conidia of spore were harvested from cultures grown for five days on complete medium (CM) plate (10 g Glucose, 2 g Triptase peptone, 1 g yeast extract, 1 g casamino acids, 50 ml 20×NO₃ Salts, 1 ml of 1000×Trace elements, 1 ml of 1000×Vitamin stock, in 1000 ml DI water, pH to 6.5), 20× NO₃ Salts contains in g/l,

Na₂NO₃, 120; KCL, 10.4 g; MgSO₄.7H₂O, 10.4 g; KH₂PO₄, 30.4 g. 1000× vitamin solution contains in per 100 ml H₂O: Biotin, 0.01 gm; pyridoxinHCL, 0.01 gm; thiamineHCL, 0.01 gm; riboflavin, 0.01 gm; paba, 0.01 gm; nicotinic acid, 0.01 gm, filtered and stock at 4° C. 1000× trace element contains in per 100 ml H₂O: ZnSO₄.7H₂O, 2.2 g; H₃BO₃, 1.1 g; MnCl₂.4H₂O, 0.5 g; FeSO₄.7H₂O, 0.5 g; CoCl₂.6H₂O, 0.17 g; CuSO₄.5H₂O, 0.16 g; Na₂MoO₄.2H₂O, 0.15 g; Na₂EDTA, 5 g, add the compounds in order, boil and cool to 60° C. Adjust pH to 6.5 with KOH. Cool to room temperature. Adjust volume to 100 ml with distilled water.

[0197] The transformants were selected for hygromycin resistance on minimum media (MM) (10 g Glucose, 50 ml 20×NO₃ Salts, 1 ml of 1000×Trace elements, 1 ml of 1000×Vitamin stock, 1000 ml DI water, pH to 6.25–6.5, hygromycin 100 ug/ml). 0.5×10⁸ conidia were inoculated into 50 ml of production media for itaconic acid production (Riscaldati medium) as described previously (100 g Glucose, 0.11 g KH₂PO₄, 2.36 g (NH₄)₂SO₄, 2.08 g MgSO₄.7H₂O, 0.074 g NaCl, 0.13 g CaCl₂.2H₂O, 1 ml of 1000× trace elements in 1000 ml DI water, adjust pH to 3.4 with H₂SO₄, 1000× trace elements contains 1.3 g/L ZnSO₄.7H₂O, 5.5 g/L FeSO₄.7H₂O, 0.2 g/L CuSO₄.5H₂O, 0.7 g/L MnCl₂.4H₂O). Cultivation was performed at 30° C. on a rotary shaker at 150rpm. At intervals during the incubation period, three single flasks were harvested for HPLC analysis, biomass measurement and RNA extraction. All experiments were replicated three times, and the standard deviation of the itaconic acid concentrations or dry weight was

always less than 10% of the mean. For collecting samples for EST analysis, *A. pseudoterreus* was grown in 20 liter stirred tank bioreactor.

[0198] Construction of deletion mutants. The deletion mutants were constructed by homologous double crossover with fusion PCR products. Synthetic oligos used for each construct are described in Table 1. Oligonucleotides were from IDT (Coraville, Iowa). Ex Taq polymerase (TaKaRa, Japan) was used to generate DNA constructs for making gene knockouts. Briefly, the 5' flanking region (~1.5 kb) of the target gene was amplified by primer pair F1 and R3. The 3' flanking region (~1.5 kb) of the target genes was amplified by primer pair F4 and R6. R3 and F4 carried 20-25 bases complementary to 5' and 3' ends of the hph cassette, respectively. The hph marker cassette was amplified from pCB1003 with the hphF and hphR primers that carried 30 bases complementary to the 3' end of the 5' flanking region and the 5' of the 3' flanking region, respectively. The three fragments, including the 5' flanking region, the hph marker cassette and the 3' flanking region were mixed in 1:3:1 molar ratio and combined by overlap PCR during the second round PCR. In the third round of PCR, the fusion PCR product was amplified with a nested primer pair (F2 and R5). This final PCR product carried a hygromycin marker cassette flanked by sequences homologous to the upstream and downstream regions of the target gene. 1-2 ug of the final product was used to transform strain *A. pseudoterreus* strain ATCC 32359.

TABLE 1

Primers for making deletion constructs		
gene targeted	primer name	primer sequence (SEQ ID NO)
tf	at tff1	gagccatagccatgcaagcg (1)
	at tff2	atagagtccttgatgagacg (2)
	at tfr5	gtggatttcagaggttccttgc (3)
	at tfr6	gaagtagaaccatgtggatcg (4)
	at hphf tfr3	tgacctccactagctccagcactactagatagggccgtttagagagtgc (5)
	at hphr tff4	aatagagttagatgcccagcgccgcttcgacgacagctctgcactctcc (6)
	at tfr3hphf	ggcactctctaaacgggctatctagtagtgcctggagctagtgagggtca (7)
	at tftff4hphr	ggagagtgcagagctgtcgtcgaagcgccggtcggcatctactctatt (8)
mttA	at motf1	gctgcatactcggattacgc (9)
	at motf2	Gaaaagggtactcggagtacg (10)
	at motr5	cagaccaaggagctttcctg (11)
	at motr6	cattaagccacaggcttgcg (12)
	athphfmotr3	tgacctccactagctccagcaatattggatgctgttcgttcgctgcgtgctgg (13)
	athphrmotf4	aatagagttagatgcccagcgtgacgaggatgtgctgagtcacaaacaaagc (14)
	at motr3hphf	ccagcacggcgaaacgaacagcatccatattgctggagctagtgagggtca (15)
	at motf4hphr	gctttgttggaactcagcacatcctcgtcacggtcggcatctactctatt (16)
cadA	at cadf1	ctccagtaacagaaccgacc (17)
	at cadf2	gaacttcactgcccattgg (18)
	at cadr5	ggacactccaaggagataagg (19)
	at cadr6	gctcatcacattgttgccg (20)
	at hphfcadr3	tgacctccactagctccagcggtaatttaaggaggacgatcttcgctgcg (21)
	at hphrcadf4	Aatagagttagatgcccagcgtcagcctggacaggctcaccgacattagcc (22)
	at cadr3hphf	cgcagcgaagatcgctcctctaaattgaccgctggagctagtgagggtca (23)
	at cadf4hphr	ggctaattgtcggtagcgtcgtccaggctgacgggtcggcatctactctatt (24)
mfsA	mfsf1	tgatgagctgaattcgttgc (25)
	mfsf2	tatagccagcttttgcgttg (26)
	mfsr5	catagcgttcagagtgttg (27)
	mfsr6	ccatttcaatgctttgtgcg (28)
	mfsr3hphf	ccataccacccttaccctcttgagtgctccgctggagctagtgagggtca (29)
	mfsf4hphr	gctgtggcctcctggcgattacgcaatattcggtcggcatctactctatt (30)
	hphfmfsr3	tgacctccactagctccagcgacactccaaggaggttaagggtggtatgg (31)
	hphrmfsf4	aatagagttagatgcccagcgaatattgctgaatcgccaggaggccacagc (32)

TABLE 1-continued

Primers for making deletion constructs		
gene targeted	primer name	primer sequence (SEQ ID NO)
p450	p450f1	tctccaaatcatcgatcatcg (33)
	p450f2	cttcaatcgaccgacatcc (34)
	p450r5	tcgtgtagacaagtcacgac (35)
	p450r6	ctataccactctagtgtatgg (36)
	p450r3hphf	cctctgctcaggtgttttcgaacaggagcgctgtagctagtggaggtca (37)
	p450f4hphr	cggaaatgcagataggcatcacagtcacagacggctggcatctactctatt (38)
	hphfp450r3	tgacctccactagctccagcgctcctgttctgaaacaacctgagcagagg (39)
	hphrp450f4	aatagagtagatgccgaccgttctggaactgtgatgcctatctgcattccg (40)

[0199] Transformation of *A. pseudoterreus* protoplasts. 10⁸ conidia of *A. pseudoterreus* ATCC 32359 were used to inoculate 300 ml Erlenmeyer baffles containing 100 ml of complete media. The cultures were grown overnight (16-18 hrs) at 30° C. and 150 rpm. The mycelia were then harvested by filtering the culture through miracloth and rinsing the mycelia mat with sterile water. The protoplasts were prepared by treating mycelia (mass of approximately 1-2 beans) with 20 mg/ml lysing enzyme (L1412, Sigma) dissolved in 20 ml of osmotic wash buffer (0.5M KCl, 10 mM sodium phosphate, pH 5.8) for 2 h. Protoplasts were collected by filtering protoplasts through sterile miracloth into a 50 ml screw cap centrifuge tube and centrifuging at 1000×g for 10 min at 4° C. Protoplasts were then washed twice with 20 ml Washing Solution (0.6M KCl, 0.1M Tris/HCl, pH 7.0) and once in 10 ml Conditioning Solution (0.6M KCl, 50 mM CaCl₂, 10 mM Tris/HCl, pH 7.5). For transformation, 1-2 µg DNA was added to 2×10⁷ protoplasts in 0.1 ml Conditioning Solution. A control reaction without added DNA was performed at the same time. 25 µl of PEG solution (25% PEG8000, 0.6M KCl, 50 mM CaCl₂, 10 mM Tris/HCl, and pH 7.5) was added and the protoplasts were incubated for 20 min on ice. An additional 500 µl of the PEG was added using a wide bore pipette tip and carefully mixed with the protoplasts by gently pipetting up and down 1-2 times. The protoplast solution was then incubated for 5 min on ice. 1 ml of cold Conditioning Solution was added and mixed by gently inverting the tube several times. Then the protoplast suspension was mixed with 12 ml of 50° C. selection agar (Minimal media+0.6M KCl+1.5% Agar+100 µg/ml hygromycin) contained in a 15 ml screwcap centrifuge tube. The tubes were then mixed by inverting the tubes 3-4 times and poured directly onto the petri dish plates. The control reaction was divided into a positive control plate (no selection antibiotics in the top agar and bottom plates) and a negative control (with selection hygromycin in top and bottom agar). The solidified plates were inverted and incubated overnight at 30° C. The next day, the plates were then overlaid with 15 ml of Minimal Medium (MM) containing 150 µg/ml hygromycin. Colonies should start to appear after 3-4 days. The transformants were excised and transferred to MM plate containing 100 µg/ml hygromycin. Correct transformants on the hygromycin plate were confirmed by PCR approaches and southern blot.

[0200] Dry mass measurement. Dry mass at each time point was determined by harvesting the mycelium on miracloth by suction filtration and washed twice with 50 ml distilled water. Subsequently, the dry weight was determined by drying it overnight in pre-weighed tubes on lyophilizer.

[0201] HPLC. The content of itaconic acid, aconitic acid, and glucose in each sample collected from filtration (0.22 µm) was assayed by a high-pressure liquid chromatography (HPLC) on a Bio-Rad Aminex HPX-87H ion exclusion column (300 mm×7.8 mm). Columns were eluted with Sulfuric acid (0.005 M) at a flow rate of 0.55 mL/min. The sample volume was 10-100 µl, and IA was detected at 210 nm with a Waters 2414 refractive index detector.

[0202] RNA isolation and transcript analysis by Quantitative real time RT-PCR. Wild type and Δ deletion strains were grown in Riscaldati medium at 30° C. After 3 days growth, mycelia were harvested, pressed dry between paper towels and immediately flash frozen in liquid nitrogen. The entire sample was then ground in a mortar and pestle with liquid nitrogen. Approximately 100 mg samples (about 0.1 ml) were extracted using Trizol® reagent (Chomczynski, *BioTechniques* 1993, 15(3):532-534, 536-537) and the resulting RNA was converted to cDNA using high capacity RNA-to-DNA kit (Applied Biosystems). Quantitative RT-PCR were performed in 50 µl reactions containing 25 µl of Power SYBR green PCR master mix (Applied Biosystems), 50 ng cDNA (from 50 ng RNA) and 0.2 uM forward and reverse primers. The RT-PCR primers used for analysis of the *mttA*, *cadA*, *mfsA* genes and *benA* (β -tubulin) as endogenous control gene are listed in Table 3. There are two additional controls, one is a no RT (without adding RT enzyme mix) control to estimate contamination from genomic DNA, and the other is no-template controls for each primer pair to measure effect from primer dimer formation. Amplification was performed using 7900HT Fast Real-Time PCR system (Applied Biosystems) programmed to initially hold at 95° C. for 10 min and then to complete 45 cycles of 95° C. for 15 s, 60° C., for 60 s. The data were analyzed using the comparative C_T method (e.g., see Schmittgen et al., *Analytical biochemistry* 2000, 285(2):194-204).

EXAMPLE 2

Expression Profile of Itaconic Acid Gene Cluster in *A. pseudoterreus*

[0203] RNA samples were prepared from three different growth stages of *A. pseudoterreus* in the itaconic acid production process. The stages were 1) “pre-production,” before itaconic acid production begins, 2) “production onset,” the beginning of itaconic acid production correlated with phosphate depletion, and 3) “production,” early in the phase of maximum itaconic acid production rate (FIG. 2).

EST data revealed four genes in the cluster having high expression frequency both in the onset phase and production phase, but not in the pre-production phase (Table 2). These genes were *tf*, *mttA*, *cadA*, and *mfsA*.

TABLE 2

Number of ESTs per gene at three stages of itaconic acid production				
Broad Institute Gene No.	Gene Description	Pre-Production	Production Onset	Production
ATEG_09968.1	upstream flanking gene; <i>lovE</i>	0	0	0
ATEG_09969.1	<i>tf</i>	0	4	4
ATEG_09970.1	<i>mttA</i>	0	81	93
ATEG_09971.1	<i>cadA</i>	0	77	110
ATEG_09972.1	<i>mfsA</i>	0	6	7
ATEG_09973.1	<i>p450</i>	0	7	11
ATEG_09974.1	downstream flanking gene	0	0	0
ATEG_09817.1	control; <i>gpdh</i>	31	51	43

[0204] *cadA* has 77 ESTs at the beginning of itaconic acid (IA) production and 110 ESTs during the IA production, while *mttA* has 81 and 93 ESTs respectively in each stage. Both have no transcript detected before IA is produced. Transcription factor (*tf*) and *mfsA*, like *cadA* and *mttA*, did not show any expression before IA production, but had significant levels of transcription following the initiation of itaconic acid production (Table 2).

[0205] When genes upstream and downstream of *tf*, *cadA*, *mttA* and *mfsA* were examined, a similar expression pattern was not observed. No transcript was detected for either upstream or downstream genes in any stage of IA production except for *p450*. Control gene *gpdh*, which is far away from this region, showed high expression through the whole growth stage. This EST data clearly demonstrated that four genes *tf*, *cadA*, *mttA* and *mfsA* have the same expression pattern and are closely related to the IA production process. In addition, these four genes are in the same cluster. They are turned on strongly at the onset of IA production and persists through the production phase (FIG. 2).

EXAMPLE 3

Effect of *tf*, *cadA*, *mttA* and *mfsA* Deletion on Itaconic Acid Production in *A. pseudoterreus*

[0206] A transformation system was developed to allow for transformation of *A. pseudoterreus* (see Example 1). This system was used to generate recombinant knockout strains for each of the endogenous *tf*, *cadA*, *mttA* and *mfsA* genes. The KO mutant strains were confirmed by PCR and southern blot. The transformation protocol gave very high frequency of homologous deletion, 8 out of 10 had the correct deletion. This high deletion frequency may be due to the presence of a *ku* gene mutation in the genome of wild-type *A. pseudoterreus*.

[0207] Biomass accumulation and itaconic acid (IA) production of each of the four knockout mutants and wild type *A. pseudoterreus* were measured at day 5. All strains, including wild type, had similar biomass accumulation (FIG. 3A). There is no significant difference in biomass among these five strains, indicating that deletion of these genes does not cause a noticeable growth defect.

[0208] However, the yield of IA was significantly lower in all four deletion strains (Δtf , $\Delta cadA$, $\Delta mttA$ and $\Delta mfsA$) when compared to wild type *A. pseudoterreus*. After 5 days growth in the Riscardi medium, the Δtf strain had only generated ~3g/l IA, compared to the wild type strain, which generated ~24 g/l of IA (about an 8-fold decrease). No detectable IA was produced by the $\Delta cadA$ and $\Delta mttA$ strains. $\Delta mfsA$ produced around 16 g/L itaconic acid, about 2/3 of wild type *A. pseudoterreus*.

[0209] These observations demonstrate that *tf*, *mttA*, *cadA* and *mfsA* genes play a role in itaconic acid production.

EXAMPLE 4

Production Kinetics of Itaconic Acid in Wild Type and *tf* Deletion Strain

[0210] To test the production kinetics in the deletion strains, Δtf and wild type *A. pseudoterreus* strain ATCC 32359 were tested for IA production during the growth on a rotary shaker for 7 days. IA was analyzed by HPLC for 2, 4, 6 and 7 day cultures.

[0211] As shown in FIG. 4, the IA yield plateaued at day 7 in both Δtf and wild type strains. Interestingly, the IA yield in Δtf (5g/l) is much lower than that of wild type (35 g/l), a decrease of about 7-fold. Thus, the Δtf strain produces IA at slower rate with a lower maximum IA yield than the wild type strain.

EXAMPLE 5

tf Regulation

[0212] The effects of *tf* gene deletion on the transcription level of other genes in the cluster were investigated by real-time reverse transcription PCR (RT-PCR). In the both Δtf and wild type strains, expression level of each gene was analyzed by RT-PCR by measuring *mttA*, *cadA*, *mfsA* mRNA levels using primers specific for those genes (Table 3).

TABLE 3

primers for real-time RT-PCR analysis of cluster gene transcript level		
Gene targeted	Primer name	Primer sequence (SEQ ID NO:)
<i>mttA</i>	<i>mttF</i>	Gctttcaatgtgggttctac (41)
	<i>mttR</i>	ctccatcacctacccttctc (42)
<i>cadA</i>	<i>cadF</i>	gaagtgtgggatctggc (43)
	<i>cadR</i>	gggttcggtatttgtgaag (44)
<i>mfsA</i>	<i>mfsF</i>	caagaacagtttggcctgag (45)
	<i>mfsR</i>	gcggacatcatacaatctgg (46)
<i>benA</i>	β -tubulinF	ttgtcgatgttctcgtcgc (47)
	β -tubulinR	tggcgttgtaaggctcaacc (48)

[0213] As shown in FIG. 5, in Δtf strains, mRNA level of *mttA* decreased 57 fold, *cadA* mRNA level decreased 37 fold, and *mfsA* decreased 23 fold, as compared to their expression in wild type *A. pseudoterreus* 32359. Thus, inactivation of the *tf* gene dramatically reduced the level of mRNA of other genes in the cluster. Within the itaconic acid biosynthesis cluster, the transcription factor potentially controls expression of other genes.

EXAMPLE 6

cadA Deletion Creates a Novel Strain that Produces Aconitic Acid

[0214] In *A. pseudoterreus*, when *cadA* was deleted, itaconic acid production was completely abolished (FIG. 3B). However, 3.5 g per liter aconitic acid in the Δ cadA strain was detected at day 5 (FIG. 6A). Aconitic acid was not produced by the wild type, Δ mttA or Δ mfsA strains (FIG. 6A). A time course analysis showed that aconitic acid started to appear in the supernatant at day 3, similar as IA in the wild type strain (FIG. 6B). At day 3, only cis-aconitic acid was detected in the supernatant. At day 4, both cis-aconitic acid and trans-aconitic acid were detected. From day 5 onward, cis-aconitic acid remained consistent at about 2 g/L, while trans-aconitic acid yield continued to increase (FIG. 6B). By day 10, 10 g/L trans-aconitic acid was detected in the supernatant from the Δ cadA strain (FIG. 6B). FIG. 6C shows a comparison of total aconitic acid production between wild type and Δ cadA fungi. Thus, Δ cadA strains of *A. pseudoterreus* and *A. terreus* can be used to produce cis- and trans-aconitic acid.

EXAMPLE 7

Materials and Methods

[0215] This example describes methods used in the experiments described in Example 8.

Transgene Expression Vector for 3-HP Production

[0216] Isolation of DNA Fragments:

[0217] Fragment 1: *A. pseudoterreus* 5'-cadA gene, 987 bp (SEQ ID NO: 59) isolated by PCR with the oligo pair 1969 and 1970 (SEQ ID NOS: 60 and 61, respectively) and *A. pseudoterreus* genomic DNA;

[0218] Fragment 2: *A. niger* gpdA promoter, 813 bp (SEQ ID NO: 62) isolated by PCR with oligo pair of 1971 and 1972 (SEQ ID NOS: 63 and 64, respectively) and *A. niger* genomic DNA;

[0219] Fragment 3: aspartate 1-decarboxylase (panD) cDNA of *Tribolium castaneum* with codon optimization for *A. pseudoterreus*, 1617 bp (SEQ ID NO: 65) was isolated by PCR with the oligo pair of 1973 and 1974 (SEQ ID NOS: 66 and 67, respectively) and the plasmid DNA containing the synthesized panD cDNA;

[0220] Fragment 4: bidirectional terminator from *A. niger* elf3/multifunctional chaperone (SEQ ID NO: 68) was iso-

lated by PCR with oligo pair of 1975 and 1976 (SEQ ID NOS: 69 and 70, respectively) and the genomic DNA of *A. niger*;

[0221] Fragment 5: codon optimized synthetic cDNA of β -alanine-pyruvate aminotransferase (BAPAT) of *Bacillus cereus*, 1350 bp (SEQ ID NO: 71) was isolated by PCR with oligo pair of 1977 and 1978 (SEQ ID NOS: 72 and 73, respectively) and the plasmid DNA containing the synthesized BABAT cDNA;

[0222] Fragment 6: *A. niger* eno1 promoter, 704 bp (SEQ ID NO: 74) isolated by PCR with oligo pair of 1979 and 1980 (SEQ ID NOS: 75 and 76, respectively) and *A. niger* genomic DNA;

[0223] Fragment 7: *A. nidulans* gpdA promoter, 885 bp (SEQ ID NO: 77) was isolated by PCR with the oligo pair of 2002 and 1982 (SEQ ID NOS: 78 and 79, respectively) and *A. nidulans* genomic DNA;

[0224] Fragment 8: the codon optimized synthetic cDNA of *E. coli* 3-hydroxypropionate dehydrogenase (HPDH), 741 bp (SEQ ID NO: 80) was isolated by PCR with oligo pair of 1983 and 1984 (SEQ ID NOS: 81 and 82, respectively) and the plasmid DNA containing the codon-optimized synthesized HPDH DNA of *E. coli*;

[0225] Fragment 9: trpC terminator of *A. nidulans*, 473 bp (SEQ ID NO: 83) isolated by PCR with oligo pair of 1985 and 2004 (SEQ ID NOS: 84 and 85, respectively) and plasmid DNA of pAN7.1;

[0226] Fragment 10: trpC terminator of *A. nidulans*, 473 bp (SEQ ID NO: 86) isolated by PCR with the oligo pair of 2005 and 1986 (SEQ ID NOS: 87 and 88, respectively) and plasmid DNA of pAN7.1;

[0227] Fragment 11: *A. oryzae* ptrA selection marker gene, 2005 bp; SEQ ID NO: 89) isolated by PCR with the oligo pair of 1987 and 1988 (SEQ ID NOS: 90 and 91, respectively) and *A. oryzae* genomic DNA;

[0228] Fragment 12: *A. pseudoterreus* 3'-cadA gene, 908 bp (SEQ ID NO: 92) isolated by PCR with the oligo pair 1989 and 2003 (SEQ ID NOS: 93 and 94, respectively) and *A. oryzae* genomic DNA;

[0229] Fragment 13 (SEQ ID NO: 95): Combination of Fragments 7 to 9 (SEQ ID NOS: 77, 80, and 83, respectively), 2099 bp isolated by PCR with oligo pair of 1981 and 1986 (SEQ ID NOS: 96 and 88, respectively) and plasmid DNA of pZD-2; and

[0230] Fragment 14 (SEQ ID NO: 97): Combination of Fragments 11 to 12 (SEQ ID NOS: 89 and 92, respectively), 2913 bp was isolated by PCR with the oligo pair of 1987 and 1990 (SEQ ID NOS: 90 and 98, respectively) and plasmid DNA of pZD-3.

[0231] The oligonucleotide primers used are shown in Table 4.

TABLE 4

Primers used to generate vector for 3-HP production	
Name	Sequence (SEQ ID NO:)
1969cad1	ccctcgagggtcgacggtatcgata GATATC GGTTGTAGCAGCGTAAACAC (60)
1970cad2	tcttttcatagttagCCTTGGTGAACATCTTGAGG (61)
1971gpdA1	atgttcaccaaggCTACTATGAAAGACCGCATG (63)
1972gpdA2	cgccggtggcgggCATTGTTTAGATGTGTCTATGTG (64)
1973pan1	catctaacaatgCCCGCCACCGGCGAGGACCA (66)

TABLE 4-continued

Primers used to generate vector for 3-HP production	
Name	Sequence (SEQ ID NO:)
1974pan2	atccaacccatcaGAGGTCGGAGCCCAGGCGTTCG (67)
1975ter1	gggctccgacctcTGATGGGTGGATGACGATG (69)
1976ter2	tctggcccagctcTGAGTCCTAGATGGGTGGTG (70)
1977bap1	catctaggactcaGAGCTGGGCCAGACATTCCTTC (72)
1978bap2	gtccatcaacatgAACTGATGATCGTCCAGGTCAC (73)
1979eno1	cgatcatcagttcCATGTTGATGGACTGGAGGG (75)
1980eno2	gaactagtggatccccgggctgcGttaaCTCGAGCTTACAAGAAGTA GCC (76)
1981gpdA1	acaggctacttcttgaagctcgagttTCTGTACAGTGACCGGTGAC (96)
1982gpdA2	tgaccagcacgatCATGGTGATGTCTGCTCAAG (79)
1983hpd1	agacatcaccatgATCGTGCTGGTCACGGGCGC (81)
1984hpd2	gccatcggtcctaTTGGCGGTGGACGTTACGGC (82)
1985trp1	cgtccaccgccaatAGGACCGATGGCTGTGTAG (84)
1986trp2	cccgtctgtcagaGAGCGGATTCTCAGTCTCG (88)
1987ptrA1	gaggaaatccgctcTCTGACAGACGGGCAATTGATTAC (90)
1988ptrA2	gaatgttgctgagGAGCCGCTCTTGCATCTTTG (91)
1989cad3	gcaagagcggctcCTCAGCAACATTCGCCATGTTC (93)
1990cad4	actaaagggaacaaaagctggagctCAGCTCCACTGCTCATAGTCTTT G (98)
2002gpdA5	ccctcgaggtcgacgggtatcgataGTTAACTCTGTACAGTGACCGGTG AC (78)
2003cad3	gaactagtggatccccgggctgcaCAGCTCCACTGCTCATAGTCTTT G (94)
2004trpR	gaactagtggatccccgggctgcaGAGCGGATTCTCAGTCTCG (85)
2005trpF	ccctcgaggtcgacgggtatcgataTAGGACCGATGGCTGTGTAG (87)

[0232] An overview of the arrangement of the Fragments is shown in FIG. 7. Fragments 1 to 6 (SEQ ID NOS: 59, 62, 65, 68, 71 and 74, respectively) were assembled into the plasmid DNA pBlueScript SK (–) linearized with HindIII and PstI via Gibson Assembly master kit to form plasmid pZD-1. A restriction enzyme site HpaI was introduced at the end of the fragment 6 for further cloning.

[0233] Fragments 7 to 9 (SEQ ID NOS: 77, 80, and 83, respectively) were assembled into the plasmid DNA pBlueScript SK (–) linearized with HindIII and PstI via Gibson Assembly master kit to form plasmid DNA pZD-2.

[0234] Fragments 10 to 12 (SEQ ID NOS: 86, 89, and 92, respectively) were assembled into the pBlueScript SK (–) vector linearized with restriction enzyme HindIII and PstI by Gibson assembly to form the plasmid vector ZD-3. (Only fragments 11 and 12 were used in the next step; SEQ ID NOS: 89 and 92).

[0235] Fragments 13 and 14 (SEQ ID NOS: 95 and 97) were assembled together into the plasmid DNA vector ZD-1

linearized with restriction enzyme HpaI/SacI via Gibson Assembly master kit to form pZD-4.

[0236] Genomic DNA isolation and Southern blotting analysis were performed as described in Example 1 (and see Dai et al., 2017, *Appl Microbiol Biotechnol* 101:6099-6110).

Detection of 3-HP

[0237] The extracellular 3-HP in the culture supernatants was quantified with HPLC method as described in Example 1.

EXAMPLE 8

Production of 3-HP

[0238] The constructs generated in Example 7 (FIG. 7) were transformed into wild type *A. pseudoterreus* strain ATCC 32359 using the methods describe in Example 1, thereby inactivating/disrupting the cadA gene in some examples.

[0239] As shown in FIG. 8, restriction fragment length polymorphism of selected transgenic strains show that the transgene expression cassette was inserted into the *cadA* locus in strain-2 (with one copy) and strain-6 (two copies), while the strain-4 and strain-5 carry the transgene expression cassette with random integration. No integration of transgene expression cassette was observed in strain-1 and strain-3.

[0240] 3-HP production was measured in several transformants. As shown in FIG. 9A, the Δ *cadA* strain did not produce 3-HP, while insertion of the transgene expression cassette that allowed for expression of *panD*, *BAPAT*, and *HPDH*, into the *cadA* locus with one copy or two copies and resulted in 0.9 or 1.7 g/l 3-HP accumulation in the strains

3HP-2 or 3HP-6. In contrast, when the transgene expression cassette was randomly inserted into the chromosome, 3HP production was substantially lower (Strains 3HP-4 and 3HP-5). FIG. 9B shows 3-HP production over 8 days in Strains 3HP-2 and 3HP-6 (strains 2 and 6, respectively). Thus, genetically inactivating *cadA* can increase 3-HP production.

[0241] In view of the many possible embodiments to which the principles of the disclosure may be applied, it should be recognized that the illustrated embodiments are only examples of the invention and should not be taken as limiting the scope of the invention. Rather, the scope of the invention is defined by the following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 98

<210> SEQ ID NO 1

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete *tf* gene in *A. pseudoterreus*

<400> SEQUENCE: 1

gagccatagc catgcaagcg

20

<210> SEQ ID NO 2

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete *tf* gene in *A. pseudoterreus*

<400> SEQUENCE: 2

atagagtcct tggatgagac g

21

<210> SEQ ID NO 3

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete *tf* gene in *A. pseudoterreus*

<400> SEQUENCE: 3

gtggatttcg aggttccttg c

21

<210> SEQ ID NO 4

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete *tf* gene in *A. pseudoterreus*

<400> SEQUENCE: 4

gaagtagaac catgtggatc g

21

<210> SEQ ID NO 5

<211> LENGTH: 50

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete tf gene in A.
pseudoterreus

<400> SEQUENCE: 5

tgacctccac tagctccagc actactagat aggcccgttt agagagtgcc 50

<210> SEQ ID NO 6
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete tf gene in A.
pseudoterreus

<400> SEQUENCE: 6

aatagagtag atgccgaccg gccgcttcga cgacagctct gcaactctcc 49

<210> SEQ ID NO 7
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete tf gene in A.
pseudoterreus

<400> SEQUENCE: 7

ggcactctct aaacgggcct atctagtagt gctggagcta gtggagggtca 50

<210> SEQ ID NO 8
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete tf gene in A.
pseudoterreus

<400> SEQUENCE: 8

ggagagtgcga gagctgtcgt cgaagcggcc ggtcggcatc tactctatt 49

<210> SEQ ID NO 9
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mttA gene in A.
pseudoterreus

<400> SEQUENCE: 9

gctgcatact cggattacgc 20

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mttA gene in A.
pseudoterreus

<400> SEQUENCE: 10

gaaaaggtac tcggagtacg 20

<210> SEQ ID NO 11

-continued

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mttA gene in A.
pseudoterreus

<400> SEQUENCE: 11

cagaccaagg agctttcctg 20

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mttA gene in A.
pseudoterreus

<400> SEQUENCE: 12

cattaagcca caggcttgcg 20

<210> SEQ ID NO 13
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mttA gene in A.
pseudoterreus

<400> SEQUENCE: 13

tgacctccac tagctccagc aatatggatg ctgttcgttc gccgtgctgg 50

<210> SEQ ID NO 14
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mttA gene in A.
pseudoterreus

<400> SEQUENCE: 14

aatagagtag atgccgaccg tgacgaggat gtgctgagtc caaacaagc 50

<210> SEQ ID NO 15
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mttA gene in A.
pseudoterreus

<400> SEQUENCE: 15

ccagcacggc gaacgaacag catccatatt gctggagcta gtggaggcca 50

<210> SEQ ID NO 16
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mttA gene in A.
pseudoterreus

<400> SEQUENCE: 16

gctttgtttg gactcagcac atcctcgtca cggtcggcat ctactctatt 50

-continued

<210> SEQ ID NO 17
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete cadA gene in A.
pseudoterreus

<400> SEQUENCE: 17

ctccagtaac agaaccgacc 20

<210> SEQ ID NO 18
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete cadA gene in A.
pseudoterreus

<400> SEQUENCE: 18

gaacttcact gccgcattgg 20

<210> SEQ ID NO 19
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete cadA gene in A.
pseudoterreus

<400> SEQUENCE: 19

ggacactcca agaggataag g 21

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete cadA gene in A.
pseudoterreus

<400> SEQUENCE: 20

gctcatcaca ttgtttgccg 20

<210> SEQ ID NO 21
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete cadA gene in A.
pseudoterreus

<400> SEQUENCE: 21

tgacctccac tagctccagc ggtcaattta agaggacgat cttcgtgcg 50

<210> SEQ ID NO 22
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete cadA gene in A.
pseudoterreus

<400> SEQUENCE: 22

-continued

aatagagtag atgccgaccg tcagcctgga caggctcacc gacattagcc 50

<210> SEQ ID NO 23
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete cadA gene in A.
pseudoterreus

<400> SEQUENCE: 23

cgcagcgaag atcgctctct taaattgacc gctggagcta gtggagggtca 50

<210> SEQ ID NO 24
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete cadA gene in A.
pseudoterreus

<400> SEQUENCE: 24

ggctaagtgc ggtgagcctg tccaggctga cggtcggcat ctactctatt 50

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 25

tgatgagctg aattcgttgc 20

<210> SEQ ID NO 26
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 26

tatagccagc ttttgctgtg 20

<210> SEQ ID NO 27
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 27

catagcggtc agagtgttg 19

<210> SEQ ID NO 28
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

-continued

<400> SEQUENCE: 28

ccatttcaat gctttgtgcg 20

<210> SEQ ID NO 29

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 29

ccataccacc cttaccctct tggagtgtcc gctggagcta gtggaggtca 50

<210> SEQ ID NO 30

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 30

gctgtggcct cctggcgatt acgcaatatt cggtcggcat ctactctatt 50

<210> SEQ ID NO 31

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 31

tgacctccac tagctccagc ggacactcca agagggttaag ggtggtatgg 50

<210> SEQ ID NO 32

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 32

aatagagtag atgccgaccg aatattgcgt aatcgccagg aggccacagc 50

<210> SEQ ID NO 33

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete p450 gene in A.
pseudoterreus

<400> SEQUENCE: 33

tctccaaatc atcgtcacg 20

<210> SEQ ID NO 34

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to delete p450 gene in A.

-continued

pseudoterreus

<400> SEQUENCE: 34

cttcaatcgc accgacatcc 20

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete p450 gene in A.
pseudoterreus

<400> SEQUENCE: 35

tcgtgtagac aagtccagtc 20

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete p450 gene in A.
pseudoterreus

<400> SEQUENCE: 36

ctataccact ctagtgatgg 20

<210> SEQ ID NO 37
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete p450 gene in A.
pseudoterreus

<400> SEQUENCE: 37

cctctgctca gggtgttttc gaacaggagc gctggagcta gtggaggcca 50

<210> SEQ ID NO 38
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete p450 gene in A.
pseudoterreus

<400> SEQUENCE: 38

cggaatgcag ataggcatca cagtccagaa cggtcggcat ctactctatt 50

<210> SEQ ID NO 39
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete p450 gene in A.
pseudoterreus

<400> SEQUENCE: 39

tgacctccac tagctccagc gctcctgttc gaaaacaacc tgagcagagg 50

<210> SEQ ID NO 40
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete p450 gene in A.
pseudoterreus

<400> SEQUENCE: 40

aatagagtag atgccgaccg ttctggactg tgatgcctat ctgcattccg 50

<210> SEQ ID NO 41
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to amplify mttA in A.
pseudoterreus

<400> SEQUENCE: 41

gctttcaatg tggttcctac 20

<210> SEQ ID NO 42
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to amplify mttA in A.
pseudoterreus

<400> SEQUENCE: 42

ctccatcacc tacccttcc 19

<210> SEQ ID NO 43
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to amplify cadA in A.
pseudoterreus

<400> SEQUENCE: 43

gaagtgtggg atctggc 17

<210> SEQ ID NO 44
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to amplify cadA in A.
pseudoterreus

<400> SEQUENCE: 44

gggttcggta tttgtgaag 19

<210> SEQ ID NO 45
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to amplify mfsA in A.
pseudoterreus

<400> SEQUENCE: 45

caagaacagt ttggcctgag 20

<210> SEQ ID NO 46
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to amplify mfsA in A.
pseudoterreus

<400> SEQUENCE: 46

gcggacatca tacaatctgg 20

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to amplify beta-tubulin in A.
pseudoterreus

<400> SEQUENCE: 47

ttgtcgatgt tgttcgtcgc 20

<210> SEQ ID NO 48
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to amplify beta-tubulin in A.
pseudoterreus

<400> SEQUENCE: 48

tggcgttgta aggetcaacc 20

<210> SEQ ID NO 49
<211> LENGTH: 2206
<212> TYPE: DNA
<213> ORGANISM: Aspergillus terreus

<400> SEQUENCE: 49

gtgggtcttg aaatcgtatg ccacacttgc tccggatgaa acacattccg gagcgcgcat 60

cgatattgct acacagtata gacccaatgg tctgcagatg ccctaaatgg tagttctcac 120

tggcctgcat taagtctctg ttgcagatca ttgtcggcct aacatcagtg taggttacgg 180

tgtgagattt acttgcatag aagattccag accacaaggt tctagatcct ttgacggcgg 240

actcccctcg aggtgccggg cgccgacgtg tgcgttgctc cgggatttgt aggacgcagc 300

tccgatacct agccgttatg ggaatcggag gttgtagcag cgtaaacaca tggatagtta 360

aataatcggg tgtacacca ctgttggaag tgacgggggc ctacaacacg agattatctg 420

atccaatttc tgttcgttgg cattctatca ttcgcagcga aaattgtcct attaaattga 480

ccatgaccaa acaatctgcg gacagcaacg caaagtcagg agttacgtcc gaaatatgtc 540

attgggcac ccaactggcc actgacgaca tcccttcgga cgtattagaa agagcaaaat 600

accttattct cgacgggtatt gcatgtgcct gggttggtgc aagagtgcct tggtcagaga 660

agtatgttca ggcaacgatg agctttgagc cgccgggggc ctgcagggtg attggatatg 720

gacaggtaaa ttttatctac tctagacggg ccacaaagta tactgacgat ccttcgtata 780

gaaactgggg cctgttgagc cagccatgac caattccgct ttcatacagg ctacggagct 840

tgacgactac cacagcgaag cccccctaca ctctgcaagc attgtccttc ctgcggtcct 900

tgcagcaagt gaggtcttag ccgagcaggg caaaacaatt tccggtatag atgttattct 960

-continued

```

agccgccatt gtggggtttg aatctggccc acggatcggc aaagcaatct acggatcggc 1020
cctcttgaac aacggctggc attgtggagc tgtgtatggc gctccagccg gtgcgctggc 1080
cacaggaaag ctctcggtc taactccaga ctccatggaa gatgctctcg gaattgcgtg 1140
cacgcaagcc tgtggtttaa tgcggcgca atacggaggc atggtaaagc gtgtgcaaca 1200
cggattcgca gcgcgtaatg gtcttcttgg gggactgttg gcccatggtg ggtacgaggc 1260
aatgaaaggt gtcttgaga gatcttacgg cggtttcctc aagatgttca ccaagggcaa 1320
cggcagagag cctccctaca aagaggagga agtggtggct ggtctcggtt cattctggca 1380
tacctttact attcgcatca agctctatgc ctgctcgga cttgtccatg gtccagtcga 1440
ggctatcgaa aaccttcagg ggagataccc cgagctcttg aatagagcca acctcagcaa 1500
cattcgccat gttcatgtac agctttcaac ggcctcgaa agtcactgtg gatggatacc 1560
agaggagaga cccatcagtt caatcgcagg gcagatgagt gtcgcataca ttctcgccgt 1620
ccagctggtc gaccagcaat gtcttttgc ccagttttct gagtttgatg acaacctgga 1680
gaggccagaa gtttgggata tggccaggaa ggttacttca tctcaaagcg aagagtttga 1740
tcaagacggc aactgtctca gtgcgggtcg cgtgaggatt gagttcaacg atggttcttc 1800
tattacggaa agtgctgaga agcctcttgg tgtcaaagag cccatgccaa acgaacggat 1860
tctccacaaa taccgaaccc ttgctggtag cgtgacggac gaatcccggtg tgaagagat 1920
tgaggatctt gtctcggcc tggacaggct caccgacatt agccattgc tggagctgct 1980
gaattgcccc gtgaaatgc cactggtata aatgggaagc gatatggaaa catttcatgt 2040
cacgggcaca aattctaggt catatcgta ctggatggtg aaaccaccag cggttttagca 2100
gatagaagat agactccttc tgctctgctg tgcgtcttga attagttcg ttcactggct 2160
taagaactta gaatgcaata cagtctctct tatttcttat taaaat 2206

```

<210> SEQ ID NO 50

<211> LENGTH: 490

<212> TYPE: PRT

<213> ORGANISM: *Aspergillus terreus*

<400> SEQUENCE: 50

```

Met Thr Lys Gln Ser Ala Asp Ser Asn Ala Lys Ser Gly Val Thr Ser
1          5          10          15

Glu Ile Cys His Trp Ala Ser Asn Leu Ala Thr Asp Asp Ile Pro Ser
20        25        30

Asp Val Leu Glu Arg Ala Lys Tyr Leu Ile Leu Asp Gly Ile Ala Cys
35        40        45

Ala Trp Val Gly Ala Arg Val Pro Trp Ser Glu Lys Tyr Val Gln Ala
50        55        60

Thr Met Ser Phe Glu Pro Pro Gly Ala Cys Arg Val Ile Gly Tyr Gly
65        70        75        80

Gln Lys Leu Gly Pro Val Ala Ala Ala Met Thr Asn Ser Ala Phe Ile
85        90        95

Gln Ala Thr Glu Leu Asp Asp Tyr His Ser Glu Ala Pro Leu His Ser
100       105       110

Ala Ser Ile Val Leu Pro Ala Val Phe Ala Ala Ser Glu Val Leu Ala
115       120       125

Glu Gln Gly Lys Thr Ile Ser Gly Ile Asp Val Ile Leu Ala Ala Ile
130       135       140

```

-continued

Val	Gly	Phe	Glu	Ser	Gly	Pro	Arg	Ile	Gly	Lys	Ala	Ile	Tyr	Gly	Ser	145	150	155	160
Asp	Leu	Leu	Asn	Asn	Gly	Trp	His	Cys	Gly	Ala	Val	Tyr	Gly	Ala	Pro	165	170	175	
Ala	Gly	Ala	Leu	Ala	Thr	Gly	Lys	Leu	Leu	Gly	Leu	Thr	Pro	Asp	Ser	180	185	190	
Met	Glu	Asp	Ala	Leu	Gly	Ile	Ala	Cys	Thr	Gln	Ala	Cys	Gly	Leu	Met	195	200	205	
Ser	Ala	Gln	Tyr	Gly	Gly	Met	Val	Lys	Arg	Val	Gln	His	Gly	Phe	Ala	210	215	220	
Ala	Arg	Asn	Gly	Leu	Leu	Gly	Gly	Leu	Leu	Ala	His	Gly	Gly	Tyr	Glu	225	230	235	240
Ala	Met	Lys	Gly	Val	Leu	Glu	Arg	Ser	Tyr	Gly	Gly	Phe	Leu	Lys	Met	245	250	255	
Phe	Thr	Lys	Gly	Asn	Gly	Arg	Glu	Pro	Pro	Tyr	Lys	Glu	Glu	Glu	Val	260	265	270	
Val	Ala	Gly	Leu	Gly	Ser	Phe	Trp	His	Thr	Phe	Thr	Ile	Arg	Ile	Lys	275	280	285	
Leu	Tyr	Ala	Cys	Cys	Gly	Leu	Val	His	Gly	Pro	Val	Glu	Ala	Ile	Glu	290	295	300	
Asn	Leu	Gln	Gly	Arg	Tyr	Pro	Glu	Leu	Leu	Asn	Arg	Ala	Asn	Leu	Ser	305	310	315	320
Asn	Ile	Arg	His	Val	His	Val	Gln	Leu	Ser	Thr	Ala	Ser	Asn	Ser	His	325	330	335	
Cys	Gly	Trp	Ile	Pro	Glu	Glu	Arg	Pro	Ile	Ser	Ser	Ile	Ala	Gly	Gln	340	345	350	
Met	Ser	Val	Ala	Tyr	Ile	Leu	Ala	Val	Gln	Leu	Val	Asp	Gln	Gln	Cys	355	360	365	
Leu	Leu	Ser	Gln	Phe	Ser	Glu	Phe	Asp	Asp	Asn	Leu	Glu	Arg	Pro	Glu	370	375	380	
Val	Trp	Asp	Leu	Ala	Arg	Lys	Val	Thr	Ser	Ser	Gln	Ser	Glu	Glu	Phe	385	390	395	400
Asp	Gln	Asp	Gly	Asn	Cys	Leu	Ser	Ala	Gly	Arg	Val	Arg	Ile	Glu	Phe	405	410	415	
Asn	Asp	Gly	Ser	Ser	Ile	Thr	Glu	Ser	Val	Glu	Lys	Pro	Leu	Gly	Val	420	425	430	
Lys	Glu	Pro	Met	Pro	Asn	Glu	Arg	Ile	Leu	His	Lys	Tyr	Arg	Thr	Leu	435	440	445	
Ala	Gly	Ser	Val	Thr	Asp	Glu	Ser	Arg	Val	Lys	Glu	Ile	Glu	Asp	Leu	450	455	460	
Val	Leu	Gly	Leu	Asp	Arg	Leu	Thr	Asp	Ile	Ser	Pro	Leu	Leu	Glu	Leu	465	470	475	480
Leu	Asn	Cys	Pro	Val	Lys	Ser	Pro	Leu	Val							485	490		

<210> SEQ ID NO 51

<211> LENGTH: 1862

<212> TYPE: DNA

<213> ORGANISM: Aspergillus vadensis

<400> SEQUENCE: 51

gttttctgtg tgtctttggg gggttataaa tagggtgtcg aatatctgga agatagggaa

60

-continued

```

ttctttctct ttcaatcaat caatcaagaa ttcttttagg gagtttctat actacatccg 120
atatggctgc catcacgcgt aaatctgaag cggcttctgc tacttcgccc attcctacca 180
attctaatac taccatgact actaccctca acggggtaga tggttcaaaa gagaaagaaa 240
aagaccagat acccccaaac aaagaggaag gaacaaaagc agaagagaaa gaaaccgaag 300
catacaactc ctccaacggc gtcaccagcc aactctgcaa ctggatcgcc tctctccagc 360
tagaagacat tccagactct gtccgcaccc gcgccaagta cctctttctc gatggcatcg 420
cctgcgcact cgtcgggtgc cgcgtcccat ggtcgcagaa ggcttttgat gcgatggctg 480
ttttcgagga gaagggaag catgtggta ttgggtatga agagcgctt ggtgctatcg 540
ccgccgaac cctcaacggc tcttgatcc aagcctgcga agtagacgac taccacagcg 600
tggcgccct gcactcgag gccgtggta tccctctct cttcgtgccc gccgtcagtg 660
cgcggaacca tccgaccgca ccgcgcacat tcgacgggcg aacacttctt ctgcctccg 720
tggtagggtt cgaggttggt ccgcgcgtgg gcatggcgt acacggcacc gagatgctcg 780
cgaagggatg gcactcgagg tctgtgttg gtggaccgc ggccgcaggc agttctgcaa 840
aactactcgg tttgtcggc ggtcaagtcg aagacgcgat cggagtagca gcgacacaag 900
catgcgact catggcggc cagtacgac ggatggtgaa gcgatgcat catggcttcg 960
cggcaaggaa tggactgttg ggcacgatgt tagcgtggg aggttatgaa gggatcaaga 1020
aggtgtttga gcgccgtat ggaggatttc tggcaatgtt tggcctaggg tcgaagcaca 1080
cgcctagttc gaagccggag gaggtggcaa aggatttggg gacgttcttg cacacggcgg 1140
agtggattcg gttgaagttg catgcgtgct gtggggggat tcatggcacg attgagtgtt 1200
tggcggagat gcaggagatg tatccagagc gatttggacg ggagaaacta ggagagatca 1260
aggagattcg gatccagttg agtgatgcgg tgtttcatca ttgtggatgg gcgccggaga 1320
cgaggccgtt gaccccgacg gggggcgaga tgaatacggc gtttgtggcg gcctcgcagt 1380
tgggtgatgg acaagtgttg ttggagcagt tctcgtcggg gaagtggat cgggatgagg 1440
tttgggaatt gattgggaag acgagttgta ttcatacggc ggagttggac aagccgaata 1500
ttggtgttg tgcgttgatc tccatcacgt ttgcggatgg cagtcagggt cagcattcgt 1560
tgttgaagcc gaagggggtg gatgaacca ttccgaatga ggagatcttg gagaagtctc 1620
gtcgttgac gggcgggttg attggggtg agaggcagga gaagattgaa aaggcgtgc 1680
tggggatgga ggagttgcag gatgtggatg agttgattga gttgctgagt gtgaatgtgg 1740
tcaatccgtt gcagtagtat actagtcac tgttttgatg cttctggcgt tggctcgtgtt 1800
gggatagtat ctcataatct tgaattaata aatcattcaa catggtgaaa atcatatttg 1860
tg

```

<210> SEQ ID NO 52

<211> LENGTH: 544

<212> TYPE: PRT

<213> ORGANISM: *Aspergillus vadensis*

<400> SEQUENCE: 52

```

Met Val Ala Ile Thr Ala Lys Ser Glu Ala Ala Ser Ala Thr Ser Pro
1           5           10           15

```

```

Ile Pro Thr Asn Ser Asn Thr Thr Met Thr Thr Thr Leu Asn Gly Val
          20           25           30

```

-continued

Asp	Gly	Ser	Lys	Glu	Lys	Glu	Lys	Asp	Gln	Ile	Pro	Pro	Asn	Lys	Glu
35															
Glu	Gly	Thr	Lys	Ala	Glu	Glu	Lys	Glu	Thr	Glu	Ala	Tyr	Asn	Ser	Ser
50															
Asn	Gly	Val	Thr	Ser	Gln	Leu	Cys	Asn	Trp	Ile	Ala	Ser	Leu	Gln	Leu
65															
Glu	Asp	Ile	Pro	Asp	Ser	Val	Arg	Thr	Arg	Ala	Lys	Tyr	Leu	Phe	Leu
85															
Asp	Gly	Ile	Ala	Cys	Ala	Leu	Val	Gly	Ala	Arg	Val	Pro	Trp	Ser	Gln
100															
Lys	Ala	Phe	Asp	Ala	Met	Ala	Val	Phe	Glu	Glu	Lys	Gly	Lys	His	Val
115															
Val	Ile	Gly	Tyr	Glu	Glu	Arg	Leu	Gly	Ala	Ile	Ala	Ala	Ala	Thr	Leu
130															
Asn	Gly	Ser	Trp	Ile	Gln	Ala	Cys	Glu	Val	Asp	Asp	Tyr	His	Ser	Val
145															
Ala	Pro	Leu	His	Ser	Gln	Ala	Val	Val	Ile	Pro	Pro	Leu	Phe	Ala	Ala
165															
Ala	Val	Ser	Ala	Arg	Asn	His	Pro	Thr	Ala	Pro	Arg	Ile	Ile	Asp	Gly
180															
Arg	Thr	Leu	Leu	Leu	Ala	Ser	Val	Val	Gly	Phe	Glu	Val	Gly	Pro	Arg
195															
Val	Gly	Met	Ala	Leu	His	Gly	Thr	Glu	Met	Leu	Ala	Lys	Gly	Trp	His
210															
Cys	Gly	Ser	Val	Phe	Gly	Gly	Pro	Ala	Ala	Ala	Gly	Ser	Ser	Ala	Lys
225															
Leu	Leu	Gly	Leu	Ser	Ala	Gly	Gln	Val	Glu	Asp	Ala	Ile	Gly	Val	Ala
245															
Ala	Thr	Gln	Ala	Cys	Gly	Leu	Met	Ala	Ala	Gln	Tyr	Asp	Gly	Met	Val
260															
Lys	Arg	Met	His	His	Gly	Phe	Ala	Ala	Arg	Asn	Gly	Leu	Leu	Gly	Thr
275															
Met	Leu	Ala	Trp	Gly	Gly	Tyr	Glu	Gly	Ile	Lys	Lys	Val	Phe	Glu	Arg
290															
Pro	Tyr	Gly	Gly	Phe	Leu	Ala	Met	Phe	Gly	Leu	Gly	Ser	Lys	His	Thr
305															
Pro	Ser	Ser	Lys	Pro	Glu	Glu	Val	Ala	Lys	Asp	Leu	Gly	Thr	Phe	Trp
325															
His	Thr	Ala	Glu	Trp	Ile	Arg	Leu	Lys	Leu	His	Ala	Cys	Cys	Gly	Gly
340															
Ile	His	Gly	Thr	Ile	Glu	Cys	Leu	Ala	Glu	Met	Gln	Glu	Met	Tyr	Pro
355															
Glu	Arg	Phe	Gly	Arg	Glu	Lys	Leu	Gly	Glu	Ile	Lys	Glu	Ile	Arg	Ile
370															
Gln	Leu	Ser	Asp	Ala	Val	Phe	His	His	Cys	Gly	Trp	Ala	Pro	Glu	Thr
385															
Arg	Pro	Leu	Thr	Pro	Thr	Gly	Ala	Gln	Met	Asn	Thr	Ala	Phe	Val	Ala
405															
Ala	Ser	Gln	Leu	Val	Asp	Gly	Gln	Val	Leu	Leu	Glu	Gln	Phe	Ser	Ser
420															
425															
430															

-continued

Gly Lys Leu Asp Arg Asp Glu Val Trp Glu Leu Ile Gly Lys Thr Ser
435 440 445

Cys Ile His Thr Ala Glu Leu Asp Lys Pro Asn Ile Gly Cys Gly Ala
450 455 460

Leu Ile Ser Ile Thr Phe Ala Asp Gly Ser Gln Val Gln His Ser Leu
465 470 475 480

Leu Lys Pro Lys Gly Val Asp Glu Pro Ile Ser Asn Glu Glu Ile Leu
485 490 495

Glu Lys Phe Arg Arg Leu Thr Gly Gly Leu Ile Gly Val Glu Arg Gln
500 505 510

Glu Lys Ile Glu Lys Ala Val Leu Gly Met Glu Glu Leu Gln Asp Val
515 520 525

Asp Glu Leu Ile Glu Leu Leu Ser Val Asn Val Val Asn Pro Leu Gln
530 535 540

<210> SEQ ID NO 53
<211> LENGTH: 1797
<212> TYPE: DNA
<213> ORGANISM: Tribolium castaneum
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (41)..(1663)

<400> SEQUENCE: 53

acttgtgaat cagtcgtgcc cccacgagga tccacacacg atg ccg gcc aca ggc 55
Met Pro Ala Thr Gly
1 5

gaa gac caa gac ctg gtg caa gac ctc atc gag gag ccc gcc acc ttc 103
Glu Asp Gln Asp Leu Val Gln Asp Leu Ile Glu Glu Pro Ala Thr Phe
10 15 20

agc gac gcc gtc ctc tcc tcc gac gag gaa ctc ttc cac cag aag tgc 151
Ser Asp Ala Val Leu Ser Ser Asp Glu Glu Leu Phe His Gln Lys Cys
25 30 35

ccc aaa ccc gcc ccc att tac tcc ccg gtc tcg aaa ccg gtc tcc ttc 199
Pro Lys Pro Ala Pro Ile Tyr Ser Pro Val Ser Lys Pro Val Ser Phe
40 45 50

gag agc ctc ccc aac agg cgc ctc cac gag gag ttc ctc cgc agc tcg 247
Glu Ser Leu Pro Asn Arg Arg Leu His Glu Glu Phe Leu Arg Ser Ser
55 60 65

gtg gac gtc ctc ctc cag gag gcg gtg ttc gag gga acg aac cgc aag 295
Val Asp Val Leu Leu Gln Glu Ala Val Phe Glu Gly Thr Asn Arg Lys
70 75 80 85

aac cgg gtg ctg caa tgg cgg gag ccg gag gag ttg agg cgt ctg atg 343
Asn Arg Val Leu Gln Trp Arg Glu Pro Glu Glu Leu Arg Arg Leu Met
90 95 100

gac ttt ggg gtg cgg agt gcg ccc tcc acg cac gag gag ttg ttg gag 391
Asp Phe Gly Val Arg Ser Ala Pro Ser Thr His Glu Glu Leu Leu Glu
105 110 115

gtg ttg aag aag gtt gta act tat tcg gtt aaa acc gga cat ccg tac 439
Val Leu Lys Lys Val Val Thr Tyr Ser Val Lys Thr Gly His Pro Tyr
120 125 130

ttc gtg aac cag ttg ttc tcg gcg gtg gat ccg tac ggt ttg gtg gca 487
Phe Val Asn Gln Leu Phe Ser Ala Val Asp Pro Tyr Gly Leu Val Ala
135 140 145

caa tgg gcc acg gat gcg ctc aat ccg agt gtt tac acc tac gag gtt 535
Gln Trp Ala Thr Asp Ala Leu Asn Pro Ser Val Tyr Thr Tyr Glu Val
150 155 160 165

-continued

tcg	ccg	gtt	ttt	gtt	ctg	atg	gag	gaa	gtg	gtt	ttg	agg	gag	atg	agg	583
Ser	Pro	Val	Phe	Val	Leu	Met	Glu	Glu	Val	Val	Leu	Arg	Glu	Met	Arg	
				170					175					180		
gcc	att	gtg	ggg	ttc	gag	ggg	gga	aag	ggc	gat	ggg	att	ttt	tgc	cca	631
Ala	Ile	Val	Gly	Phe	Glu	Gly	Gly	Lys	Gly	Asp	Gly	Ile	Phe	Cys	Pro	
			185					190					195			
gga	ggg	tcc	att	gcc	aat	gga	tat	gcc	atc	agt	tgt	gcc	aga	tac	agg	679
Gly	Gly	Ser	Ile	Ala	Asn	Gly	Tyr	Ala	Ile	Ser	Cys	Ala	Arg	Tyr	Arg	
		200					205				210					
ttt	atg	ccc	gat	att	aag	aaa	aaa	ggc	ctc	cac	tct	ctc	ccc	cgt	ttg	727
Phe	Met	Pro	Asp	Ile	Lys	Lys	Lys	Gly	Leu	His	Ser	Leu	Pro	Arg	Leu	
	215				220						225					
gtc	ctc	ttc	acc	tct	gaa	gat	gcc	cac	tat	tcc	atc	aaa	aaa	ctc	gcc	775
Val	Leu	Phe	Thr	Ser	Glu	Asp	Ala	His	Tyr	Ser	Ile	Lys	Lys	Leu	Ala	
230					235					240				245		
tct	ttc	caa	ggc	atc	ggc	acc	gac	aac	gtc	tac	ttg	ata	cga	acg	gac	823
Ser	Phe	Gln	Gly	Ile	Gly	Thr	Asp	Asn	Val	Tyr	Leu	Ile	Arg	Thr	Asp	
			250					255						260		
gcc	cga	ggg	cgc	atg	gac	gtc	tcg	cac	ctg	gtg	gag	gaa	atc	gag	cgt	871
Ala	Arg	Gly	Arg	Met	Asp	Val	Ser	His	Leu	Val	Glu	Glu	Ile	Glu	Arg	
		265						270					275			
tcg	ctc	cgt	gaa	ggc	gcc	gct	cct	ttc	atg	gtc	agt	gcc	acc	gct	gga	919
Ser	Leu	Arg	Glu	Gly	Ala	Ala	Pro	Phe	Met	Val	Ser	Ala	Thr	Ala	Gly	
		280					285					290				
acc	aca	gtg	att	ggg	gcc	ttt	gac	ccc	atc	gaa	aaa	atc	gca	gat	gtg	967
Thr	Thr	Val	Ile	Gly	Ala	Phe	Asp	Pro	Ile	Glu	Lys	Ile	Ala	Asp	Val	
	295				300					305						
tgc	caa	aaa	tac	aaa	ctg	tgg	ttg	cac	gtg	gat	gcc	gcc	tgg	gga	ggg	1015
Cys	Gln	Lys	Tyr	Lys	Leu	Trp	Leu	His	Val	Asp	Ala	Ala	Trp	Gly	Gly	
310					315					320				325		
ggc	gcg	ctt	gtc	tct	gcc	aaa	cac	cgc	cac	ctc	ctc	aaa	ggg	att	gag	1063
Gly	Ala	Leu	Val	Ser	Ala	Lys	His	Arg	His	Leu	Leu	Lys	Gly	Ile	Glu	
		330						335						340		
agg	gcc	gac	tcg	gtc	acc	tgg	aac	cct	cac	aaa	ctc	cta	aca	gcc	ccc	1111
Arg	Ala	Asp	Ser	Val	Thr	Trp	Asn	Pro	His	Lys	Leu	Leu	Thr	Ala	Pro	
		345					350						355			
cag	caa	tgt	tcc	aca	ctt	tta	ctg	cga	cat	gag	ggg	gtc	ctc	gcc	gag	1159
Gln	Gln	Cys	Ser	Thr	Leu	Leu	Leu	Arg	His	Glu	Gly	Val	Leu	Ala	Glu	
		360				365						370				
gcg	cat	tcc	acg	aac	gcc	gct	tac	ctc	ttc	caa	aaa	gac	aaa	ttc	tac	1207
Ala	His	Ser	Thr	Asn	Ala	Ala	Tyr	Leu	Phe	Gln	Lys	Asp	Lys	Phe	Tyr	
	375				380					385						
gac	acc	aaa	tac	gac	acg	ggc	gac	aag	cac	atc	cag	tgc	ggc	cgc	agg	1255
Asp	Thr	Lys	Tyr	Asp	Thr	Gly	Asp	Lys	His	Ile	Gln	Cys	Gly	Arg	Arg	
	390				395					400				405		
gcc	gac	gtc	ctc	aag	ttc	tgg	ttc	atg	tgg	aag	gcg	aag	gga	aca	tca	1303
Ala	Asp	Val	Leu	Lys	Phe	Trp	Phe	Met	Trp	Lys	Ala	Lys	Gly	Thr	Ser	
		410						415						420		
ggg	ttg	gag	aaa	cac	gtc	gat	aaa	gtg	ttc	gaa	aat	gcg	aga	ttt	ttc	1351
Gly	Leu	Glu	Lys	His	Val	Asp	Lys	Val	Phe	Glu	Asn	Ala	Arg	Phe	Phe	
		425				430							435			
acc	gat	tgt	ata	aaa	aat	cgg	gaa	ggg	ttt	gaa	atg	gtg	ata	gcg	gag	1399
Thr	Asp	Cys	Ile	Lys	Asn	Arg	Glu	Gly	Phe	Glu	Met	Val	Ile	Ala	Glu	
	440				445							450				
ccc	gaa	tac	aca	aac	atc	tgc	ttt	tgg	tac	gtg	ccg	aag	agt	ctg	agg	1447
Pro	Glu	Tyr	Thr	Asn	Ile	Cys	Phe	Trp	Tyr	Val	Pro	Lys	Ser	Leu	Arg	
	455				460					465						

-continued

```

ggg cgc aag gac gaa gcc gat tac aaa gac aag ctg cat aag gtg gcc      1495
Gly Arg Lys Asp Glu Ala Asp Tyr Lys Asp Lys Leu His Lys Val Ala
470                               475                               480                               485

ccc agg att aag gag agg atg atg aag gag ggc tcc atg atg gtc acg      1543
Pro Arg Ile Lys Glu Arg Met Met Lys Glu Gly Ser Met Met Val Thr
                               490                               495                               500

tac cag gcg caa aag gga cac ccg aat ttt ttc agg att gtg ttc cag      1591
Tyr Gln Ala Gln Lys Gly His Pro Asn Phe Phe Arg Ile Val Phe Gln
                               505                               510                               515

aat tcg ggg ctt gac aag gct gat atg gtg cac ctt gtt gag gag att      1639
Asn Ser Gly Leu Asp Lys Ala Asp Met Val His Leu Val Glu Glu Ile
                               520                               525                               530

gag cgg ttg ggg agc gat ctt taa ggccttgaat ggtgctagtt gtagattgtg      1693
Glu Arg Leu Gly Ser Asp Leu
535                               540

taattaatgt aaaaagtatt atttaaaaaa tgtaaat ttt gatgtattta ttctcattag      1753
ttgtagttta ttcaaataaaa agtttaaaaa aaaaaaaaaa aaaa                        1797

<210> SEQ ID NO 54
<211> LENGTH: 540
<212> TYPE: PRT
<213> ORGANISM: Tribolium castaneum

<400> SEQUENCE: 54
Met Pro Ala Thr Gly Glu Asp Gln Asp Leu Val Gln Asp Leu Ile Glu
1                               5                               10                               15
Glu Pro Ala Thr Phe Ser Asp Ala Val Leu Ser Ser Asp Glu Glu Leu
20                               25                               30
Phe His Gln Lys Cys Pro Lys Pro Ala Pro Ile Tyr Ser Pro Val Ser
35                               40                               45
Lys Pro Val Ser Phe Glu Ser Leu Pro Asn Arg Arg Leu His Glu Glu
50                               55                               60
Phe Leu Arg Ser Ser Val Asp Val Leu Leu Gln Glu Ala Val Phe Glu
65                               70                               75                               80
Gly Thr Asn Arg Lys Asn Arg Val Leu Gln Trp Arg Glu Pro Glu Glu
85                               90                               95
Leu Arg Arg Leu Met Asp Phe Gly Val Arg Ser Ala Pro Ser Thr His
100                              105                              110
Glu Glu Leu Leu Glu Val Leu Lys Lys Val Val Thr Tyr Ser Val Lys
115                              120                              125
Thr Gly His Pro Tyr Phe Val Asn Gln Leu Phe Ser Ala Val Asp Pro
130                              135                              140
Tyr Gly Leu Val Ala Gln Trp Ala Thr Asp Ala Leu Asn Pro Ser Val
145                              150                              155                              160
Tyr Thr Tyr Glu Val Ser Pro Val Phe Val Leu Met Glu Glu Val Val
165                              170                              175
Leu Arg Glu Met Arg Ala Ile Val Gly Phe Glu Gly Gly Lys Gly Asp
180                              185                              190
Gly Ile Phe Cys Pro Gly Gly Ser Ile Ala Asn Gly Tyr Ala Ile Ser
195                              200                              205
Cys Ala Arg Tyr Arg Phe Met Pro Asp Ile Lys Lys Lys Gly Leu His
210                              215                              220
Ser Leu Pro Arg Leu Val Leu Phe Thr Ser Glu Asp Ala His Tyr Ser
225                              230                              235                              240
Ile Lys Lys Leu Ala Ser Phe Gln Gly Ile Gly Thr Asp Asn Val Tyr

```


-continued

245								250				255					
Leu	Ile	Arg	Thr	Asp	Ala	Arg	Gly	Arg	Met	Asp	Val	Ser	His	Leu	Val		
260								265				270					
Glu	Glu	Ile	Glu	Arg	Ser	Leu	Arg	Glu	Gly	Ala	Ala	Pro	Phe	Met	Val		
275								280				285					
Ser	Ala	Thr	Ala	Gly	Thr	Thr	Val	Ile	Gly	Ala	Phe	Asp	Pro	Ile	Glu		
290								295				300					
Lys	Ile	Ala	Asp	Val	Cys	Gln	Lys	Tyr	Lys	Leu	Trp	Leu	His	Val	Asp		
305								310				315					
Ala	Ala	Trp	Gly	Gly	Gly	Ala	Leu	Val	Ser	Ala	Lys	His	Arg	His	Leu		
325								330				335					
Leu	Lys	Gly	Ile	Glu	Arg	Ala	Asp	Ser	Val	Thr	Trp	Asn	Pro	His	Lys		
340								345				350					
Leu	Leu	Thr	Ala	Pro	Gln	Gln	Cys	Ser	Thr	Leu	Leu	Leu	Arg	His	Glu		
355								360				365					
Gly	Val	Leu	Ala	Glu	Ala	His	Ser	Thr	Asn	Ala	Ala	Tyr	Leu	Phe	Gln		
370								375				380					
Lys	Asp	Lys	Phe	Tyr	Asp	Thr	Lys	Tyr	Asp	Thr	Gly	Asp	Lys	His	Ile		
385								390				395					
Gln	Cys	Gly	Arg	Arg	Ala	Asp	Val	Leu	Lys	Phe	Trp	Phe	Met	Trp	Lys		
405								410				415					
Ala	Lys	Gly	Thr	Ser	Gly	Leu	Glu	Lys	His	Val	Asp	Lys	Val	Phe	Glu		
420								425				430					
Asn	Ala	Arg	Phe	Phe	Thr	Asp	Cys	Ile	Lys	Asn	Arg	Glu	Gly	Phe	Glu		
435								440				445					
Met	Val	Ile	Ala	Glu	Pro	Glu	Tyr	Thr	Asn	Ile	Cys	Phe	Trp	Tyr	Val		
450								455				460					
Pro	Lys	Ser	Leu	Arg	Gly	Arg	Lys	Asp	Glu	Ala	Asp	Tyr	Lys	Asp	Lys		
465								470				475					
Leu	His	Lys	Val	Ala	Pro	Arg	Ile	Lys	Glu	Arg	Met	Met	Lys	Glu	Gly		
485								490				495					
Ser	Met	Met	Val	Thr	Tyr	Gln	Ala	Gln	Lys	Gly	His	Pro	Asn	Phe	Phe		
500								505				510					
Arg	Ile	Val	Phe	Gln	Asn	Ser	Gly	Leu	Asp	Lys	Ala	Asp	Met	Val	His		
515								520				525					
Leu	Val	Glu	Glu	Ile	Glu	Arg	Leu	Gly	Ser	Asp	Leu						
530								535				540					

<210> SEQ ID NO 55

<211> LENGTH: 1356

<212> TYPE: DNA

<213> ORGANISM: Bacillus cereus

<400> SEQUENCE: 55

```

ttaaagttga gctaacatt ctttcattgt tttaacgata aaagtaaagt cttcctctgt      60
gatgcttaat ggaggtgcaa gctgcaaat attattgtaa cctgcaacag tgtcaccatt      120
tttaccaata attagacctt tttctttaca agcattgatg actttgttca tcttttcaat      180
ggaagccggt tcttttgttt gcttatcttc cactagttca atacctaaaa gaaggccttt      240
tccgcaaca tctcctacgt ttggatgctc ttttacatct tctagttcat ataacagtcg      300
ttcacccaat tctttggaac gttcaatgag tttctcattc tccataattt ctaaattctt      360

```

-continued

```

caaagctaag gcgcaagcag caggatttcc tccaaacgta tttacatggc ggaagcgatc 420
ataatcatca ctgcctacga atgcctcata aacctctcgt ctaactgctg ttgctgacaa 480
aggaagatac gcacttgtaa taccttttgc cattgtaatg atatctgggt tgacgccata 540
attcataaat ccaaacggct tcctgttcg tccaaatcca catataactt catcacaaat 600
gagcaacgca ccatgcttct cgcaaatttc ttttactttt tccatatatc catcaggagg 660
cattaaaatt cgcceccag taatgattgg ctccataatc acacgggcta ctgtttggct 720
taactcccat gtcatgacac gatcgatttc ctccagcactt gccagtgtat gaacatcctc 780
tggattgcga tacgtatcag gcggtgctac atgcaaaaaa ccttgctcta atggctcata 840
tttatacttt ctttgtgctt gccctgttgc tgcaagagca cccattgagt taccgtgata 900
agcgcggtag cgggaaataa acttatagcg tccatgatca cctttttgct gatgatattg 960
acgagcaatt ttaaatgctg ttctatttgc ttctgatcca ctgttagaaa agaaaatgac 1020
gtattcatca tccagccatt cattcaattt ctctgctaat ttaatggcag gaacatgact 1080
ttgtgtcaga gggaaatatg gcatttcttc aagttgctca aatgccgctc ttgcaagctc 1140
ttttcggcg tatccaacat tcacacacca aagaccagac ataccgtcta aataacgggt 1200
tccatcaata tccgtcacc atgcccttc tgcttttggtg ataattaaat tcgttggaact 1260
aggggcccgt cctctcatcg catgccaaag gtacttttca tctgtttttt tcaaactttg 1320
tgtttgcctc gtcacttgca caatcatcag ctccat 1356

```

<210> SEQ ID NO 56

<211> LENGTH: 451

<212> TYPE: PRT

<213> ORGANISM: *Bacillus cereus*

<400> SEQUENCE: 56

```

Met Glu Leu Met Ile Val Gln Val Thr Glu Gln Thr Gln Ser Leu Lys
1             5             10             15

Lys Thr Asp Glu Lys Tyr Leu Trp His Ala Met Arg Gly Ala Ala Pro
20          25          30

Ser Pro Thr Asn Leu Ile Ile Thr Lys Ala Glu Gly Ala Trp Val Thr
35          40          45

Asp Ile Asp Gly Asn Arg Tyr Leu Asp Gly Met Ser Gly Leu Trp Cys
50          55          60

Val Asn Val Gly Tyr Gly Arg Lys Glu Leu Ala Arg Ala Ala Phe Glu
65          70          75          80

Gln Leu Glu Glu Met Pro Tyr Phe Pro Leu Thr Gln Ser His Val Pro
85          90          95

Ala Ile Lys Leu Ala Glu Lys Leu Asn Glu Trp Leu Asp Asp Glu Tyr
100         105         110

Val Ile Phe Phe Ser Asn Ser Gly Ser Glu Ala Asn Glu Thr Ala Phe
115         120         125

Lys Ile Ala Arg Gln Tyr His Gln Gln Lys Gly Asp His Gly Arg Tyr
130         135         140

Lys Phe Ile Ser Arg Tyr Arg Ala Tyr His Gly Asn Ser Met Gly Ala
145         150         155         160

Leu Ala Ala Thr Gly Gln Ala Gln Arg Lys Tyr Lys Tyr Glu Pro Leu
165         170         175

Gly Gln Gly Phe Leu His Val Ala Pro Pro Asp Thr Tyr Arg Asn Pro

```

-continued

180					185					190					
Glu	Asp	Val	His	Thr	Leu	Ala	Ser	Ala	Glu	Glu	Ile	Asp	Arg	Val	Met
	195						200					205			
Thr	Trp	Glu	Leu	Ser	Gln	Thr	Val	Ala	Gly	Val	Ile	Met	Glu	Pro	Ile
	210					215					220				
Ile	Thr	Gly	Gly	Gly	Ile	Leu	Met	Pro	Pro	Asp	Gly	Tyr	Met	Glu	Lys
225					230					235				240	
Val	Lys	Glu	Ile	Cys	Glu	Lys	His	Gly	Ala	Leu	Leu	Ile	Cys	Asp	Glu
				245					250					255	
Val	Ile	Cys	Gly	Phe	Gly	Arg	Thr	Gly	Lys	Pro	Phe	Gly	Phe	Met	Asn
			260					265					270		
Tyr	Gly	Val	Lys	Pro	Asp	Ile	Ile	Thr	Met	Ala	Lys	Gly	Ile	Thr	Ser
	275						280					285			
Ala	Tyr	Leu	Pro	Leu	Ser	Ala	Thr	Ala	Val	Arg	Arg	Glu	Val	Tyr	Glu
	290					295					300				
Ala	Phe	Val	Gly	Ser	Asp	Asp	Tyr	Asp	Arg	Phe	Arg	His	Val	Asn	Thr
305					310					315				320	
Phe	Gly	Gly	Asn	Pro	Ala	Ala	Cys	Ala	Leu	Ala	Leu	Lys	Asn	Leu	Glu
				325					330					335	
Ile	Met	Glu	Asn	Glu	Lys	Leu	Ile	Glu	Arg	Ser	Lys	Glu	Leu	Gly	Glu
		340						345					350		
Arg	Leu	Leu	Tyr	Glu	Leu	Glu	Asp	Val	Lys	Glu	His	Pro	Asn	Val	Gly
	355						360					365			
Asp	Val	Arg	Gly	Lys	Gly	Leu	Leu	Leu	Gly	Ile	Glu	Leu	Val	Glu	Asp
	370				375						380				
Lys	Gln	Thr	Lys	Glu	Pro	Ala	Ser	Ile	Glu	Lys	Met	Asn	Lys	Val	Ile
385					390					395				400	
Asn	Ala	Cys	Lys	Glu	Lys	Gly	Leu	Ile	Ile	Gly	Lys	Asn	Gly	Asp	Thr
			405						410					415	
Val	Ala	Gly	Tyr	Asn	Asn	Ile	Leu	Gln	Leu	Ala	Pro	Pro	Leu	Ser	Ile
			420					425					430		
Thr	Glu	Glu	Asp	Phe	Thr	Phe	Ile	Val	Lys	Thr	Met	Lys	Glu	Cys	Leu
	435						440					445			
Ala	Gln	Leu													
	450														

<210> SEQ ID NO 57

<211> LENGTH: 747

<212> TYPE: DNA

<213> ORGANISM: Escherichia coli

<400> SEQUENCE: 57

```

atgatcgttt tagtaactgg agcaacggca ggttttggtg aatgcattac tcgtcgtttt    60
attcaacaag ggcataaagt tatcgccact ggccgtcgcc aggagcggtt gcaggagtta    120
aaagacgaac tgggagataa tctgtatatc gcccaactgg acgttcgcaa ccgcgccgct    180
attgaagaga tgctggcatc gcttcctgcc gagtggtgca atattgatat cctggtaaat    240
aatgccggct tggcgttggg catggagcct gcgcataaag ccagcgttga agactgggaa    300
acgatgattg ataccaacaa caaaggcctg gtatatatga cgcgcgccgt cttaccgggt    360
atggttgaac gtaatcatgg tcatattatt aacattggct caacggcagg tagctggccg    420
tatgccggtg gtaacgttta cggtgcgacg aaagcgtttg ttcgtcagtt tagcctgaat    480

```

-continued

```

ctgcgtacgg atctgcatgg tacggcgggtg cgcgtcacccg acatcgaacc gggctcgggtg 540
gggtggcacccg agttttccaa tgtccgcttt aaaggcgatg acggtaaagc ggaaaaaacc 600
tatcaaaata ccgttgcatg gacgccagaa gatgtcagcg aagccgtctg gtgggtgtca 660
acgctgcctg ctcacgtcaa tatcaatacc ctggaaatga tgccggttac ccaaagctat 720
gccggactga atgtccaccg tcagtaa 747

```

```

<210> SEQ ID NO 58
<211> LENGTH: 248
<212> TYPE: PRT
<213> ORGANISM: Proteobacteria

```

```

<400> SEQUENCE: 58

```

```

Met Ile Val Leu Val Thr Gly Ala Thr Ala Gly Phe Gly Glu Cys Ile
1           5           10          15
Thr Arg Arg Phe Ile Gln Gln Gly His Lys Val Ile Ala Thr Gly Arg
20          25          30
Arg Gln Glu Arg Leu Gln Glu Leu Lys Asp Glu Leu Gly Asp Asn Leu
35          40          45
Tyr Ile Ala Gln Leu Asp Val Arg Asn Arg Ala Ala Ile Glu Glu Met
50          55          60
Leu Ala Ser Leu Pro Ala Glu Trp Cys Asn Ile Asp Ile Leu Val Asn
65          70          75          80
Asn Ala Gly Leu Ala Leu Gly Met Glu Pro Ala His Lys Ala Ser Val
85          90          95
Glu Asp Trp Glu Thr Met Ile Asp Thr Asn Asn Lys Gly Leu Val Tyr
100         105         110
Met Thr Arg Ala Val Leu Pro Gly Met Val Glu Arg Asn His Gly His
115        120        125
Ile Ile Asn Ile Gly Ser Thr Ala Gly Ser Trp Pro Tyr Ala Gly Gly
130        135        140
Asn Val Tyr Gly Ala Thr Lys Ala Phe Val Arg Gln Phe Ser Leu Asn
145        150        155        160
Leu Arg Thr Asp Leu His Gly Thr Ala Val Arg Val Thr Asp Ile Glu
165        170        175
Pro Gly Leu Val Gly Gly Thr Glu Phe Ser Asn Val Arg Phe Lys Gly
180        185        190
Asp Asp Gly Lys Ala Glu Lys Thr Tyr Gln Asn Thr Val Ala Leu Thr
195        200        205
Pro Glu Asp Val Ser Glu Ala Val Trp Trp Val Ser Thr Leu Pro Ala
210        215        220
His Val Asn Ile Asn Thr Leu Glu Met Met Pro Val Thr Gln Ser Tyr
225        230        235        240
Ala Gly Leu Asn Val His Arg Gln
245

```

```

<210> SEQ ID NO 59
<211> LENGTH: 987
<212> TYPE: DNA
<213> ORGANISM: Aspergillus pseudoterreus

```

```

<400> SEQUENCE: 59

```

```

ggttgtagca gcgtaaaccac atggatagtt aaataatcgg atgtacaccc actgttggaa 60

```

-continued

```

atgacggggg cctacaacac gagattatct gatecaattt ctgttcgttg gcattctatc 120
attcgacgcg aaaattgtcc tattaaattg accatgacca aacaatctgc ggacagcaac 180
gcaaagtcag gagttacgtc cgaatatgtt cattgggcat ccaacctggc cactgacgac 240
atcccttcgg acgtattaga aagagcaaaa taccttattc tcgacggtat tgcattgtgc 300
tgggttggtg caagagtgcc ttggtcagag aagtatgttc aggcaacgat gagctttgag 360
ccgcgggggg cctgcagggt gattggatat ggacaggtaa attttattca ctctagacgg 420
tccacaaagt atactgaaga tccttcgtat agaaactggg gcctgttgca gcagccatga 480
ccaattccgc ttctacacag gctacggagc ttgacgacta ccacagcgaa gccccctac 540
actctgcaag cattgtcctt cctgcggtct ttgcagcaag tgaggcttta gccgagcagg 600
gcaaaacaat ttccggtata gatgttattc tagccgccat tgtggggttt gaatctggcc 660
cacggatcgg caaagcaatc tacggatcgg acctcttgaa caacggctgg cattgtggag 720
ctgtgtatgg cgctccagcc ggtgcgctgg ccacaggaaa gctcctcggc ctaactccag 780
actccatgga agatgctctc ggaattgctg gcacgcaagc ctgtggttta atgtcggcgc 840
aatacggagg catggttaaag cgtgtgcaac acggattcgc agcgcgtaat ggtcttcttg 900
ggggactgtt ggcccatggt ggtgtacgag caatgaaagg tgctctggag agatcttacg 960
gcggtttcct caagatgttc accaagg 987

```

```

<210> SEQ ID NO 60
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate A. pseudoterreus 5'-cadA
gene

```

```

<400> SEQUENCE: 60
ccctcgaggt cgacgggtatc gatagatatc ggttgtagca gcgtaaacac 50

```

```

<210> SEQ ID NO 61
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate A. pseudoterreus 5'-cadA
gene

```

```

<400> SEQUENCE: 61
tctttcatag tagccttggt gaacatcttg agg 33

```

```

<210> SEQ ID NO 62
<211> LENGTH: 813
<212> TYPE: DNA
<213> ORGANISM: Aspergillus niger

```

```

<400> SEQUENCE: 62
ctactatgaa agaccgcgat gggccgatag tagtagttac ttccattaca tcatctcatc 60
cgccccgttc ctcgcctccg cggcagtcta cgggtaggat cgtagcaaaa acccggggga 120
tagaccgctc gtcccgagct ggagttccgt ataacctagg tagaaggat caattgaacc 180
cgaacaactg gcaaaacatt ctcgagatcg taggagttag taccggcgt gatggagggg 240
gagcacgctc attggtccgt acggcagctg ccgaggggga gcaggagatc caaatatcgt 300

```

-continued

gagttctctg ctttgccggg tgtatgaaac cggaaaggac tgctggggaa ctggggagcg	360
gcgcaagcgg ggaatcccag ctgacaattg acccatcctc atgcctgggc agagcttgag	420
gtagcttttg ccccgctgtg ctecccggtg tgcgcattcg actgggcgcg gcattctgtg	480
ctctccagg agcggaggac ccagtagtaa gtaggcctga cctggctggt gcgtcagtc	540
agaggttccc tcccctaccc tttttctact tcccctcccc cgccgctcaa cttttcttcc	600
ccttttactt tctctctctc ttctcttca tccatcctct cttcatcact tccctcttcc	660
cttcaccaa ttcattcttc aagtgaagtct tctctcccat ctgtccctcc atctttccca	720
tcatcatctc cttcccagc tctctccctc ctctcgtctc ctacgaagc ttgactaacc	780
attaccccg ccatagaca catctaaaca atg	813

<210> SEQ ID NO 63
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. niger gpdA promoter

<400> SEQUENCE: 63

atgttcacca aggctactat gaaagaccgc gatg	34
---------------------------------------	----

<210> SEQ ID NO 64
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. niger gpdA promoter

<400> SEQUENCE: 64

cgccggtggc gggcattggt tagatgtgtc tatgtg	36
---	----

<210> SEQ ID NO 65
 <211> LENGTH: 1617
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: panD cDNA of Tribolium castaneum with codon optimization for A. pseudoterreus

<400> SEQUENCE: 65

cccgccaccg gcgaggacca ggacctggtg caggacctga tcgaggaaac cgccaccttc	60
tccgacgcgg tctgtctctc cgacgaggaa ctgttccacc agaagtgtcc caagccggct	120
ccgatctaca gccccgtcag caagcccgtc agcttcgagt ccctgccgaa ccgccgctg	180
cacgaagagt tctccgctc ctccgtcgac gtctgtgtgc aagaggccgt gttcgagggc	240
accaaccgca agaaccgct cctgcagtgg cgcgagccg aagaactgcg ccgcctgatg	300
gacttcggcg tccgcagcgc cccgtccacg catgaggaaac tgctcgaggt cctgaagaag	360
gtcgtacct actccgtcaa gaccggccat ccgtacttcg tcaaccagct gttctccgcc	420
gtcgatccct acggcctggg cgcccagtgg gccaccgacg cgctgaacct ctccgtctac	480
acctacgagg tcagcccggt gttcgtctcg atggaagagg tcgtctcgcg cgagatgcgc	540
gccatcgtcg gcttcgaagg cggcaaaggc gacggcatct tctgccctgg cggtctgatc	600

-continued

```

gccaaacggct acgccatcag ctgcgcccgc taccgcttca tgcccacat caagaagaag 660
ggcctgcaact cctgcgcgag cctggctcctg ttcacctccg aggacgccc ctactcgatc 720
aagaagctgg cctcgttcca aggcacggc accgacaacg tctacctgat cgcaccgac 780
gtcgcgggtc gcattggacgt cagccacctg gtcgaagaga tcgagcgctc cctccgcgag 840
ggcgctgccc cgttcattgt cagcgccacc gccggcacca ccgctcctcg cgcttcgat 900
cccatcgaga agatcgccga cgtctgccag aagtacaagc tctgggtgca cgtcgacgcc 960
gcctggggcg gagcgctctt ggtgtccgcc aagcaccgcc atctgtgaa gggcctcgag 1020
cgcgccgact ccgtcacctg gaattccccc aagctgtgca ccgctccgca gcagtgcagc 1080
acctgtctgc tgcgccacga gggcgctcctg gccgaggcgc actccaccaa cgccgcctac 1140
ctgttccaga aggacaagtt ctacgacacc aagtagaca ccggcgacaa gcacatccag 1200
tgcgcccgtc gcgcgcagct gctgaagttc tggttcatgt ggaaggccaa gggcacctcc 1260
ggcctcgaga agcacgtgga caaggtgttc gagaacgccc gcttcttcac cgactgcac 1320
aagaaccgtg agggcttcga gatggtgatc gccgagcctg agtacacaa catctgtttc 1380
tggtacgtcc ccaagagcct gcgcggacgc aaggacgagg ccgactacaa ggacaagctg 1440
cacaaggtcg cccctcgcat caaagaacgc atgatgaagg aaggctccat gatggtcacc 1500
taccaggcgc agaagggcca tccgaatttc ttccgcatcg tctttcagaa ctccggcctg 1560
gacaaggcgc acatggtcca tctggtcgag gaaatcgaa gcctgggctc cgacctc 1617

```

```

<210> SEQ ID NO 66
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate panD cDNA of Tribolium
castaneum with codon optimization for A. pseudoterreus

```

```

<400> SEQUENCE: 66

```

```

catctaaaca atgcccgcga ccggcgagga cca 33

```

```

<210> SEQ ID NO 67
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate panD cDNA of Tribolium
castaneum with codon optimization for A. pseudoterreus

```

```

<400> SEQUENCE: 67

```

```

atccaaccca tcagaggtcg gagcccaggc gttcg 35

```

```

<210> SEQ ID NO 68
<211> LENGTH: 1040
<212> TYPE: DNA
<213> ORGANISM: Aspergillus niger

```

```

<400> SEQUENCE: 68

```

```

tgatgggttg gatgacgatg acttcatgtg attttgttat ttagaatatt ttatatttcc 60
ttttttctct ctcaccaccc atccccttaa cactcttget tcatttgctt cagatttctc 120
ggttttctct tttttctctt ccccagttat ccaatatatc tttgctagac cggcctgcgc 180
cctggcatgc atcataaaat catgtccgtt ggtcatcatc tgttttgtat atccgtcata 240

```

-continued

taaagtattc ttttattccc tccccctcg gtcgtcttcc gctgtcccgc ttcctacctc	300
cgggtttatag agcatgggtc atctcttccg tacatttccg ttggtactag cattttatgtc	360
ttcagctagt atagaagctg ccgcagttgt tcgcttacta cctgcctaag tccttaactt	420
tttaaagtgt ttaacctata cgtagtgtta aacgagtact gggaggtggg gaggtagaaa	480
atgtttctgca cgggcagtggt gtatttggtg gtgtgtaagg cggttattta tcaggctgac	540
gctaagact tctatgggag cagtatggga tcgcggctca tagaagtaca caaaatctaa	600
gagtcgtttg ataattaatt gattcccggc agggctctct tgggattgag agaactggtt	660
actttgattt gagatattgt aaagcttaag gctcttaaca cgtacgagcg aaacagcagg	720
ggggaaatcg ggaaaagggt cgtggggtga ataaaaagt tgaataaga cactgtatct	780
tgctgggggt gaataaagag agaataaaag agaggtaaat tccactcagc cccttttctt	840
cgctctccaa acatcaaact ccgccggcgc acccacagga tcccgaacaa gtggaagata	900
tgtgccggtc cagacccttc gcacagctaa aagcagacct tcataagcgt ttccgggtag	960
tattcgacac cctgaactgg cagctcgggg acacaactgt ttttgatata caagaacaca	1020
caccacccat ctaggactca	1040

<210> SEQ ID NO 69
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate bidirectional terminator
 from A. niger elf3/multifunctional chaperone

<400> SEQUENCE: 69

gggctccgac ctctgatggg ttggatgacg atg	33
--------------------------------------	----

<210> SEQ ID NO 70
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate bidirectional terminator
 from A. niger elf3/multifunctional chaperone

<400> SEQUENCE: 70

tctggcccag ctctgagtc tagatgggtg gtg	33
-------------------------------------	----

<210> SEQ ID NO 71
 <211> LENGTH: 1350
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: codon optimized synthetic cDNA of
 beta-alanine-pyruvate aminotransferase (BAPAT) of Bacillus cereus

<400> SEQUENCE: 71

gagctgggcc agacattcct tcatagtctt gacgatgaag gtgaagtcct cttcggtgat	60
ggacagcgga ggggcgagct gcaggatggt gttgtagccg gccacgggtg cgcggttctt	120
gccgatgatc agacccttct ctttgaggc gttgatgacc ttgttcattt tttegatgga	180
ggcgggctct ttggtctgct taccctcgac gagttcgata cccagcagga ggcccttgcc	240
gcggacgtcc ccgacgttgg ggtgctcttt gacgtcctcc aactcgtaca gcaggcgctc	300
gcccagttct ttggaccgct cgatgagctt ctcgttttcc atgatctcga ggttcttcag	360

-continued

```

ggccagcgcg caggcggcag ggttgccgcc gaaggtgttg acatggcgga agcggtcgta 420
gtcgtcggag ccgacgaagg cctcgtagac ctccggcgcc acccggtggg cagacagcgg 480
caggtaggcc gaggtgatac ccttgcccat ggtgataatg tcgggcttga cgccgtagtt 540
catgaagccg aagggcttgc cgtgcgacc gaagccgag atgacctcgt cgcagatcag 600
cagggcgccg tgcttttcgc agatctcttt gaccttttcc atgtagccgt ccggcgccat 660
caggatgccca ccaccggtga tgatgggttc catgatgacg ccggcgacgg tctgggacag 720
ctcccaggtc atgacgcggt cgattttctc ggccggaggcc aggggtgtgca cgtcctcggg 780
gttgcgatag gtgtccggag gggccacgtg caggaagccc tgaccgaggg gctcgtactt 840
gtacttgccg tgggcctgac cgttcgcggc cagggcaccc atggagtgc cgtggtaggc 900
gcggtagcga gagatgaact tgtagcggcc gtggtcaccc ttctgctggg ggtactggcg 960
ggcgatcttg aagggcggtt cgttgccctc cgagccggag ttggagaaga agatgacgta 1020
ctcgtcgtcc agccactcgt tcagcttctc ggccagcttg atggcgggga cgtgcgactg 1080
cgtcagcggg aagtacggca tctcttcag ctgctcgaag gcagcgcgag ccagctcttt 1140
gcggccgtag ccgacgttga cgcaccacag gccggacatg ccgtccaggt agcggttgcc 1200
gtcgatgtcg gtgacccacg cgccctcggc cttggtgatg atcaggttgg tcggactcgg 1260
agcggcaccc cgcattggct gccacaggta cttctcgtcg gttttcttca ggctctgggt 1320
ctgctcggtg acctggacga tcacagttc 1350

```

```

<210> SEQ ID NO 72
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate codon optimized
synthetic cDNA of beta-alanine-pyruvate aminotransferase (BAPAT)
of Bacillus cereus

```

```

<400> SEQUENCE: 72
catctaggac tcagagctgg gccagacatt ccttc 35

```

```

<210> SEQ ID NO 73
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate codon optimized
synthetic cDNA of beta-alanine-pyruvate aminotransferase (bapat)
of bacillus cereus

```

```

<400> SEQUENCE: 73
gtccatcaac atggaactga tgatcgtcca gggtcac 36

```

```

<210> SEQ ID NO 74
<211> LENGTH: 704
<212> TYPE: DNA
<213> ORGANISM: Aspergillus niger

```

```

<400> SEQUENCE: 74
catgttgatg gactggaggg ggatgagtta tggatcagtg aaactgggag aaaacaaaga 60
tggcaaaggg agaacatggc ccagatatag gaaaaaacgg aggaggcaaa aatgtaagcg 120
ctccggactt gctgtttcgg tgtgcactag cagcagcggg ggggaagggt gtgagtgttc 180

```

-continued

accgaggacc caaaaagaat gagcggatgg cggatgagtg acggagaagg gaaggacggg	240
gggggaatta gaggtggaga ggtccgatcc atcaaataga ccaggctcgg cacagccaag	300
tttcccaaat gatcaactaa tcaatgggac ttggtgctaa atccggagat gccagatcat	360
tgatagacag acaggatgga gtgatggcat atagacagga ggatggatgg atggatagat	420
ggaggggtca agcacaacat ggtgggatga tggcggggtc atgactagca gctaagagga	480
agaagaggag gatgaaatgg acagagaaaag atgggagggg tgataaaatg agtatatggg	540
acaagtcata cttacaggac cttgaagatg gtggttgtag tatctaagaa aggccttttt	600
tgagagtact cttaacacaa gaggaggagg gaggaggggg aagtagtaga taaataataa	660
acacgaccac agacttgcta caggctactt cttgtaagct cgag	704

<210> SEQ ID NO 75
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. niger eno1 promoter

<400> SEQUENCE: 75

cgatcatcag ttccatgttg atggactgga ggg	33
--------------------------------------	----

<210> SEQ ID NO 76
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. niger eno1 promoter

<400> SEQUENCE: 76

gaactagtgg atccccggg ctgcgttaac tcgagcttac aagaagtagc c	51
---	----

<210> SEQ ID NO 77
 <211> LENGTH: 885
 <212> TYPE: DNA
 <213> ORGANISM: Aspergillus nidulans

<400> SEQUENCE: 77

tctgtacagt gaccggtgac tctttctggc atgcggagag acggacggac gcagagagaa	60
gggctgagta ataagccact ggccagacag ctctggcggc tctgaggtgc agtggatgat	120
tattaatccg ggaccggccg cccctccgcc ccgaagtgga aaggctgggtg tgcccctcgt	180
tgaccaagaa tctattgcat catcggagaa tatggagctt catcgaatca ccggcagtaa	240
gcgaaggaga atgtgaagcc aggggtgtat agccgtcgcc gaaatagcat gccattaacc	300
taggtacaga agtccaattg cttccgatct ggtaaaagat tcacgagata gtaccttctc	360
cgaagtaggt agagcgagta cccggcgcggt aagctcccta attggcccat ccggcatctg	420
tagggcgctc aaatatcgtg cctctcctgc ttgcccgggt gtatgaaacc ggaaaggccg	480
ctcaggagct ggccagcggc gcagaccggg aacacaagct ggcagtcgac ccatccggtg	540
ctctgcactc gacctgtgta ggtccctcag tccctggtag gcagctttgc cccgtctgtc	600
cgcccggtgt gtcggcgggg ttgacaaggt cgttgcgta gtccaacatt tgttgccata	660
ttttctgct ctcaccaaca gctgctcttt tcttttctct ttcttttccc atcttcagta	720

-continued

tattcatctt cccatccaag aaccttttatt tcccctaagt aagtactttg ctacatccat	780
actccatcct tcccacccct tattcctttg aacctttcag ttcgagcttt cccacttcat	840
cgcagcttga ctaacagcta ccccgcttga gcagacatca ccatg	885

<210> SEQ ID NO 78
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. nidulans gpdA promoter

<400> SEQUENCE: 78

ccctcgaggt cgacggtatc gatagttaac tctgtacagt gaccggtgac	50
--	----

<210> SEQ ID NO 79
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. nidulans gpdA promoter

<400> SEQUENCE: 79

tgaccagcac gatcatgggtg atgtctgctc aag	33
---------------------------------------	----

<210> SEQ ID NO 80
 <211> LENGTH: 741
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: codon optimized synthetic cDNA of E. coli HPDH

<400> SEQUENCE: 80

atcgtgctgg tcacggggcg gaccgcccgt ttcggcgagt gcatcaccgg ccgcttcac	60
cagcagggcc acaagggtgat cgctaccgga cgccgccaag agcgctcca agagctgaag	120
gatgagctgg ggcacaacct gtacattgcc cagctggacg tgcgcaaccg ggctgccatc	180
gaagaaatgc tcgcctcgct gcccgccgag tgggtgcaaca tcgacatcct ggtcaacaac	240
gccggtctgg ccctcggeat ggaaccggcg cacaaggcca gcgtcgagga ctgggaaacc	300
atgatcgaca ccaacaacaa gggactcgtc tacatgaccc gcgctgtgct gcccggcacg	360
gtcgagcgca accacggcca catcatcaac atcggtccca ccgctggcag ctggccctac	420
gctggcgga acgtctatgg cgcgaccaag gcgttcgtcc gccagttctc cctgaacctg	480
cgcaccgacc tgcacggcac cgccgtccgc gtgaccgaca ttgagcccg tctggctggc	540
ggcaccgagt tcagcaacgt ccgcttcaag ggcgacgacg gcaaggccga gaaaacctac	600
cagaacacgg tcgctctgac ccctgaggat gtcagcgagg ccgtctggtg ggctcagcact	660
ctgcccgcg acgtcaacat caacaccctc gagatgatgc ccgtcacgca gtcctacgcc	720
ggcctgaacg tccaccgcca a	741

<210> SEQ ID NO 81
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate a codon optimized

-continued

synthetic cDNA of E. coli HPDH

<400> SEQUENCE: 81

agacatcacc atgategtgc tggtcacggg cgc 33

<210> SEQ ID NO 82
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate a codon optimized
synthetic cDNA of E. coli HPDH

<400> SEQUENCE: 82

gccatcggtc ctattggcgg tggacgttca ggc 33

<210> SEQ ID NO 83
<211> LENGTH: 473
<212> TYPE: DNA
<213> ORGANISM: Aspergillus nidulans

<400> SEQUENCE: 83

taggaccgat ggctgtgtag aagtactcgc cgatagtggg aaccgacgcc ccagcactcg 60
tccgagggca aaggaataga gtagatgccg accgcgggat ccacttaacg ttactgaaat 120
catcaaacag cttgacgaat ctggatataa gatcgttggt gtcgatgtca gctccggagt 180
tgagacaaat ggtgttcagg atctcgataa gatacgttca tttgtccaag cagcaaagag 240
tgccttctag tgatttaata gtcctatgtc aacaagaata aaacgcgttt tcgggtttac 300
ctcttcacga tacagctcat ctgcaatgca ttaatgcatt gactgcaacc tagtaacgcc 360
ttcaggctcc ggcgaagaga agaatagctt agcagagcta ttttcatttt cgggagacga 420
gatcaagcag atcaacgggc gtcaagagac ctacgagact gaggaatccg ctc 473

<210> SEQ ID NO 84
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate the trpC terminator of
A. nidulans

<400> SEQUENCE: 84

cgtccaccgc caataggacc gatggctgtg tag 33

<210> SEQ ID NO 85
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate the trpC terminator of
A. nidulans

<400> SEQUENCE: 85

gaactagtgg atcccccggg ctgcagagcg gattcctcag tctcg 45

<210> SEQ ID NO 86
<211> LENGTH: 510
<212> TYPE: DNA
<213> ORGANISM: Aspergillus nidulans

<400> SEQUENCE: 86

-continued

cacctcgaggt cgacgggtatc gatataggac cgatggctgt gtagaagtac tcgccgatag	60
tggaaaccga cgcgccagca ctcgccagag ggcaaaggaa tagagtagat gccgaccgcg	120
ggatccactt aacgttactg aaatcatcaa acagcttgac gaatctggat ataagatcgt	180
tggtgtcgat gtcagctccg gagttgagac aaatgggtgtt caggatctcg ataagatacg	240
ttcatttgtc caagcagcaa agagtgcctt ctagtgtttt aatagctcca tgtcaacaag	300
aataaaacgc gttttcgggt ttacctcttc cagatacagc tcctctgcaa tgcattaatg	360
cattgactgc aacctagtaa cgccttcagg ctccggcgaa gagaagaata gcttagcaga	420
gctattttca ttttcgggag acgagatcaa gcagatcaac ggctcgtaag agacctacga	480
gactgaggaa tccgctctct gacagacggg	510

<210> SEQ ID NO 87
 <211> LENGTH: 44
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate a trpC terminator of
 A. nidulans

<400> SEQUENCE: 87

cacctcgaggt cgacgggtatc gatataggac cgatggctgt gtag	44
--	----

<210> SEQ ID NO 88
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate a trpC terminator of
 A. nidulans

<400> SEQUENCE: 88

cccgtctgtc agagagcgga ttctctagtc tcg	33
--------------------------------------	----

<210> SEQ ID NO 89
 <211> LENGTH: 2005
 <212> TYPE: DNA
 <213> ORGANISM: Aspergillus oryzae

<400> SEQUENCE: 89

tctgacagac gggcaattga ttacgggac ccatggtaa cgaaatgtaa aagctaggag	60
atcgteccgc gatgtcagga tgatttcaact tgtttcttgt ccggctcacc ggtcaaagct	120
aaagaggagc aaaaggaacg gatagaatcg ggtgccgctg atctatacgg tatagtgcgc	180
ttatcacgtt gactcaaccc atgctattta actcaacccc tccttctgaa cccaccatc	240
ttcttctttt tcctctcatc ccacacaatt ctctatctca gatttgaatt ccaaaagtcc	300
tcggacgaaa ctgaacaagt ctctctcct tcgataaacc tttggtgatt ggaataactg	360
accatcttct atagtccca aaccaaccga caatgtaaat aactcctcg attagccctc	420
tagagggcat acgatggaag tcatggaata cttttggctg gactctcaca atgatcaagg	480
tatcttaggt aacgtctttg gcgtgggccc gtgttcgttc ccagtcacg atgcattcac	540
atgcctctcc taagctgggc cctagactct aggatcctag tctagaagga catggcatcg	600
atggactggg ttcgttctga gattatacgg ctaaaacttg atctggataa taccagcgaa	660
aagggtcatg cttctctctg ttcttctgt tgatggaatg gctaacagat gatagtcatt	720

-continued

```

gcaacttgaa acatgtctcc tccagctgcc atctacgaac ccaactgtggc cgctaccggc 780
ctcaagggtta aggtcgtggt ttctgagacc gtccccgttg agggagcttc tcagaccaag 840
ctgttggaacc atttcggtgg caagtgggac gaggttcaagt tcgcccctat ccgcgaaagc 900
caggctctctc gtgccatgac cagacgttac ttgaggacc tggacaagta cgctgaaagt 960
gacgttgtea ttgttggtgc tggttcctgc ggtctgagca ctgcgtacgt cttggccaag 1020
gtcgtcccg acctgaagat tgctatcgtc gaggccagcg tctctcctgg tcagtagtcc 1080
atgatggatt gccttgcaact cagctttccg gaactaacgt gcaatagggt gcggtgctg 1140
gttggtggc caactctttt ctgctatggt catgcgccgt cccgcggaag tcttctgaa 1200
cgagctgggt gtctcttacg aagaggacgc aaaccccaac tacgttgctc tcaagcacgc 1260
ctccctgttt acctgcacac tcatgtcgaa ggttctctcc tccccaatg tcaagctctt 1320
caatgctacc gctgttgagg acttgatcac ccgtccgacc gagaacggca accccagat 1380
tgctggtgtt gtcgtcaact ggacgctggt cacccttcac cacgatgac actcctgcat 1440
ggaccccaac actatcaacg ctctgtcat catcagtacc actggtcacg atgggccatt 1500
cggcgccctc tgtgcgaagc gcttggtgct catgggcagc gtcgacaagc taggtggcat 1560
gcgtggtctc gacatgaact cggccgagga tgccatcgtc aagaacaccc gcgaggttac 1620
taagggcttg ataateggcg gtatggagct gtctgaaatt gatggcttta accgcatggg 1680
ccctaccttc ggtgccatgg ttctcagtgg tgtcaaggct gccgaggagg cattgaaggt 1740
gttcgacgag cgtcagcgcg agtgtgctga gtaaatgact cactaccga atgggttcag 1800
tgcatgaacc ggatttgtct tacggtcttt gacgataggg gaatgatgat tatgtgatag 1860
ttctgagatt tgaatgaact cgtagctcg taatccacat gcatatgtaa atggctgtgt 1920
cccgtatgta acggtggggc attctagaat aattatgtgt aacaagaaag acagtataat 1980
acaaacaaag atgcaagagc ggctc 2005

```

```

<210> SEQ ID NO 90
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate the A. oryzae ptrA
selection marker gene

```

```

<400> SEQUENCE: 90

```

```

gaggaatccg ctctctgaca gacgggcaat tgattac 37

```

```

<210> SEQ ID NO 91
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate the A. oryzae ptrA
selection marker gene

```

```

<400> SEQUENCE: 91

```

```

gaatgttgct gaggagccgc tcttgcatct ttg 33

```

```

<210> SEQ ID NO 92
<211> LENGTH: 908
<212> TYPE: DNA
<213> ORGANISM: Aspergillus pseudoterreus

```

-continued

<400> SEQUENCE: 92

```

ctcagcaaca ttcgccatgt tcatgtacag ctttcaacgg cctcgaacag tcactgtgga      60
tggataccag aggagagacc catcagttca atcgcagggc agatgagtgt cgcatacatt      120
ctcgccgtcc agctggtoga ccagcaatgt cttttgtccc agttttctga gtttgatgac      180
aacctggaga ggccagaagt ttgggatctg gccaggaagg ttacttcac tcaaagcgaa      240
gagtttgatc aagacggcaa ctgtctcagt gcgggtcgcg tgaggattga gttcaacgat      300
ggttcttcta ttacggaaag tgctgagaag cctcttggtg tcaaagagcc catgccaaac      360
gaacggattc tccacaaata ccgaaccctt gctggtagcg tgacggacga atcccgggtg      420
aaagagattg aggatcttgt cctcggcctg gacaggtcca ccgacattag cccattgctg      480
gagctgctga attgccccgt gaaatcgcca ctggtataaa tgggaagcga tatggaaaca      540
tttcatgtca cgggcacaaa ttctaggtca tatcgtacct ggatggtgaa accaccagcg      600
gtttagcaga tagaagatag actccttctg ctctgcgttg cgtcttgaat ttagttcggt      660
cactggctta agaacttaga atgcaataca gtctctctta tttcttatta aaatcacgta      720
ttccacattc cggcgactgg aggatacgaa agcagtgttg gtggtgctcc ccgtaatgga      780
tatgattttg ctgactggac tattctatga ccattccctc caacggagat cctttctcga      840
cactttagat gttgacgctg tctggaggaa ctacttttgc gctgcaaaga ctatgagcag      900
tggagctg                                     908

```

<210> SEQ ID NO 93

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to isolate an *A. pseudoterreus* 3'-cad1 gene fragment

<400> SEQUENCE: 93

```

gcaagagcgg ctctcagca acattcgcca tgttc                                     35

```

<210> SEQ ID NO 94

<211> LENGTH: 49

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to isolate an *A. pseudoterreus* 3'-cad1 gene fragment

<400> SEQUENCE: 94

```

gaactagtgg atccccggg ctgcacagct ccaactgtca tagtctttg                     49

```

<210> SEQ ID NO 95

<211> LENGTH: 2099

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: combination of SEQ ID NOS: 77, 80, and 83

<400> SEQUENCE: 95

```

tctgtacagt gaccggtgac tctttctggc atcgcgagag acggacggac gcagagagaa      60
gggctgagta ataagccact ggccagacag ctctggcggc tctgaggtgc agtggatgat      120
tattaatccg ggaccggcgg cccctccgcc ccgaagtgga aaggctggtg tgcccctcgt      180

```

-continued

tgaccaagaa tctattgcat catcgagaa tatggagctt catcgaatca cggcgagtaa	240
gcgaaggaga atgtgaagcc aggggtgtat agccgtcggc gaaatagcat gccattaacc	300
taggtacaga agtccaattg cttccgatct ggtaaaagat tcacgagata gtaccttctc	360
cgaagtaggt agagcgagta cccggcgcgt aagctcccta attggcccat cggcgatctg	420
tagggcgctc aaatatcgtg cctctcctgc ttgcccgggt gtatgaaacc ggaaaggccg	480
ctcaggagct ggccagcggc gcagaccggg aacacaagct ggagtcgac ccacccgggtg	540
ctctgcactc gacctgctga ggtccctcag tccctggtag gcagctttgc cccgtctgtc	600
cggccgggtg gtcggcgggg ttgacaaggt cgttgctgca gtccaacatt tgttgccata	660
ttttctgct ctcaccacca gctgctcttt tcttttctct ttcttttccc atcttcagta	720
tattcatctt cccatccaag aacctttatt tccctaagt aagtactttg ctacatccat	780
actccatcct tcccatcctt tattcctttg aacctttcag ttcgagcttt cccacttcat	840
cgcagcttga ctaacagcta ccccgcttga gcagacatca ccatgatcgt gctggtcacg	900
ggcgcgaccg cgggtttcgg cgagtgcatc acccgccgct tcatccagca gggccacaag	960
gtgatcgcta ccggacgcgc ccaagagcgc ctccaagagc tgaaggatga gctggcgac	1020
aacctgtaca ttgcccagct ggagctgcgc aaccgggctg ccacgaaga aatgctcgcc	1080
tcgctgcccc cgagtggtg caacatcgac atcctggtea acaacgcggg tctggccctc	1140
ggcatggaac cggcgacaaa ggccagcgtc gaggactggg aaaccatgat cgacaccaac	1200
aacaagggac tcgtctacat gaccgcgcgt gtgctgcccc gcatggtega gcgcaaccac	1260
ggccacatca tcaacatcgg ctccaccgct ggagctggc cctacgctgg cggcaacgtc	1320
tatggcgca ccaaggcgtt cgtccgcag ttctccctga acctgcgcac cgacctgcac	1380
ggcaccgcgc tccgctgac cgacattgag cccggtctgg tcggcggcac cgagttcagc	1440
aacgtccgct tcaagggcga cgacggcaag gccgagaaaa cctaccagaa caccgtcgct	1500
ctgacccctg aggatgtcag cgaggcgtc tgggtggtea gcaactctgcc cgcgcacgtc	1560
aacatcaaca cctcagat gatgccgtc acgcagctct acgccggcct gaacgtccac	1620
cgccaatagg accgatggct gtgtagaagt actcgccgat agtggaaacc gacgccccag	1680
cactcgctcg agggcaaagg aatagagtag atgccgaccg cgggatccac ttaacgttac	1740
tgaatcatc aaacagcttg acgaatctgg atataagatc gttgggtgctg atgtcagctc	1800
cggagttgag acaaatgggtg ttcaggatct cgataagata cgttcatttg tccaagcagc	1860
aaagagtgcc ttctagtgat ttaatagtc catgtcaaca agaataaac gcgttttcgg	1920
gtttacctct tccagatata gctcatctgc aatgcattaa tgcattgact gcaacctagt	1980
aacgccttca ggctccggcg aagagaagaa tagcttagca gagctatctt cattttcggg	2040
agacgagatc aagcagatca acggtcgtca agagacctac gagactgagg aatccgctc	2099

<210> SEQ ID NO 96

<211> LENGTH: 47

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer to isolate Fragments 7 to 9

<400> SEQUENCE: 96

acaggctact tcttgtaagc tcgagtttct gtacagtgc cggtgac

47

-continued

```

<210> SEQ ID NO 97
<211> LENGTH: 2913
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: combination of SEQ ID NOS: 89 and 92

<400> SEQUENCE: 97

tctgacagac gggcaattga ttacgggac ccattggtaa cgaaatgtaa aagctaggag      60
atcgtecgcc gatgtcagga tgatttcact tgtttcttgt cgggtccacc ggtcaaagct      120
aaagaggagc aaaaggaacg gatagaatcg ggtgccgctg atctatacgg tatagtgcgc      180
ttatcacgtt gactcaaccc atgctattta actcaacccc tccttctgaa cccaccatc      240
ttcttctctt tcctctcatc ccacacaatt ctctatctca gatttgaatt ccaaagtc      300
tcggacgaaa ctgaacaagt ctctctccct tcgataaacc tttggtgatt ggaataactg      360
accatcttct atagtcccca aaccaaccga caatgtaaat acactcctcg attagccctc      420
tagagggcat acgatggaag tcattggaata cttttggctg gactctcaca atgatcaagg      480
tatcttaggt aacgtctttg gcgtgggccc gtgttcgttc ccagtcacg atgcattcac      540
atgcctctcc taagctgggc cctagactct aggatcctag tctagaagga catggcatcg      600
atggactggg ttcgttctga gattatacgg ctaaaacttg atctggataa taccagcgaa      660
aagggtcatg ccttctctcg ttcttctctg tgatggaatg gctaacagat gatagtcatt      720
gcaacttgaa acatgtctcc tcacagctgc atctacgaac ccactgtggc cgctaccggc      780
ctcaagggta aggtcgtggt ttctgagacc gtcccggttg agggagcttc tcagaccaag      840
ctgttggaac atttcggtgg caagtgggac gagttcaagt tcgcccctat ccgcgaaagc      900
caggctcttc gtgccatgac cagacgttac tttgaggacc tggacaagta cgctgaaagt      960
gacgttgta ttgttggtgc tggttcctgc ggtctgagca ctgcgtacgt cttggccaag      1020
gtcgtcccg accgaagat tgctatctgc gaggccagcg tctctcctgg tcagtagtcc      1080
atgatggatt gccttgcaat cagcttcccg gaactaacgt gcaatagggt gcggtgcctg      1140
gttgggtggc caactctttt ctgctatggt catgcgcgt cccgcggaag tcttctgaa      1200
cgagctgggt gttccttacg aagaggacgc aaacccaac tacgttgtcg tcaagcacgc      1260
ctcctgctt accctgacac tcattgtgaa ggttctctcc tccccaatg tcaagctctt      1320
caatgctacc gctgttgagg acttgatcac ccgtccgacc gagaacggca accccagat      1380
tgctgggtgt gtcgtcaact ggacgctggt cacccttcac cagcatgac actcctgcat      1440
ggacccaac actatcaacg ctctgtcat catcagtacc actggtcacg atgggccatt      1500
cggcgcttc tgtcggaagc gcttggtgtc catgggcagc gtcgacaagc taggtggcat      1560
gcgtggtctc gacatgaact cggccgagga tgccatcgtc aagaacaccc gcgaggttac      1620
taagggtctg ataatcgcg gtatggagct gtctgaaatt gatggcttta accgcatggg      1680
ccctaccttc ggtgccatgg ttctcagtgg tgtcaaggct gccgaggagg cattgaagg      1740
gttcgacgag cgtcagcgcg agtggtctga gtaaatgact cactaccga atgggttcag      1800
tgcatgaacc ggatttgtct tacggtcttt gacgataggg gaatgatgat tatgtgatg      1860
ttctgagatt tgaatgaact cgttagctcg taatccacat gcatatgtaa atggctgtgt      1920
cccgtatgta acggtggggc attctagaat aattatgtgt aacaagaaag acagtataat      1980

```

-continued

```

acaaacaaag atgcaagagc ggctctcag caacattcgc catgttcacg tacagctttc 2040
aacggcctcg aacagtcact gtggatggat accagaggag agacccatca gttcaatcgc 2100
agggcagatg agtgtcgcat acattctcgc cgtccagctg gtcgaccagc aatgtctttt 2160
gtcccagttt tctgagtttg atgacaacct ggagaggcca gaagtttggg atctggccag 2220
gaaggttact tcattctaaa gcgaagagtt tgatcaagac ggcaactgtc tcagtgcggg 2280
tcgcgtgagg attgagtcca acgatggttc ttctattacg gaaagtgtcg agaagcctct 2340
tggtgtcaaa gagcccatgc caaacgaacg gattctccac aaataccgaa ccttgctgg 2400
tagcgtgacg gacgaatccc gggtgaaaga gattgaggat cttgtcctcg gcctggacag 2460
gctcaccgac attagcccat tgctggagct gctgaattgc cccgtgaaat cgccactggt 2520
ataaatggga agcगतatgg aaacatttca tgtcacgggc acaaattcta ggcatatcg 2580
tacctggatg gtgaaaccac cagcggttta gcagatagaa gatagaactcc ttctgctctg 2640
cgttgcgtct tgaatttagt tcgttccactg gcttaagaac ttagaatgca atacagtctc 2700
tcttattttt tattaaaac acgtattccc acattcggcg actggaggat acgaaagcag 2760
tgttggtggt gctccccgta atggatatga ttttgtgac tggactattc tatgaccatt 2820
ccctccaacg gagatccttt ctgcacactt tagatgttga cgctgtctgg aggaactact 2880
tttgcgtgc aaagactatg agcagtggag ctg 2913

```

```

<210> SEQ ID NO 98
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate Fragments 11-12

```

```

<400> SEQUENCE: 98

```

```

actaaaggga acaaaagctg gagctcagct ccaactgctca tagtctttg 49

```

We claim:

1. An isolated recombinant *Aspergillus* fungus comprising a genetic inactivation of an endogenous cis-aconitic acid decarboxylase (cadA) gene.

2. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the *Aspergillus* fungus is *Aspergillus pseudoterreus*.

3. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the *Aspergillus* fungus is *Aspergillus terreus*.

4. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the endogenous cadA gene is genetically inactivated by complete or partial deletion mutation or by insertional mutation.

5. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the cadA gene prior to its genetic inactivation encodes a protein having at least 80% sequence identity to SEQ ID NO: 50 or 52.

6. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the cadA gene prior to its genetic inactivation

comprises a coding sequence comprising at least 80% sequence identity to SEQ ID NO: 49, 51, 59 or 92.

7. A composition comprising the isolated recombinant *Aspergillus* fungus of claim 1.

8. A kit, comprising:

the isolated recombinant *Aspergillus* fungus of claim 1; and
a medium for culturing the fungus.

9. A method of making aconitic acid, comprising:

culturing the isolated recombinant *Aspergillus* fungus of claim 1 under conditions that permit the fungus to make aconitic acid; thereby making aconitic acid.

10. The method of claim 9, wherein the fungus is cultured in Riscaldati medium.

11. The method of claim 9, further comprising isolating the aconitic acid from culture media or from the fungus.

12. The method of claim 9, wherein the aconitic acid is cis-aconitic acid.

* * * * *